



Fungal Planet description sheets: 1284–1382

P.W. Crous^{1,2}, E.R. Osieck³, Ž. Jurjević⁴, J. Boers⁵, A.L. van Iperen¹, M. Starink-Willemse¹, B. Dima⁶, S. Balashov⁴, T.S. Bulgakov⁷, P.R. Johnston⁸, O.V. Morozova⁹, U. Pinruan¹⁰, S. Sommai¹⁰, P. Alvarado¹¹, C.A. Decock¹², T. Lebel¹³, S. McMullan-Fisher¹⁴, G. Moreno¹⁵, R.G. Shivas¹⁶, L. Zhao¹, J. Abdollahzadeh¹⁷, M. Abrinbana¹⁸, D.V. Ageev¹⁹, G. Akhmetova⁶, A.V. Alexandrova²⁰, A. Altés¹⁵, A.G.G. Amaral²¹, C. Angelini^{22,23}, V. Antonín²³, F. Arenas²⁴, P. Asselman²⁵, F. Badali¹⁸, A. Baghela^{26,27}, A. Bañares²⁸, R.W. Barreto²⁹, I.G. Baseia³⁰, J.-M. Bellanger³¹, A. Berraf-Tebbal³², A.Yu. Biketova^{33,34}, N.V. Bukharova³⁵, T.I. Burgess³⁶, J. Cabero³⁷, M.P.S. Câmara²¹, J.F. Cano-Lira³⁸, P. Ceryngier³⁹, R. Chávez⁴⁰, D.A. Cowan⁴¹, A.F. de Lima²¹, R.L. Oliveira⁴², S. Denman⁴³, Q.N. Dang⁴⁴, F. Dovana⁴⁵, I.G. Duarte²¹, A. Eichmeier³², A. Erhard⁴, F. Esteve-Raventós¹⁵, A. Fellin⁴⁶, G. Ferisin⁴⁷, R.J. Ferreira⁴⁸, A. Ferrer⁴⁹, P. Finy⁵⁰, E. Gaya⁵¹, A.D.W. Geering⁵², C. Gil-Durán⁴⁰, K. Glässnerová⁵³, A.M. Glushakova^{20,54}, D. Gramaje⁵⁵, F.E. Guard⁵⁶, A.L. Guarnizo²⁴, D. Haelewaters^{25,57}, R.E. Halling⁵⁸, R. Hill⁵¹, Y. Hirooka⁵⁹, V. Hubka^{53,60}, V.A. Iliushin⁹, D.D. Ivanova⁶¹, N.E. Ivanushkina⁶², P. Jangsantear⁶³, A. Justo⁶⁴, A.V. Kachalkin^{20,62}, S. Kato⁵⁹, P. Khamsuntorn⁶⁵, I.Y. Kirtsideli⁹, D.G. Knapp⁶, G.A. Kochkina⁶², O. Koukol⁶⁶, G.M. Kovács⁶, J. Kruse⁶⁷, T.K.A. Kumar⁶⁸, I. Kušan⁶⁹, T. Læssøe⁷⁰, E. Larsson⁷¹, R. Lebeuf⁷², G. Levicán⁴⁰, M. Loizides⁷³, P. Marinho⁷⁴, J.J. Luangsa-ard¹⁰, E.G. Lukina⁷⁵, V. Magaña-Dueñas³⁸, G. Maggs-Kölling⁷⁶, E.F. Malysheva⁹, V.F. Malysheva⁹, B. Martín⁷⁷, M.P. Martín⁷⁸, N. Matočec⁶⁹, A.R. McTaggart⁷⁹, M. Mehrabi-Koushki^{80,81}, A. Mešić⁶⁹, A.N. Miller⁸², P. Mironova²⁵, P.-A. Moreau⁸³, A. Morte²⁴, K. Müller⁸⁴, L.G. Nagy³³, S. Nanu⁶⁸, A. Navarro-Ródenas²⁴, W.J. Nel², T.H. Nguyen⁴⁴, T.F. Nóbrega²⁹, M.E. Noordeloos⁸⁵, I. Olariaga⁸⁶, B.E. Overton⁸⁷, S.M. Ozerskaya⁶², P. Palani⁸⁸, F. Pancorbo⁸⁹, V. Papp⁹⁰, J. Pawłowska⁹¹, T.Q. Pham⁴⁴, C. Phosri⁹², E.S. Popov⁹, A. Portugal^{93,94}, A. Pošta⁶⁹, K. Reschke⁹⁵, M. Reul⁹⁶, G.M. Ricci⁸⁷, A. Rodríguez²⁴, J. Romanowski³⁹, N. Ruchikachorn⁹⁷, I. Saar⁹⁸, A. Safi⁸⁰, B. Sakolrak⁶³, F. Salzmann⁹⁹, M. Sandoval-Denis¹, E. Sangwichein¹⁰⁰, L. Sanhueza⁴⁹, T. Sato¹⁰¹, A. Sastoque³⁸, B. Senn-Irlet¹⁰², A. Shibata⁵⁹, K. Siepe¹⁰³, S. Somrithipol¹⁰, M. Spetik³², P. Sridhar⁸⁸, A.M. Stchigel³⁸, K. Stuskova³², N. Suwannasai¹⁰⁴, Y.P. Tan¹⁰⁵, R. Thangavel¹⁰⁶, I. Tiago⁹³, S. Tiwari^{26,27}, Z. Tkalčec⁶⁹, M.A. Tomashevskaya⁶², C. Tonegawa⁵⁹, H.X. Tran⁴⁴, N.T. Tran⁵², J. Trovão⁹³, V.E. Trubitsyn⁶², J. Van Wyk¹⁰⁷, W.A.S. Vieira²¹, J. Vila¹⁰⁸, C.M. Visagie², A. Vizzini¹⁰⁹, S.V. Volobuev⁹, D.T. Vu¹¹⁰, N. Wangsawat¹¹¹, T. Yaguchi⁶⁰, E. Ercole¹¹², B.W. Ferreira²⁹, A.P. de Souza¹¹³, B.S. Vieira¹¹³, J.Z. Groenewald¹

Key words

ITS nrDNA barcodes
LSU
new taxa
systematics

Abstract Novel species of fungi described in this study include those from various countries as follows: **Antartica**, *Cladosporium austrolitorale* from coastal sea sand. **Australia**, *Austroboletus yourkae* on soil, *Crepidotus innuopurpureus* on dead wood, *Curvularia stenotaphri* from roots and leaves of *Stenotaphrum secundatum* and *Thecaphora stajscii* from capsules of *Oxalis radicata*. **Belgium**, *Paraxerochrysium coryli* (incl. *Paraxerochrysium* gen. nov.) from *Corylus avellana*. **Brazil**, *Calvatia nordestina* on soil, *Didymella tabebuicola* from leaf spots on *Tabebuia aurea*, *Fusarium subflagellisporum* from hypertrophied floral and vegetative branches of *Mangifera indica* and *Microdochium maculosum* from living leaves of *Digitaria insularis*. **Canada**, *Cuphophyllus bondii* from a grassland. **Croatia**, *Mollisia inferiseptata* from a rotten *Laurus nobilis* trunk. **Cyprus**, *Amanita exilis* on calcareous soil. **Czech Republic**, *Cytospora hippophaicola* from wood of symptomatic *Vaccinium corymbosum*. **Denmark**, *Lasiosphaeria deviata* on pieces of wood and herbaceous debris. **Dominican Republic**, *Calocybella goethei* among grass on a lawn. **France (Corsica)**, *Inocybe corsica* on wet ground. **France (French Guiana)**, *Trechispora patawaensis* on decayed branch of unknown angiosperm tree and *Trechispora subregularis* on decayed log of unknown angiosperm tree. **Germany**, *Paramicrothecium sambuci* (incl. *Paramicrothecium* gen. nov.) on dead stems of *Sambucus nigra*. **India**, *Aureobasidium microtermis* from the gut of a *Microtermes* sp. termite, *Laccaria diospyricola* on soil and *Phylloporia tamilnadensis* on branches of *Catunaregam spinosa*. **Iran**, *Pythium serotinoosporum* from soil under *Prunus dulcis*. **Italy**, *Pluteus brunneovenosus* on twigs of broadleaved trees on the ground. **Japan**, *Heterophoma rehmanniae* on leaves of *Rehmannia glutinosa* f. *hueichingensis*. **Kazakhstan**, *Murisporea kazachstanica* from healthy roots of *Triticum aestivum*. **Namibia**, *Caespitomonium euphorbiae* (incl. *Caespitomonium* gen. nov.) from

Abstract (cont.)

stems of an *Euphorbia* sp. **Netherlands**, *Alfaria junci*, *Myrmecridium junci*, *Myrmecridium juncicola*, *Myrmecridium juncigenum*, *Ophioceras junci*, *Paradinemasporium junci* (incl. *Paradinemasporium* gen. nov.), *Phialoseptomonium junci*, *Sporidesmiella juncicola*, *Xenopyricularia junci* and *Zaenenomyces quadripartis* (incl. *Zaenenomyces* gen. nov.), from dead culms of *Juncus effusus*, *Cylindromonium everniae* and *Rhodoveronaea everniae* from *Evernia prunastri*, *Cyphellophora sambuci* and *Myrmecridium sambuci* from *Sambucus nigra*, *Kiflimonium junci*, *Sarocladium junci*, *Zaenenomyces moderatricis-academiae* and *Zaenenomyces versatilis* from dead culms of *Juncus inflexus*, *Microcera physciae* from *Physcia tenella*, *Myrmecridium dactylidis* from dead culms of *Dactylis glomerata*, *Neochalara spiraeae* and *Sporidesmium spiraeae* from leaves of *Spiraea japonica*, *Neofabraea salicina* from *Salix* sp., *Paradissoconium narthecii* (incl. *Paradissoconium* gen. nov.) from dead leaves of *Narthecium ossifragum*, *Polyscytalum vaccinii* from *Vaccinium myrtillus*, *Pseudosoloacrosporiella cryptomeriae* (incl. *Pseudosoloacrosporiella* gen. nov.) from leaves of *Cryptomeria japonica*, *Ramularia pararhabdospora* from *Plantago lanceolata*, *Sporidesmiella pini* from needles of *Pinus sylvestris* and *Xenoacrodontium juglandis* (incl. *Xenoacrodontium* gen. nov. and *Xenoacrodontiaceae* fam. nov.) from *Juglans regia*. **New Zealand**, *Cryptometrion metrosideri* from twigs of *Metrosideros* sp., *Coccomyces pycnophyllocladi* from dead leaves of *Phyllocladus alpinus*, *Hypoderma aliforme* from fallen leaves *Fuscopora solandri* and *Hypoderma subiculatum* from dead leaves *Phormium tenax*. **Norway**, *Neodevriesia kalakoutskii* from permafrost and *Variabilispora viridis* from driftwood of *Picea abies*. **Portugal**, *Entomortierella hereditatis* from a biofilm covering a deteriorated limestone wall. **Russia**, *Colpoma junipericola* from needles of *Juniperus sabina*, *Entoloma cinnamomeum* on soil in grasslands, *Entoloma verae* on soil in grasslands, *Hyphodermella pallidostraminea* on a dry dead branch of *Actinidia* sp., *Lepiota sayanensis* on litter in a mixed forest, *Papiliotrema horticola* from *Malus communis*, *Paramacroventuria ribis* (incl. *Paramacroventuria* gen. nov.) from leaves of *Ribes aureum* and *Paramyrothecium lathyri* from leaves of *Lathyrus tuberosus*. **South Africa**, *Harzia combreti* from leaf litter of *Combretum collinum* ssp. *sulvense*, *Penicillium xyleborini* from *Xyleborinus saxesenii*, *Phaeoisaria dalbergiae* from bark of *Dalbergia armata*, *Protocreopsis euphorbiae* from leaf litter of *Euphorbia ingens* and *Roi-giella syzygii* from twigs of *Syzygium chordatum*. **Spain**, *Genea zamorana* on sandy soil, *Gymnopus nigrescens* on *Scleropodium touretii*, *Hesperomyces parexochomi* on *Parexochomus quadriplagiatus*, *Paraphoma variabilis* from dung, *Phaeococcomyces kinklidomatophilus* from a blackened metal railing of an industrial warehouse and *Tuber suaveolens* in soil under *Quercus faginea*. **Svalbard and Jan Mayen**, *Inocybe nivea* associated with *Salix polaris*. **Thailand**, *Biscogniauxia whalleyi* on corticated wood. **UK**, *Parasitella quercicola* from *Quercus robur*. **USA**, *Aspergillus arizonicus* from indoor air in a hospital, *Caeliomyces tampanus* (incl. *Caeliomyces* gen. nov.) from office dust, *Cippumomyces mortalis* (incl. *Cippumomyces* gen. nov.) from a tombstone, *Cylindrium desperesense* from air in a store, *Tetracoccosporium pseudoaerium* from air sample in house, *Toxicocladosporium glendoranum* from air in a brick room, *Toxicocladosporium losalamitosense* from air in a classroom, *Valsonectria portsmouthensis* from air in men's locker room and *Varicosporellopsis americana* from sludge in a water reservoir. **Vietnam**, *Entoloma kovalenkoi* on rotten wood, *Fusarium chuoi* inside seed of *Musa itinerans*, *Microspallia albobellina* on soil in tropical evergreen mixed forests and *Phytophthora docyniae* from soil and roots of *Docynia indica*. Morphological and culture characteristics are supported by DNA barcodes.

Citation: Crous PW, Osieck ER, Jurjević Ž, et al. 2021. Fungal Planet description sheets: 1284–1382. Persoonia 47: 178–374.

<https://doi.org/10.3767/persoonia.2021.47.06>.

Effectively published online: 24 December 2021 [Received: 1 October 2021; Accepted: 4 November 2021].

¹ Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands.

² Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa.

³ Jkvr. C.M. van Asch van Wijcklaan 19, 3972 ST Driebergen-Rijsenburg, Netherlands.

⁴ EMSL Analytical, Inc., 200 Route 130 North, Cinnaminson, NJ 08077 USA.

⁵ Conventstraat 13A, 6701 GA Wageningen, Netherlands.

⁶ Department of Plant Anatomy, Institute of Biology, Eötvös Loránd University, Pázmány Péter sétány 1/C, H-1117, Budapest, Hungary.

⁷ Department of Plant Protection, Federal Research Centre the Subtropical Scientific Centre of the Russian Academy of Sciences, Yana Fabritsiusa street 2/28, 354002 Sochi, Krasnodar region, Russia.

⁸ Manaaki Whenua – Landcare Research, P. Bag 92170, Auckland 1142, New Zealand.

⁹ Komarov Botanical Institute of the Russian Academy of Sciences, 197376, 2 Prof. Popov Str., Saint Petersburg, Russia.

¹⁰ Plant Microbe Interaction Research Team (APMT), BIOTEC, National Science and Technology Development Agency, Pathum Thani, Thailand, 113 Thailand Science Park, Phahonyothin Rd., Khlong Nueng, Khlong Luang, Pathum Thani Thailand.

¹¹ ALVALAB, C/ Dr. Fernando Bongera, Severo Ochoa bldg. S1.04, 33006 Oviedo, Spain.

¹² Mycothèque de l'Université catholique de Louvain (MUCL, BCCMTM), Earth and Life Institute – ELIM – Mycology, Université catholique de Louvain, Croix du Sud 2 bte L7.05.06, B-1348 Louvain-la-Neuve, Belgium.

¹³ State Herbarium of South Australia, Adelaide, South Australia 5000 Australia.

¹⁴ Royal Botanic Gardens, Victoria, Melbourne, Victoria, Australia.

¹⁵ Universidad de Alcalá, Facultad de Ciencias, Departamento de Ciencias de la Vida (Botánica), 28805 Alcalá de Henares, Madrid, Spain.

¹⁶ Centre for Crop Health, University of Southern Queensland, Toowoomba 4350, Queensland, Australia.

¹⁷ Department of Plant Protection, Agriculture Faculty, University of Kurdistan, P.O. Box 416, Sanandaj, Iran.

¹⁸ Department of Plant Protection, Faculty of Agriculture, Urmia University, P.O. Box 165, Urmia, Iran.

¹⁹ LLC 'Signatec', 630090, Inzhenernaya Str. 22, Novosibirsk, Russia.

²⁰ Lomonosov Moscow State University (MSU), 119234, 1, 12 Leninskie Gory Str., Moscow, Russia.

²¹ Departamento de Agronomia, Universidade Federal Rural de Pernambuco, Recife, Pernambuco, Brazil.

²² Herbario Jardín Botánico Nacional Dr. Rafael Ma. Moscoso, Santo Domingo, Dominican Republic and Via Cappuccini, 78/8 – 33170 Pordenone, Italy.

²³ Department of Botany, Moravian Museum, Zelný trh 6, 659 37 Brno, Czech Republic.

²⁴ Departamento de Biología Vegetal (Botánica), Facultad de Biología, Universidad de Murcia, 30100 Murcia, Spain.

²⁵ Research Group Mycology, Department of Biology, Ghent University, K.L. Ledeganckstraat 35, 9000 Gent, Belgium.

²⁶ National Fungal Culture Collection of India (NFCCI).

²⁷ Biodiversity and Palaeobiology Group, MACS-Agharkar Research Institute, G.G. Agarkar Road, Pune 411004, Maharashtra, India.

²⁸ Departamento de Botánica, Ecología y Fisiología Vegetal, Universidad de La Laguna. Apdo. 456, E-38200 La Laguna, Tenerife, Islas Canarias, Spain.

²⁹ Departamento de Fitopatologia, Universidade Federal de Viçosa, Viçosa, 36570-900, MG, Brazil.

³⁰ Departamento Botânica e Zoologia, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Campus Universitário, 59072–970 Natal, RN, Brazil.

³¹ CEFÉ, CNRS, Université de Montpellier, Université Paul-Valéry Montpellier 3, EPHE, IRD, INSERM, 1919 route de Mende, F-34293 Montpellier Cedex 5, France.

- ³² Mendeleum – Institute of Genetics, Faculty of Horticulture, Mendel University in Brno, Valtická 334, Lednice, 69144, Czech Republic.
- ³³ Institute of Biochemistry, Biological Research Centre of the Eötvös Lóránd Research Network, Temesvári Blvd. 62, H-6726 Szeged, Hungary.
- ³⁴ Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3DS, UK.
- ³⁵ Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far Eastern Branch of the Russian Academy of Sciences, Pr-t 100-let Vladivostoka 159, 690022 Vladivostok, Russia.
- ³⁶ Phytophthora Science and Management, Harry Butler Institute, Murdoch University, 90 South Street, Murdoch, WA 6150, Australia.
- ³⁷ C/ El Sol 6, 49800 Toro, Zamora, Spain.
- ³⁸ Mycology Unit, Medical School, Universitat Rovira i Virgili (URV), Sant Llorenç 21, 43201 Reus, Tarragona, Spain.
- ³⁹ Institute of Biological Sciences, Cardinal Stefan Wyszyński University, Wóycickiego 1/3, 01-938 Warsaw, Poland.
- ⁴⁰ Facultad de Química y Biología, Universidad de Santiago de Chile (USACH), Alameda 3363, Estación Central, 9170022, Santiago, Chile.
- ⁴¹ Centre for Microbial Ecology and Genomics, Department of Biochemistry, Genetics and Microbiology, University of Pretoria, Private Bag X20, Hatfield 0028, Pretoria, South Africa.
- ⁴² Programa de Pós-Graduação em Sistemática e Evolução, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Av. Senador Salgado Filho, 3000, 59072-970 Natal, RN, Brazil.
- ⁴³ Forest Research, Alice Holt Lodge, Farnham, Surrey, UK.
- ⁴⁴ Forest Protection Research Centre, Vietnamese Academy of Forest Sciences, 46 Duc Thang Ward, Bac Tu Liem District, Hanoi City, Vietnam.
- ⁴⁵ Via Quargnento, 17, 15029, Solero (AL), Italy.
- ⁴⁶ Via G. Canestrini 10/B, I-38028, Novella (TN), Italy.
- ⁴⁷ Associazione Micologica Bassa Friulana, 33052 Cervignano del Friuli, Italy.
- ⁴⁸ Programa de Pós-Graduação em Biologia de Fungos, Departamento de Micologia, Universidade Federal de Pernambuco, 50670-420 Recife, PE, Brazil.
- ⁴⁹ Facultad de Estudios Interdisciplinarios, Núcleo de Química y Bioquímica, Universidad Mayor, Santiago, Chile.
- ⁵⁰ Zsombolyai u. 56, 8000 Székesfehérvár, Hungary.
- ⁵¹ Comparative Fungal Biology, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3DS, UK.
- ⁵² Centre for Horticultural Science, Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Dutton Park 4102, Queensland, Australia.
- ⁵³ Department of Botany, Faculty of Science, Charles University, Benátská 2, 128 01 Prague 2, Czech Republic.
- ⁵⁴ Mechnikov Research Institute for Vaccines and Sera, 105064, Moscow, Maly Kazenny by-street, 5A, Russia.
- ⁵⁵ Instituto de Ciencias de la Vid y del Vino (ICVV), Consejo Superior de Investigaciones Científicas (CSIC) – Universidad de La Rioja – Gobierno de La Rioja, Ctra. LO-20, Salida 13, 26007, Logroño, Spain.
- ⁵⁶ Maleny, Queensland, Australia.
- ⁵⁷ Faculty of Science, University of South Bohemia, Branišovská 31, 370 05 České Budějovice, Czech Republic.
- ⁵⁸ Inst. Systematic Botany, New York Botanical Garden, 2900 Southern Blvd, Bronx, NY, USA 10458-5126.
- ⁵⁹ Department of Clinical Plant Science, Hosei University, 3-7-2 Kajino-cho, Koganei, Tokyo, Japan.
- ⁶⁰ Medical Mycology Research Center, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba, 260-8673, Japan.
- ⁶¹ The Herzen State Pedagogical University of Russia, 191186, 48 Moyka Embankment, Saint Petersburg, Russia.
- ⁶² All-Russian Collection of Microorganisms, G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Pushchino Center for Biological Research of the Russian Academy of Sciences, 142290, Pushchino, pr. Nauki, 5, Russia.
- ⁶³ Forest and Plant Conservation Research Office, Department of National Parks, Wildlife and Plant Conservation, Chatuchak District, Bangkok, Thailand.
- ⁶⁴ Department of Natural History, New Brunswick Museum, 277 Douglas Ave. Saint John, NB – E2K 1E5 Canada.
- ⁶⁵ Microbe Interaction and Ecology Laboratory (BMIE), National Center for Genetic Engineering and Biotechnology (BIOTEC), 113 Thailand Science Park, Phahonyothin Rd., Khlong Nueng, Khlong Luang, Pathum Thani Thailand.
- ⁶⁶ Department of Botany, Charles University, Faculty of Science, Benátská 2, 128 01 Prague 2, Czech Republic.
- ⁶⁷ Pfalzmuseum für Naturkunde – POLLICHIA-Museum, Hermann-Schäfer-Str. 17, 67098 Bad Dürkheim, Germany.
- ⁶⁸ Department of Botany, The Zamorin's Guruvayurappan College, Kozhikode, Kerala, India.
- ⁶⁹ Laboratory for Biological Diversity, Ruđer Bošković Institute, Bijenička cesta 54, HR-10000 Zagreb, Croatia.
- ⁷⁰ Globe Inst./Department of Biology, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen Ø, Denmark, Denmark.
- ⁷¹ Biological and Environmental Sciences, University of Gothenburg, and Gothenburg Global Biodiversity Centre, Box 461, SE40530 Göteborg, Sweden.
- ⁷² 775, rang du Rapide Nord, Saint-Casimir, Quebec, G0A 3L0, Canada.
- ⁷³ P.O. Box 58499, 3734 Limassol, Cyprus.
- ⁷⁴ Departamento de Biologia Celular e Genética, Universidade Federal do Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil.
- ⁷⁵ Saint Petersburg State University, 199034, 7-9 Universitetskaya emb., St. Petersburg, Russia.
- ⁷⁶ Gobabeb-Namib Research Institute, Walvis Bay, Namibia.
- ⁷⁷ Servicio Territorial de Agricultura, Ganadería y Desarrollo Rural de Zamora, C/ Prado Tuerto 17, 49019 Zamora, Spain.
- ⁷⁸ Real Jardín Botánico RJB-CSIC, Plaza de Murillo 2, 28014 Madrid, Spain.
- ⁷⁹ Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Brisbane 4001, Australia.
- ⁸⁰ Department of Plant Protection, Faculty of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Khuzestan Province, Iran.
- ⁸¹ Biotechnology and Bioscience Research Center, Shahid Chamran University of Ahvaz, Ahvaz, Iran.
- ⁸² University of Illinois Urbana-Champaign, Illinois Natural History Survey, 1816 South Oak Street, Champaign, Illinois, 61820, USA.
- ⁸³ Université de Lille, Faculté de pharmacie de Lille, EA 4483, F-59000 Lille, France.
- ⁸⁴ Falkstraße 103, D-47058 Duisburg, Germany.
- ⁸⁵ Naturalis Biodiversity Center, section Botany, P.O. Box 9517, 2300 RA Leiden, The Netherlands.
- ⁸⁶ Rey Juan Carlos University, Dep. Biology and Geology, Physics and Inorganic Chemistry, C/ Tulipán s/n, 28933 Móstoles, Madrid, Spain.
- ⁸⁷ 205 East Campus Science Center, Lock Haven University, Department of Biology, Lock Haven, PA 17745, USA.
- ⁸⁸ Centre for Advanced Studies in Botany, University of Madras, Guindy Campus, Chennai 600 025, India.
- ⁸⁹ Sociedad Micológica de Madrid, Real Jardín Botánico, C/ Claudio Moyano 1, 28014 Madrid, Spain.
- ⁹⁰ Department of Botany, Hungarian University of Agriculture and Life Sciences, Ménézi út 44. H-1118 Budapest, Hungary.
- ⁹¹ Institute of Evolutionary Biology, Faculty of Biology, Biological and Chemical Research Centre, University of Warsaw, ul. Zwirki i Wigury 101, 02-089 Warsaw, Poland.
- ⁹² Biology programme, Faculty of Science, Nakhon Phanom University, Nakhon Phanom, 48000, Thailand.
- ⁹³ Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, 3004-531 Coimbra, Portugal.
- ⁹⁴ Fitolab - Laboratory for Phytopathology, Instituto Pedro Nunes, 3030-199 Coimbra, Portugal.
- ⁹⁵ Mycology Research Group, Faculty of Biological Sciences, Goethe University Frankfurt am Main, Max-von-Laue Straße 13, 60439 Frankfurt am Main, Germany.
- ⁹⁶ Ostenstraße 19, D-95615 Marktredwitz, Germany.
- ⁹⁷ The Institute for the Promotion of Teaching Science and Technology, Bangkok, 10110, Thailand.
- ⁹⁸ Institute of Ecology and Earth Sciences, University of Tartu, Ravila Street 14A, 50411 Tartu, Estonia.
- ⁹⁹ Kloosterweg 5, 6301WK, Valkenburg a/d Geul, The Netherlands.
- ¹⁰⁰ Department of Biology, Faculty of Science, Ramkhamhaeng University, Bangkok, 10240, Thailand.
- ¹⁰¹ Department of Agro-Food Science, Niigata Agro-Food University, 2416 Hiranedai, Tainai, Niigata Prefecture, Japan.
- ¹⁰² Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland.
- ¹⁰³ Geeste 133, D-46342 Velen, Germany.
- ¹⁰⁴ Department of Microbiology, Faculty of Science, Srinakharinwirot University, Bangkok, 10110 Thailand.
- ¹⁰⁵ Plant Pathology Herbarium, Department of Agriculture and Fisheries, Dutton Park 4102, Queensland, Australia.
- ¹⁰⁶ Plant Health and Environment Laboratory, Ministry for Primary Industries, P.O. Box 2095, Auckland 1140, New Zealand.
- ¹⁰⁷ Department of Plant Soil and Microbial Sciences, 1066 Bogue Street, Michigan State University, East Lansing, MI, 48824 USA.
- ¹⁰⁸ Passatge del Torn, 4, 17800 Olot, Spain.
- ¹⁰⁹ Department of Life Sciences and Systems Biology, University of Torino, Viale P.A. Mattioli 25, I-10125 Torino, Italy.
- ¹¹⁰ Research Planning and International Cooperation Department, Plant Resources Center, An Khanh, Hoai Duc, Hanoi 152900, Vietnam.
- ¹¹¹ Department of Biology, Faculty of Science, Srinakharinwirot University, Bangkok, 10110 Thailand.
- ¹¹² Via Murazzano 11, I-10141, Torino (TO), Italy.
- ¹¹³ Laboratório de Microbiologia e Fitopatologia, Universidade Federal de Uberlândia, Monte Carmelo, 38500-000, MG, Brazil.

Acknowledgements P.R. Johnston thanks J. Sullivan (Lincoln University) for the habitat image of Kowai Bush, Duckchul Park (Manaaki Whenua – Landcare Research) for the DNA sequencing, and the New Zealand Department of Conservation for permission to collect the specimens; this research was supported through the Manaaki Whenua – Landcare Research Biota Portfolio with funding from the Science and Innovation Group of the New Zealand Ministry of Business, Innovation and Employment. V. Hubka was supported by the Czech Ministry of Health (grant number NU21-05-00681), and is grateful for the support from the Japan Society for the Promotion of Science – grant-in-aid for JSPS research fellow (grant no. 20F20772). K. Glässnerová was supported by the Charles University Grant Agency (grant No. GAUK 140520). J. Tróvão and colleagues were financed by FEDER-Fundo Europeu de Desenvolvimento Regional funds through the COMPETE 2020 – Operational Programme for Competitiveness and Internationalisation (POCI), and by Portuguese funds through FCT – Fundação para a Ciência e a Tecnologia in the framework of the project POCI-01-0145-FEDER-PTDC/EPH-PAT/3345/2014. This work was carried out at the R&D Unit Centre for Functional Ecology – Science for People and the Planet (CFE), with reference UIDB/04004/2020, financed by FCT/MCTES through national funds (PIDDAC). J. Tróvão was also supported by POCH – Programa Operacional Capital Humano (co-funding by the European Social Fund and national funding by MCTES), through a ‘FCT – Fundação para a Ciência e Tecnologia’ PhD research grant (SFRH/BD/132523/2017). D. Haelewaters acknowledges support from the Research Foundation – Flanders (Junior Postdoctoral Fellowship 1206620N). M. Loizides and colleagues are grateful to Y. Cherniavsky for contributing collections AB A12-058-1 and AB A12-058-2, and Á. Kovács and B. Kiss for their help with molecular studies of these specimens. C. Zmuda is thanked for assisting with the collection of ladybird specimens infected with *Hesperomyces parexochomi*. A.V. Kachalikin and colleagues were supported by the Russian Science Foundation (grant No. 19-74-10002). The study of A.M. Glushakova was carried out as part of the Scientific Project of the State Order of the Government of Russian Federation to Lomonosov Moscow State University No. 121040800174-6. S. Nanu acknowledges the Kerala State Council for Science, Technology and Environment (KSCSTE) for granting a research fellowship and is grateful to the Chief Conservator of Forests and Wildlife for giving permission to collect fungal samples. A. Bafiães and colleagues thank L. Monje and A. Pueblos of the Department of Drawing and Scientific Photography at the University of Alcalá for their help in the digital preparation of the photographs, and J. Rejos, curator of the AH herbarium for his assistance with the specimens examined in the present study. The research of V. Antonín received institutional support for long-term conceptual development of research institutions provided by the Ministry of Culture (Moravian Museum, ref. MK000094862). The studies of E.F. Malysheva, V.F. Malysheva, O.V. Morozova, and S.V. Volobuev were carried out within the framework of a research project of the Komarov Botanical Institute RAS, St Petersburg, Russia (AAAA-A18-118022090078-2) using equipment of its Core Facility Centre ‘Cell and Molecular Technologies in Plant Science’. The study of A.V. Alexandrova was carried out as part of the Scientific Project of the State Order of the Government of Russian Federation to Lomonosov Moscow State University No. 121032300081-7. The Kits van Waveren Foundation (Rijks-herbariumfonds Dr E. Kits van Waveren, Leiden, Netherlands) contributed substantially to the costs of sequencing and travelling expenses for M.E. Noordeloos. The work of B. Dima was partly supported by the ÚNKP-20-4 New National Excellence Program of the Ministry for Innovation and Technology from the source of the National Research, Development and Innovation Fund. The work of L. Nagy was supported by the ‘Momentum’ program of the Hungarian Academy of Sciences (contract No. LP2019-13/2019 to L.G.N.). G.A. Kochkina and colleagues acknowledge N. Demidov for the background photograph, and N. Suzina for the SEM photomicrograph. The research of C.M. Visagie and W.J. Nel was supported by the National Research Foundation grant no 118924 and SFH170610239162. C. Gil-Durán acknowledges Agencia Nacional de Investigación y Desarrollo, Ministerio de Ciencia, Tecnología, Conocimiento e Innovación, Gobierno de Chile, for grant ANID – Fondecyt de Postdoctorado 2021 – N° 3210135. R. Chávez and G. Levicán thank DICYT-USACH and acknowledges the grants INACH RG_03-14 and INACH RT_31-16 from the Chilean Antarctic Institute, respectively. S. Tiwari and A. Baghela would like to acknowledge R. Avchar and K. Balasubramanian from the Agharkar Research Institute, Pune, Maharashtra for helping with the termite collection. S. Tiwari is also thankful to the University Grants Commission, Delhi (India) for a junior research fellow-

ship (827/(CSIR-UGC NET DEC.2017)). R. Lebeuf and I. Saar thank D. and H. Spencer for collecting and photographing the holotype of *C. bondii*, and R. Smith for photographing the habitat. A. Voitek is thanked for helping with the colour plate and review of the manuscript, and the Foray Newfoundland and Labrador for providing the paratype material. I. Saar was supported by the Estonian Research Council (grant PRG1170) and the European Regional Development Fund (Centre of Excellence EcolChange). M.P.S. Câmara acknowledges the ‘Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq’ for the research productivity fellowship, and financial support (Universal number 408724/2018-8). W.A.S. Vieira acknowledges the ‘Coordenação de Aperfeiçoamento Pessoal de Ensino Superior – CAPES’ and the ‘Programa Nacional de Pós-Doutorado/CAPES – PNPD/CAPES’ for the postdoctoral fellowship. A.G.G. Amaral acknowledges CNPq, and A.F. Lima and I.G. Duarte acknowledge CAPES for the doctorate fellowships. F. Esteve-Raventós and colleagues were financially supported by FEDER/Ministerio de Ciencia, Innovación y Universidades – Agencia Estatal de Investigación (Spain)/ Project CGL2017-86540-P. The authors would like to thank L. Hugot and N. Suberbielle (Conservatoire Botanique National de Corse, Office de l’Environnement de la Corse, Corti) for their help. The research of E. Larsson is supported by The Swedish Taxonomy Initiative, SLU Artdatabanken, Uppsala. Financial support was provided to R.J. Ferreira by the National Council for Scientific and Technological Development (CNPq), and to I.G. Baseia, P.S.M. Lúcio and M.P. Martín by the National Council for Scientific and Technological Development (CNPq) under CNPq-Universal 2016 (409960/2016-0) and CNPq-visiting researcher (407474/2013-7). J. Cabero and colleagues wish to acknowledge A. Rodríguez for his help to describe *Genea zamorana*, as well as H. Hernández for sharing information about the vegetation of the type locality. S. McMullan-Fisher and colleagues acknowledge K. Syme (assistance with illustrations), J. Kellermann (translations), M. Barrett (collection, images and sequences), T. Lohmeyer (collection and images) and N. Karunajeewa (for prompt accessioning). This research was supported through funding from Australian Biological Resources Study grant (TTC217-06) to the Royal Botanic Gardens Victoria. The research of M. Spetik and co-authors was supported by project No. CZ.02.1.01/0.0/0.0/16_017/0002334. N. Wangsawat and colleagues were partially supported by NRCIT and the Royal Golden Jubilee Ph.D. programme, grant number PHD/0218/2559. They are thankful to M. Kamsook for the photograph of the Phu Khiao Wildlife Sanctuary and P. Thamvithayakorn for phylogenetic illustrations. The study by N.T. Tran and colleagues was funded by Hort Innovation (Grant TU19000). They also thank the turf growers who supported their surveys and specimen collection. N. Matočec, I. Kušan, A. Pošta, Z. Tkalčec and A. Mešić thank the Croatian Science Foundation for their financial support under the project grant HRZZ-IP-2018-01-1736 (ForFun-giDNA). A. Pošta thanks the Croatian Science Foundation for their support under the grant HRZZ-2018-09-7081. A. Morte is grateful to Fundación Séneca – Agencia de Ciencia y Tecnología de la Región de Murcia (20866/PI/18) for financial support. The research of G. Akhmetova, G.M. Kovács, B. Dima and D.G. Knapp was supported by the National Research, Development and Innovation Office, Hungary (NKFIH KH-130401 and K-139026), the ELTE Thematic Excellence Program 2020 supported by the National Research, Development and Innovation Office (TKP2020-IKA-05) and the Stipendium Hungaricum Programme. The support of the János Bolyai Research Scholarship of the Hungarian Academy of Sciences and the Bolyai+ New National Excellence Program of the Ministry for Innovation and Technology to D.G. Knapp is highly appreciated. F.E. Guard and colleagues are grateful to the traditional owners, the Jirral and Warungu people, as well as L. and P. Hales, Reserve Managers, of the Yourka Bush Heritage Reserve. Their generosity, guidance, and the opportunity to explore the Bush Heritage Reserve on the Einasleigh Uplands in far north Queensland is greatly appreciated. The National Science Foundation (USA) provided funds (DBI#1828479) to the New York Botanical Garden for a scanning electron microscope used for imaging the spores. V. Papp was supported by the ÚNKP-21-5 New National Excellence Program of the Ministry for Innovation and Technology from the National Research, Development and Innovation Fund of Hungary. A.N. Miller thanks the WM Keck Center at the University of Illinois Urbana – Champaign for sequencing *Lasiosphaeria devitata*. J. Pawłowska acknowledges support from National Science Centre, Poland (grant Opus 13 no 2017/25/B/NZ8/00473). The research of T.S. Bulgakov was carried out as part of the State Research Task of the Subtropical Scientific Centre of the Russian Academy of Sciences (Theme No. 0492-2021-0007). K. Bensch (Westerdijk Fungal Biodiversity Institute, Utrecht) is thanked for correcting the spelling of various Latin epithets.

Caespitomonium euphorbiae



Fungal Planet 1284 – 24 December 2021

Caespitomonium* Crous, gen. nov.Etymology.* Name refers to the tufted conidiophores and *Acremonium*.Classification — *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*.*Conidiophores* solitary to aggregated, branching extensively, aggregating, becoming tuft-like, arising from superficial mycelium, subcylindrical, hyaline, smooth. *Stipe* hyaline, smooth,subcylindrical, septate. Primary and secondary branches subcylindrical, hyaline, smooth. *Conidiogenous cells* phialidic, hyaline, smooth, terminal and intercalary, subcylindrical to narrowly ampulliform, with apical taper. *Conidia* in long unbranched chains, aseptate, hyaline, smooth, ellipsoid.*Type species.* *Caespitomonium euphorbiae* Crous
MycoBank MB 841776.***Caespitomonium euphorbiae* Crous, sp. nov.***Etymology.* Name refers to the host genus *Euphorbia* from which it was isolated.*Conidiophores* solitary to aggregated, penicillate, branching extensively, aggregating, becoming tuft-like, arising from superficial mycelium, subcylindrical, hyaline, smooth, 40–100 µm tall, 3–4 µm diam at base. *Stipe* hyaline, smooth, subcylindrical, septate, 15–30 × 3–4 µm. Primary and secondary branches subcylindrical, hyaline, smooth, aseptate, 8–12 × 2.5–3 µm. *Conidiogenous cells* phialidic, hyaline, smooth, terminal and intercalary, subcylindrical to narrowly ampulliform, with apical taper, 12–22 × 2.5–3 µm; apex with short, 1–1.5 µm long, non-flared collarette. *Conidia* in long unbranched chains, aseptate, hyaline, smooth, ellipsoid, apex subobtuse, base with truncate hilum, (5–)5.5–6(–6.5) × 2.5–3 µm.

Culture characteristics — Colonies flat, spreading, surface with concentric zone lines, with sparse to moderate aerial mycelium and smooth, lobate margin, reaching 40 mm diam after 2 wk at 25 °C. On malt extract agar (MEA), potato dextrose agar (PDA) and oatmeal agar (OA) surface and reverse buff.

Typus. NAMIBIA, Walvis Bay, on *Euphorbia* sp. (*Euphorbiaceae*), 20 Nov. 2019, P.W. Crous, HPC 3127 (holotype CBS H-24539, culture ex-type CPC 39083 = CBS 147075, ITS, LSU, *actA*, *rpb2*, *tef1* (second part) and *tub2* sequences GenBank OK664698.1, OK663737.1, OK651122.1, OK651157.1, OK651197.1 and OK651201.1, MycoBank MB 841777).***Caespitomonium squamicola* (Berk. & Broome) Crous, comb. nov.** MycoBank MB 841778*Basionym.* *Fusidium squamicola* Berk. & Broome, as '*squamicolum*', J. Linn. Soc., Bot. 14: 98. 1873 (1875).*Synonyms.* *Monosporium squamicola* (Berk. & Broome) Petch, as '*squamicolum*', Ann. R. bot. Gdns Peradeniya 9: 169. 1924.*Nalanthamala squamicola* (Berk. & Broome) W. Gams, Trans. Br. Mycol. Soc. 64: 402. 1975.*Material examined.* SRI LANKA, Peradeniya Gardens, from sooty mould and aphids on *Clerodendron monahassa*, collection date and collector unknown, CBS H-14970, culture CBS 701.73 = IMI 185382.***Caespitomonium hyalinulum* (Sacc.) Crous, comb. nov.**
MycoBank MB 841879*Basionym.* *Torula hyalinula* Sacc., *Michelia* 1 (no. 2): 265. 1878.*Synonyms.* *Oospora hyalinula* (Sacc.) Sacc., Fungi italica autogr. del. 17–28: t. 878. 1881.*Colour illustrations.* *Euphorbia virosa* in Namibia (photo Oliver Halsey). Conidiophores on synthetic nutrient-poor agar (SNA); conidiophores and conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.*Alysidium hyalinulum* (Sacc.) Pound & Clem., Minn. bot. Stud. 1 (Bulletin 9): 650. 1896.*Acremonium hyalinulum* (Sacc.) W. Gams, *Cephalosporium-artige Schimmelpilze* (Stuttgart): 104. 1971.*Culture examined.* USA, Maryland, Baltimore, air in hospital, date unknown, Walker No. 589, CBS H-8191, culture CBS 271.36.Notes — *Caespitomonium* represents a new genus in *Bionectriaceae* that is *acremonium*-like in morphology but distinguished from *Acremonium* s.str. based on its extensively branched, tuft-like conidiophores (Gams 1971, 1975).Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Acremonium polychromum* (strain BXT10-7, GenBank JX076941.1; Identities = 498/501 (99 %), no gaps), *Hydropisphaera erubescens* (strain 5141/HMAS 91779, GenBank FJ969800.1; Identities = 527/578 (91 %), 17 gaps (2 %)), and *Paracylindrocarpon aloicola* (strain CBS 141300, GenBank NR_154346.1; Identities = 530/583 (91 %), 17 gaps (2 %)). Closest hits using the **LSU** sequence are *Nalanthamala squamicola* (GenBank AF373281.1; Identities = 868/869 (99 %), no gaps), *Acremonium hyalinulum* (strain CBS 271.36, GenBank HQ232045.1; Identities = 846/848 (99 %), one gap (0 %)), and *Acremonium persicinum* (strain CBS 203.73, GenBank MH877816.1; Identities = 865/873 (99 %), four gaps (0 %)). Distant hits obtained using the **actA** sequence had highest similarity to *Volutella consors* (strain CBS 122767, GenBank KM231160.1; Identities = 384/400 (96 %), no gaps), *Gliocladiopsis irregularis* (strain CBS 755.97, GenBank KM231113.1; Identities = 384/400 (96 %), no gaps), and *Gliocladiopsis pseudotenuis* (strain CBS 116074, GenBank KM231112.1; Identities = 384/400 (96 %), no gaps). Distant hits obtained using the **rpb2** sequence had highest similarity to *Heleococcum aurantiacum* (strain CBS 201.35, GenBank JX158463.1; Identities = 708/845 (84 %), six gaps (0 %)), *Hydropisphaera peziza* (strain CBS 102038, GenBank DQ522444.1; Identities = 682/823 (83 %), eight gaps (0 %)), and *Gliomastix murorum* (as *Acremonium murorum*; strain AFTOL-ID 1393, GenBank FJ238363.1; Identities = 602/731 (82 %), six gaps (0 %)).

(Notes continues on Supplementary material page FP1284)

Supplementary material**FP1284** Phylogenetic tree.

Pedro W. Crous, Lin Zhao & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl, l.zhao@wi.knaw.nl & e.groenewald@wi.knaw.nl

Don Cowan, Centre for Microbial Ecology and Genomics, Department of Biochemistry, Genetics and Microbiology, University of Pretoria, Private Bag X20, Hatfield 0028, Pretoria, South Africa; e-mail: don.cowan@up.ac.za

Gillian Maggs-Kölling, Gobabeb-Namib Research Institute, Walvis Bay, Namibia; e-mail: gillian@gobabeb.org



Fungal Planet 1285 – 24 December 2021

Roigiella syzygii Crous, sp. nov.

Etymology. Name refers to the host genus *Syzygium* from which it was isolated.

Classification — *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*.

Mycelium consisting of hyaline, smooth, branched, septate, 2.5–3 µm diam hyphae. *Conidiophores* solitary, erect, arising from superficial hyphae, branched, septate, subcylindrical, up to 300 µm tall, 4–5 µm wide. *Conidiogenous cells* terminal and intercalary, subcylindrical, hyaline, smooth, 50–100 × 3–4 µm, with subdenticulate loci aggregated in dense, swollen clusters; denticles 1 µm diam, peg-like, not thickened nor darkened. *Conidia* in clusters, straight, obovoid, apex subobtuse, tapering to a truncate hilum, 1 µm diam, not constricted at median septum, enclosed in mucoid sheath, hyaline, smooth, granular, (16–)18–20(–22) × (4–)4.5–5 µm.

Culture characteristics — Colonies flat, spreading, with fluffy, moderate aerial mycelium and smooth, even margin, reaching 30 mm diam after 2 wk at 25 °C. On MEA, PDA and OA surface and reverse pale luteous.


Typus. SOUTH AFRICA, Northern Province, Nelspruit, Buffelskloof Nature Reserve, on twigs of *Syzygium cordatum* (Myrtaceae), Nov. 2018, P.W. Crous, HPC 3149 (holotype CBS H-24536, culture ex-type CPC 39044 = CBS 147072, ITS and LSU sequences GenBank OK664699.1 and OK663738.1, MycoBank MB 841779).

Notes — *Roigiella syzygii* was isolated from synnematal conidiophores on host tissue, but these structures did not develop in culture. Phylogenetically it clusters with *Roigiella lignicola*, which is a synnematous genus in the *Arthrobotrys* complex (Castañeda-Ruiz 1984). Morphologically *R. syzygii* is distinct from *R. lignicola* (conidia 11–17 × 2–3 µm) in that it has larger conidia that are encased in a mucilaginous sheath.


Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Roigiella lignicola* (strain CBS 222.85, GenBank MH861869.1; Identities = 511/519 (98 %), three gaps (0 %)), *Arthrobotrys dendroides* (strain CBS 431.85, GenBank NR_159642.1; Identities = 545/568 (96 %), five gaps (0 %)), and *Arthrobotrys ellipsospora* (strain NBRC 31826, GenBank LC146721.1; Identities = 538/561 (96 %), nine gaps (1 %)). Closest hits using the LSU sequence are *Roigiella lignicola* (strain CBS 222.85, GenBank MH873557.1; Identities = 882/886 (99 %), no gaps), *Arthrobotrys superba* (strain YNUCC 3628, GenBank AY261154.1; Identities = 862/872 (99 %), no gaps), and *Arthrobotrys botryosporus* (strain YNUCC 2728, GenBank AY261146.1; Identities = 862/872 (99 %), no gaps).

Consensus phylogram (50 % majority rule) obtained from the maximum likelihood analysis with IQ-TREE v. 2.1.3 (Minh et al. 2020) of the *Orbiliomycetes* LSU nucleotide alignment. Bootstrap support values (> 74 % are shown) from 1000 non-parametric bootstrap replicates are shown at the nodes. Culture collection numbers and GenBank accession numbers (superscript) are indicated for all species. The tree was rooted to *Candida broadrunensis* (culture CBS 11838; GenBank KY106372.1) and the species described here is highlighted with **bold face**. Families, orders and classes are shown in coloured blocks. Alignment statistics: 44 strains including the outgroup; 855 characters including alignment gaps analysed: 209 distinct patterns, 191 parsimony-informative, 59 singleton sites, 605 constant sites. The best model identified for the entire alignment in IQ-TREE using the TESTNEW option was: TN+F+I+G4. The alignment and tree were deposited in figshare. com (10.6084/m9.figshare.16918582).

Colour illustrations. *Syzygium cordatum* in Buffelskloof Nature Reserve. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia with mucoid sheath. Scale bars = 10 µm



Protocreopsis euphorbiae



Fungal Planet 1286 – 24 December 2021

***Protocreopsis euphorbiae* Crous, sp. nov.**

Etymology. Name refers to the host genus *Euphorbia* from which it was isolated.

Classification — *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*.

Conidiophores solitary to penicillate, aggregated in clusters forming superficial sporodochia (up to 450 µm diam), with orange mucoid spore masses. *Conidiophores* branched, septate, subcylindrical, up to 150 µm tall, 2.5–3.5 µm diam. *Conidiogenous cells* integrated, terminal and intercalary, phialidic, hyaline, smooth, subcylindrical to aculiform, flexuous with slight apical taper towards truncate apex, 1–1.5 µm diam, with periclinal thickening lacking collarette, 20–35 × 2.5–3 µm. *Conidia* solitary, aseptate, hyaline, smooth, granular, ellipsoid, apex subobtuse, base truncate, 1–1.5 µm diam, (3.5–)6–7(–7.5) × (3–)4 µm.

Culture characteristics — Colonies flat, spreading, with sparse to moderate aerial mycelium and feathery, lobate margin, reaching 20 mm diam after 2 wk at 25 °C. On MEA, PDA and OA surface and reverse orange.

Typus. SOUTH AFRICA, Northern Province, Nelspruit, Buffelskloof Nature Reserve, on leaf litter of *Euphorbia ingens* (*Euphorbiaceae*), Nov. 2018, P.W. Crous, HPC 3140 (holotype CBS H-24484, culture ex-type CPC 38896 = CBS 146972, ITS, LSU, *actA*, *tef1* (second part) and *tub2* sequences GenBank OK664700.1, OK663739.1, OK651123.1, OK651198.1 and OK651202.1, MycoBank MB 841780).

Notes — *Protocreopsis* (based on *P. musicola* = syn. *P. fusigera*) is a nectria-like genus with a hyphal stroma and acremonium-like asexual morph that was treated by Rossman et al. (1999). The present collection, known only from its asexual morph, is best accommodated in this genus based on its phylogenetic relationship to other acremonium-like species of *Protocreopsis*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Acremonium rutilum* (strain JCM 23088, GenBank NR_077124.1; Identities = 565/587 (96 %), 11 gaps (1 %)), *Protocreopsis phormiicola* (strain CBS 567.76, GenBank MH861001.1; Identities = 520/597 (87 %), 24 gaps (4 %)), and *Beauveria bassiana* (strain BbN06B02, GenBank MK952492.1; Identities = 484/558 (87 %), 21 gaps (3 %)). Closest hits using the **LSU** sequence are *Acremonium rutilum* (strain MY6, GenBank KY873378.1; Identities = 860/863 (99 %), no gaps), *Acremonium persicinum* (strain CBS 378.70C, GenBank HQ232080.1; Identities = 809/812 (99 %), one gap (0 %)), and *Pronectria robergei* (strain CBS 128021, GenBank MH876199.1; Identities = 855/869 (98 %), one gap (0 %)). Distant hits obtained using the **actA** sequence had highest similarity to *Penicillifer bipapillatus* (strain CBS 420.88, GenBank KM231105.1; Identities = 375/406 (92 %), no gaps), *Dactylonectria novozelandica* (strain CBS 113552, GenBank KM231157.1; Identities = 380/416 (91 %), two gaps (0 %)), and *Dactylonectria alcacerensis* (strain CBS 129087, GenBank KM231158.1; Identities = 380/416 (91 %), two gaps (0 %)). Distant hits obtained using the **tef1** (second part) sequence had highest similarity to *Podospora curvicolla* (strain VLV, GenBank X96614.1; Identities = 439/478 (92 %), two gaps (0 %)), *Clonostachys rosea* (strain GJS 90-227, GenBank AY489611.1; Identities = 438/477 (92 %), no gaps), and *Engyodontium parvisporum* (strain IHEM 22910, GenBank LC425558.1; Identities = 426/464 (92 %), two gaps (0 %)). Distant hits obtained using the **tub2** sequence had highest similarity to *Septofusidium herbarum* (strain CBS 265.58, GenBank KM232113.1; Identities = 310/367 (84 %), 19 gaps (5 %)), *Atractium crassum* (as *Fusarium merismoides* var. *crassum*; strain F-241,346, GenBank EU860033.1; Identities = 303/359 (84 %), 13 gaps (3 %)), and *Allantonectria miltina* (strain CBS 125499, GenBank KM232107.1; Identities = 303/359 (84 %), 15 gaps (4 %)).

Colour illustrations. Buffelskloof Nature Reserve. Colonies forming sporodochia on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1284.

Cryptometrion metrosideri



Fungal Planet 1287 – 24 December 2021

***Cryptometrion metrosideri* Crous, sp. nov.**

Etymology. Name refers to the host genus *Metrosideros* from which it was isolated.

Classification — *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*.

Conidiomata on PNA erumpent, solitary, globose, pycnidial with central ostiole, punctiform, 100–150 µm diam, red-brown to orange, exuding a creamy conidial mass. **Conidiophores** lining the inner cavity, hyaline, smooth, subcylindrical, septate, 10–20 × 2–3.5 µm, intermingled among subcylindrical, hyaline, smooth, septate paraphyses with obtuse ends. **Conidiogenous cells** subcylindrical, integrated, terminal and intercalary, phialidic, hyaline, smooth, 3–7 × 2–3 µm. **Conidia** solitary, subcylindrical, hyaline, smooth, aseptate, (2.5–)3–4(–6) × 1.5 µm, exuding in orange droplets.

Culture characteristics — Colonies flat, spreading, with moderate aerial mycelium and smooth, even margin, covering dish after 2 wk at 25 °C. On MEA surface cinnamon to sepia, reverse sepia; on PDA surface and reverse orange; on OA surface sepia.

Typus. NEW ZEALAND, Auckland, Devonport, Lake Road, on *Metrosideros* sp. (*Myrtaceae*), 10 Apr. 2019, K. Hofer (holotype CBS H-24413, culture ex-type CPC 38512 = T19_03462C = CBS 146785, ITS, LSU, *rpb2*, *tef1* (first part) and *tub2* sequences GenBank OK664701.1, OK663740.1, OK651158.1, OK651182.1 and OK651203.1, MycoBank MB 841781).

Additional material examined. NEW ZEALAND, Auckland, Mangere, on *Metrosideros excelsa*, 20 Aug. 2013, N. Scott, culture T13_03336.

Notes — *Cryptometrion metrosideri* is closely related to *C. aestuescens* (CBS 124014, *Eucalyptus grandis*, Indonesia; Gryzenhout et al. 2010). Morphologically the asexual morph is similar to *C. aestuescens*, as are the cultures, which on MEA turn sienna to umber with age. The present collection is thus best accommodated in *Cryptometrion*, pending additional collections to better resolve this generic complex.


Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Aurapex penicillata* (strain CR1207B, GenBank JQ864529.1; Identities = 615/650 (95 %), 13 gaps (2 %)), *Diversimorbus metrosideri* (strain CMW 37322, GenBank NR_153977.1; Identities = 577/618 (93 %), 15 gaps (2 %)), and *Corticimorbus sinomyrti* (strain CERC3634, GenBank KT167171.1; Identities = 583/630 (93 %), 22 gaps (3 %)). Closest hits using the **LSU** sequence are *Cryptometrion aestuescens* (strain CBS 124014, GenBank MH874864.1; Identities = 804/805 (99 %), no gaps), *Holocryphia eucalypti* (strain CMW 7037, GenBank JQ862753.1; Identities = 770/772 (99 %), no gaps), and *Diversimorbus metrosideri* (strain CMW 37322, GenBank NG_070365.1; Identities = 770/772 (99 %), no gaps). Closest hits using the **rpb2** sequence are *Cryptometrion aestuescens* (strain CBS 124007, GenBank MN271798.1; Identities = 649/662 (98 %), no gaps), *Holocryphia eucalypti* (strain CBS 115842, GenBank MN271804.1; Identities = 734/756 (97 %), no gaps), and *Corticimorbus sinomyrti* (strain CBS 140205, GenBank MN271794.1; Identities = 428/441 (97 %), no gaps). Closest hits using the **tef1** (first part) sequence had highest similarity to *Cryptometrion aestuescens* (strain CBS 124007, GenBank MN271851.1; Identities = 509/563 (90 %), 17 gaps (3 %)), *Ursicollum fallax* (strain CBS 118663, GenBank MN271897.1; Identities = 512/576 (89 %), 19 gaps (3 %)), and *Diversimorbus metrosiderotis* (strain CBS 132866, GenBank MN271857.1; Identities = 515/580 (89 %), 30 gaps (5 %)). Distant hits using the **tub2** sequence had highest similarity to *Cryphonectria macrospora* (voucher YMJ 94031513, GenBank EF025618.1; Identities = 699/811 (86 %), 41 gaps (5 %)), *Diversimorbus metrosideri* (as *Cryphonectriaceae* sp. SFC-2012a; strain CMW 37320, GenBank JQ862951.1; Identities = 414/444 (93 %), two gaps (0 %)), and *Parvomorbus guangdongensis* (strain CSF10738, GenBank MN258814.1; Identities = 388/424 (92 %), 11 gaps (2 %)).

Colour illustrations. Devonport, New Zealand. Conidioma on pine needle agar (PNA); conidiogenous cells giving rise to conidia; conidia. Scale bars = 150 µm (conidioma), 10 µm (all others).

Supplementary material

FP1287 Phylogenetic tree.

Harzia combreti



Fungal Planet 1288 – 24 December 2021

***Harzia combreti* Crous, sp. nov.**

Etymology. Name refers to the host genus *Combretum* from which it was isolated.

Classification — *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*.

Mycelium consisting of hyaline to olivaceous, smooth, branched, septate, 3.5–4 µm diam hyphae. **Conidiophores** macronematous, hyaline, smooth, subcylindrical, multiseptate with terminal and intercalary conidiogenous cells; **conidiogenous cells** hyaline, smooth, subcylindrical with apical taper, 20–70 × 3–4.5 µm. **Conidia** pale brown, smooth to finely roughened, granular, aseptate, dry, ovoid, thick-walled, (18–)20–22(–23) × (13–)15–17(–18) µm, with marginal frill.


Culture characteristics — Colonies flat, spreading, with moderate aerial mycelium and feathery, lobate margin, reaching 50 mm diam after 2 wk at 25 °C. On MEA surface pale luteous, reverse luteous; on PDA surface and reverse saffron; on OA surface pale luteous.

Typus. SOUTH AFRICA, Northern Province, Nelspruit, Lowveld Botanical Garden, on leaf litter of *Combretum collinum* ssp. *sulvense* (*Combretaceae*), 24 Nov. 2018, P.W. Crous, HPC 3170 (holotype CBS H-24881, culture ex-type CPC 39059 = CBS 148439, ITS and LSU sequences GenBank OK664702.1 and OK663741.1, MycoBank MB 841782).

Notes — *Harzia* is characterised by sympodially branched, hyaline superficial mycelium, hyaline to brown ovoid conidia and a *Proteophiala* synasexual morph (Domsch et al. 2007, Schulthes et al. 2017). *Harzia combreti* is closely related to an isolate identified as *H. patula*. However, *H. patula* has larger conidia ((16–)25–37.5(–50) × (12.5–)15–28(–37.5) µm; Holubová-Jechová 1974) and thus easily distinguished from *H. combreti*. Other similar species include *H. acremonioides* (conidia 20–30 × 15–20 µm; Domsch et al. 2007) and *H. metrosideri* (conidia (15–)16–18(–20) × (12–)15–16 µm; Crous et al. 2019d), from which it is also morphologically and phylogenetically distinct.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to '*Harzia patula*' (as *Olpitrichum patulum*; strain CBS 121524, GenBank KY628686.1; Identities = 644/657 (98 %), two gaps (0 %)), *Harzia patula* (strain CBS 379.88, GenBank NR_161009.1; Identities = 636/658 (97 %), seven gaps (1 %)), and *Harzia acremonioides* (strain NRRL 54327, GenBank HQ698593.1; Identities = 594/615 (97 %), five gaps (0 %)). Closest hits using the **LSU** sequence are '*Harzia patula*' (as *Olpitrichum patulum*; strain CBS 121524, GenBank KY628687.1; Identities = 829/832 (99 %), no gaps), *Harzia acremonioides* (strain NRRL 54327, GenBank HQ698593.1; Identities = 502/505 (99 %), one gap (0 %)), and *Harzia macrospora* (strain CBS 343.67, GenBank MH870687.1; Identities = 827/832 (99 %), one gap (0 %)).

The first of 61 equally most parsimonious trees obtained from a phylogenetic analysis of the *Harzia* ITS nucleotide alignment. The tree was rooted to *Sphaerodes retispora* var. *retispora* (culture CBS 994.72; GenBank KY628702.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1 000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and **bold** text. GenBank accession (superscript) and/or culture collection/voucher numbers (in **bold** face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 24 strains including the outgroup; 598 characters including alignment gaps analysed: 300 constant, 212 variable and parsimony-uninformative and 86 parsimony-informative. Tree statistics: Tree Length = 466, Consistency Index = 0.856, Retention Index = 0.740, Rescaled Consistency Index = 0.634. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).



10

Colour illustrations. Leaves of *Combretum collinum* ssp. *sulvense*. Conidiophores on PNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.



Fungal Planet 1289 – 24 December 2021

***Phaeoisaria dalbergiae* Crous, sp. nov.**

Etymology. Name refers to the host genus *Dalbergia* from which it was isolated.

Classification — *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** erect, subcylindrical, hyaline, smooth, 0–2-septate, unbranched or branched at apex, 10–50 × 1.5–2.5 µm, giving rise to 1–4 apical conidiogenous cells; **conidiogenous cells** lateral, arising in apical clusters, 7–15 × 1.5–2.5 µm, with terminal whorl of denticles, subcylindrical, hyaline, smooth, 0.5–2 × 1 µm, not thickened nor darkened. **Conidia** solitary, hyaline, smooth, aseptate, thin-walled, guttulate, subcylindrical to obovoid, tapering towards both ends, apex subobtuse, base with truncate hilum, 0.5 µm diam, (5–)6–7 × (1.5–)2 µm.


Culture characteristics — Colonies flat, spreading, surface folded, with sparse aerial mycelium and smooth, lobate margin, reaching 6–12 mm diam after 2 wk at 25 °C. On MEA, PDA and OA surface and reverse isabelline.

Typus. SOUTH AFRICA, Northern Province, Nelspruit, Buffelskloof Nature Reserve, on bark of *Dalbergia armata* (*Fabaceae*), 23 Nov. 2018, P.W. Crous, HPC 3147 (holotype CBS H-24882, culture ex-type CPC 39540 = CBS 148440, ITS, LSU, *rpb2* and SSU sequences GenBank OK664703.1, OK663742.1, OK663796.1 and OK651159.1, MycoBank MB 841783).

Notes — *Phaeoisaria* includes species with solitary conidiophores and synnemata, with polyblastic, denticulate conidiogenous cells (Seifert et al. 2011). *Phaeoisaria dalbergiae* was originally isolated as a synnematous mycophylic fungus on *Dalbergia armata* in South Africa. In culture however, only solitary conidiophores were observed. *Phaeoisaria dalbergiae* is closely related to *P. clematidis* (conidia 5–7 × 2–3 µm; Luo et al. 2018) and *P. annesophieae* (conidia 4.5–9 × 2–3.5 µm; Crous et al. 2017), from which it is phylogenetically distinct.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Phaeoisaria clematidis* (strain CBS 113340, Gen-

Bank EU552148.1; Identities = 554/570 (97 %), one gap (0 %)), *Phaeoisaria annesophieae* (voucher MFLU 19-0531, GenBank MT559109.1; Identities = 518/549 (94 %), 11 gaps (2 %)), and *Phaeoisaria filiformis* (voucher MFLU 18-1462, GenBank NR_168819.1; Identities = 488/525 (93 %), 12 gaps (2 %)). Closest hits using the **LSU** sequence are *Phaeoisaria annesophieae* (voucher MFLU 19-0531, GenBank MT559084.1; Identities = 806/816 (99 %), no gaps), *Phaeoisaria* sp. JB-2020a (strain FCCUGF 03, GenBank MT375866.1; Identities = 805/815 (99 %), no gaps), and *Phaeoisaria clematidis* (strain MFLUCC 18-1017, GenBank MW132065.1; Identities = 783/793 (99 %), no gaps). Closest hits using the **rpb2** sequence had highest similarity to *Phaeoisaria clematidis* (strain MFLUCC 17-1341, GenBank MF401400.1; Identities = 455/489 (93 %), no gaps), *Phaeoisaria aquatica* (as *Phaeoisaria* sp. ZLL-2017a; strain MFLUCC 16-1298, GenBank MF401406.1; Identities = 611/669 (91 %), no gaps), and *Phaeoisaria microspora* (strain M0104, GenBank MF167352.1; Identities = 666/757 (88 %), seven gaps (0 %)). Closest hits using the **SSU** sequence are *Phaeoisaria clematidis* (strain MFLUCC 17-1968, GenBank MG837027.1; Identities = 987/992 (99 %), no gaps), *Phaeoisaria fasciculata* (strain DAOM 230055, GenBank KT278694.1; Identities = 919/927 (99 %), no gaps), and *Phaeoisaria guttulate* (as *Phaeoisaria* sp. JY-2018; voucher MFLU 18-0139, GenBank MG837026.1; Identities = 983/992 (99 %), no gaps).



The first of two equally most parsimonious trees obtained from a phylogenetic analysis of the *Phaeoisaria* ITS nucleotide alignment. The tree was rooted to *Pleurotheciella aquatica* (voucher MFLU 17-0911; GenBank NR_160591.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1 000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and bold text. GenBank accession (superscript) and/or culture collection/voucher numbers (in bold face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 33 strains including the outgroup; 520 characters including alignment gaps analysed: 321 constant, 96 variable and parsimony-uninformative and 101 parsimony-informative. Tree statistics: Tree Length = 334, Consistency Index = 0.778, Retention Index = 0.894, Rescaled Consistency Index = 0.696. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Colour illustrations. Stem of *Dalbergia armata*. Conidiophores and conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

10



Fungal Planet 1290 – 24 December 2021

Pseudosoloacrosporiella* Crous, gen. nov.Etymology.* Morphologically similar to *Soloacrosporiella*.Classification — *Incertae sedis*, *Microthyriales*, *Dothideomycetes*.*Mycelium* consisting of pale brown, smooth, branched, septate hyphae. *Conidiophores* solitary, erect, pale brown, smooth, subcylindrical with apical taper, 0–1-septate, mostly reduced to conidiogenous cells. *Conidiogenous cells* with 1–3 sympodialapical loci with rhexolytic conidiogenesis, leaving minute collarettes on denticulate loci. *Ramoconidia* fusoid-ellipsoid, tapering towards both ends, pale brown, smooth, guttulate, septate. *Conidia* in long unbranched chains, septate, fusoid-ellipsoid, pale brown, smooth; loci rhexolytic, with minute marginal frill.*Type species.* *Pseudosoloacrosporiella cryptomeriae* Crous
MycoBank MB 841784.***Pseudosoloacrosporiella cryptomeriae* Crous, sp. nov.***Etymology.* Name refers to the host genus *Cryptomeria* from which it was isolated.*Mycelium* consisting of pale brown, smooth, branched, septate, 1.5–2 µm diam hyphae. *Conidiophores* solitary, erect, pale brown, smooth, subcylindrical with apical taper, 0–1-septate, mostly reduced to conidiogenous cells. *Conidiogenous cells* 15–30 × 3–4 µm, with 1–3 sympodial apical loci with rhexolytic conidiogenesis, leaving minute collarettes on denticulate loci, 0.5 µm diam. *Ramoconidia* fusoid-ellipsoid, tapering towards both ends, pale brown, smooth, guttulate, 1–3-septate, 15–20 × 3–4 µm, with 1–3 apical loci. *Conidia* in long (–15) unbranched chains, 0–1-septate, fusoid-ellipsoid, pale brown, smooth, (7–)12–13(–15) × (2.5–)3(–3.5) µm; loci rhexolytic, 0.5 µm diam, with minute marginal frill.

Culture characteristics — Colonies flat, spreading, surface folded, with moderate aerial mycelium and smooth, lobate margin, reaching 8 mm diam after 7 d at 25 °C. On MEA, PDA and OA surface and reverse ochreous.

Typus. NETHERLANDS, Gelderland Province, Wageningen, Belmonte Botanical Garden, on leaves of *Cryptomeria japonica* (Cupressaceae), 28 July 2020, P.W. Crous, HPC 3301 (holotype CBS H-24883, culture ex-type CPC 39587 = CBS 148441, ITS, LSU and *tef1* (first part) sequences GenBank OK664704.1, OK663743.1 and OK651183.1, MycoBank MB 841785).Notes — *Pseudosoloacrosporiella* is phylogenetically closely related to *Soloacrosporiella*, but distinct in that it lacks setae, and its conidia do not have thickened and darkened hila, but rather have a characteristic marginal frill (also on conidiogenous loci), which also distinguishes it from *Soloacrospora* and *Neocladophialophora* (Castañeda Ruiz et al. 1997, Crous et al. 2014b).Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Neocladophialophora quercina* (strain CBS 138874, GenBank NR_137948.1; Identities = 497/557 (89 %), 17 gaps (3 %)) and *Soloacrosporiella acaciae* (strain CBS 139894, GenBank NR_137986.1; Identities = 413/494 (84 %), 38 gaps (7 %)). Closest hits using the LSU sequence are *Neocladophialophora quercina* (strain CBS 138874, GenBank MH877670.1; Identities = 772/810 (95 %), one gap (0 %)), *Soloacrosporiella acaciae* (strain CBS 139894, GenBank NG_058164.1; Identities = 761/814 (93 %), two gaps (0 %)), and *Neoscolecobasidium agapanthi* (strain CPC 28778, GenBank NG_059748.1; Identities = 695/793 (88 %), 14 gaps (1 %)). No significant hits were obtained when the *tef1* (first part) sequence was used in blastn and megablast searches.*Colour illustrations.* *Cryptomeria japonica*. Conidiophores on SNA, giving rise to branched chains of conidia; conidia. Scale bars = 10 µm.**Supplementary material****FP1290** Phylogenetic tree.



Fungal Planet 1291 – 24 December 2021

***Parasitella quercicola* Crous & Denman, sp. nov.**

Etymology. Name refers to the host genus *Quercus* from which it was isolated.

Classification — *Mucoraceae*, *Mucorales*, *Mucoromycetes*.


Mycelium consisting of rhizoid-like hyphae, 4–7 µm diam, smooth, hyaline. **Sporangiophores** arising from aerial mycelium, simple or sympodially branched, erect, smooth-walled, hyaline, 5–7 µm diam. **Sporangia** olivaceous, globose to subglobose, 25–60 µm diam, shortly echinulate; wall evanescent, leaving columella with short collarettes. **Columellae** hyaline, smooth-walled, globose to subglobose, 11–17 µm diam. **Sporangiospores** hyaline, smooth-walled, aseptate, ellipsoid to reniform, (5–)6–7(–9) × (3–)3.5(–4) µm. **Oidia** and **zygosporengia** not observed.

Culture characteristics — Colonies erumpent, spreading, with abundant aerial mycelium, reaching 7 cm diam after 7 d at 25 °C. On MEA surface smoke grey, reverse luteous; on PDA surface and reverse dirty white; on OA surface white.

Typus. UK, E379015.3 N243233.1, on *Quercus robur* (*Fabaceae*), 29 July 2020, S. Denman, FPPH 459 (holotype CBS H-24884, culture ex-type CPC 39612 = CBS 148442, ITS and LSU sequences GenBank OK664705.1 and OK663744.1, MycoBank MB 841786).

Notes — *Parasitella quercicola* is closely related to *P. parasitica* (neotype CBS 412.66), a species known to be myco-parasitic on other fungi, including *Mucorales* (Schipper 1978, O'Donnell et al. 2001). *Parasitella* is presently seen as monotypic, with *P. quercicola* being phylogenetically distinct from *P. parasitica*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Parasitella parasitica* (strain CBS 208.28, GenBank MH854982.1; Identities = 540/593 (91 %), 21 gaps (3 %)), *Mucor plumbeus* (strain LF9, GenBank KX011014.1; Identities = 489/603 (81 %), 54 gaps (8 %)), and *Mucor racemosus* f. *chibinensis* (strain LT4-3, GenBank MW850517.1; Identities = 488/603 (81 %), 52 gaps (8 %)). Closest hits using the LSU sequence are *Parasitella parasitica* (strain CBS 153.69, GenBank MH871012.1; Identities = 853/880 (97 %), nine gaps (1 %)), *Mucor mousanensis* (strain CBS 999.70, GenBank MH871807.1; Identities = 823/879 (94 %), 13 gaps (1 %)), and *Mucor plumbeus* (strain FMR 16013, GenBank LT984540.1; Identities = 821/879 (93 %), 13 gaps (1 %)).



The first of 11 equally most parsimonious trees obtained from a phylogenetic analysis of the *Mucor* ITS nucleotide alignment. The tree was rooted to *Absidia montepascoalii* (voucher URM 8218; GenBank NR_172995.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and bold text. GenBank accession (superscript) and/or culture collection/voucher numbers (in bold face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 62 strains including the outgroup; 809 characters including alignment gaps analysed: 153 constant, 164 variable and parsimony-uninformative and 492 parsimony-informative. Tree statistics: Tree Length = 3931, Consistency Index = 0.364, Retention Index = 0.612, Rescaled Consistency Index = 0.223. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Colour illustrations. *Quercus robur*. Sporangioophores; sporangioophores with sporangia, sporangiospores, and columella; sporangiospores. Scale bars = 10 µm.

10



Fungal Planet 1292 – 24 December 2021

***Paramacroventuria* Crous & Bulgakov, gen. nov.**

Etymology. Name refers to the fact that it is closely related to *Macroventuria*.

Classification — *Didymellaceae*, *Pleosporales*, *Dothideomycetes*.

Conidiomata pycnidial, globose, brown, with central periphysate ostiole; wall of 3–6 layers of medium brown *textura angularis*.

Conidiophores reduced to conidiogenous cells lining inner cavity, hyaline, smooth, ampulliform, proliferating percurrently at apex. **Conidia** solitary, brown, finely roughened, medianly septate, guttulate, ellipsoid, apex obtuse, base with truncate hilum.

Type species. *Paramacroventuria ribis* Crous & Bulgakov
Mycobank MB 841787.

***Paramacroventuria ribis* Crous & Bulgakov, sp. nov.**

Etymology. Name refers to the host genus *Ribes* from which it was isolated.

Leaf spots superficial, round, merging, 1–6 mm diam, firstly medium brown, later fair grey, with raised, dark brown-purple edge. **Conidiomata** pycnidial, globose, brown, 150–250 µm diam, with central periphysate ostiole; wall of 3–6 layers of medium brown *textura angularis*. **Conidiophores** reduced to conidiogenous cells lining inner cavity, hyaline, smooth, ampulliform, 5–8 × 4–7 µm, proliferating percurrently at apex. **Conidia** solitary, brown, finely roughened, medianly septate, guttulate, ellipsoid, apex obtuse, base with truncate hilum, 2–3 µm diam, (9–)10–12(–15) × 5(–6) µm.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, even margin, reaching 35 mm diam after 7 d at 25 °C. On MEA surface umber, reverse isabelline; on PDA surface and reverse isabelline; on OA surface isabelline.

Typus. RUSSIA, Rostov region, Shakhty city district, urban street greenery, on *Ribes aureum* (*Grossulariaceae*), 18 July 2020, T.S. Bulgakov, HPC 3326 = PC 107 (holotype CBS H-24885, isotype LE F-332405, culture ex-type CPC 39692 = CBS 148443, ITS, LSU, *rpb2* and *tub2* sequences GenBank OK664706.1, OK663745.1, OK651160.1 and OK651204.1, MycoBank MB 841788).

Notes — *Paramacroventuria* clusters close to *Juxtiphoma* and *Macroventuria*. *Juxtiphoma* was introduced to accommodate *Phoma eupyrena* reported on stems of *Solanum tuberosum* (Valenzuela-Lopez et al. 2018), while the sexual genus *Macroventuria* is based on *M. anomochaeta*, which was described from decayed canvas (Hou et al. 2020b). *Paramacroventuria* is an asexual phoma-like genus, distinct from *Juxtiphoma* in that it lacks chlamydospores, and has percurrently proliferating conidiogenous cells.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Briansuttonomyces eucalypti* (strain CBS 114879, GenBank NR_153618.1; Identities = 535/541 (99 %), no gaps), *Macroventuria wentii* (strain CBS 526.71, GenBank MH860250.1; Identities = 534/540 (99 %), no gaps), and *Leptosphaerulina australis* (strain CBS 297.54, GenBank MH857342.1; Identities = 536/543 (99 %), one gap (0 %)). Closest hits using the **LSU** sequence are *Microsphaeropsis hellebori* (strain CBS 569.82, GenBank MH873273.1; Identities = 820/821 (99 %), no gaps), *Didymella aurea* (strain CBS 269.93, GenBank NG_069042.1; Identities = 819/821 (99 %), no gaps), and *Microsphaeropsis ononidicola* (strain MFLUCC 15-0459, GenBank MG967668.1; Identities = 819/821 (99 %), no gaps). Closest hits using the **rpb2** sequence had highest similarity to *Macroventuria anomochaeta* (strain CBS 525.71, GenBank GU456346.1; Identities = 691/751 (92 %), no gaps), *Paraboeremia adianticola* (strain CBS 187.83, GenBank KP330401.1; Identities = 693/760 (91 %), no gaps), and *Juxtiphoma eupyrena* (as *Phoma eupyrena*; strain CBS 374.91, GenBank LT623268.1; Identities = 692/760 (91 %), no gaps). Closest hits using the **tub2** sequence had highest similarity to *Ascochyta pilosella* (strain CBS 583.97, GenBank MT005696.1; Identities = 316/333 (95 %), no gaps), *Paramicrosphaeropsis ellipsoidea* (strain CBS 197.97, GenBank MT005680.1; Identities = 314/333 (94 %), no gaps), and *Stagonosporopsis helianthi* (strain CBS 200.87, GenBank KT389848.1; Identities = 314/334 (94 %), three gaps (0 %)).

Colour illustrations. *Ribes aureum*. Conidiomata on PNA; colony sporulating on OA; conidioma; conidiogenous cells giving rise to conidia; conidia. Scale bars = 250 µm (conidioma), 10 µm (all others).

Supplementary material

FP1292 Phylogenetic tree.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Timur S. Bulgakov, Department of Plant Protection, Federal Research Centre the Subtropical Scientific Centre of the Russian Academy of Sciences, Yana Fabritsiusa street 2/28, 354002 Sochi, Krasnodar region, Russia; e-mail: fungi-on-don@yandex.ru

Myrmecridium sambuci



Fungal Planet 1293 – 24 December 2021

***Myrmecridium sambuci* Crous, sp. nov.**

Etymology. Name refers to the host genus *Sambucus* from which it was isolated.

Classification — *Myrmecridiaceae*, *Myrmecridiales*, *Sordariomycetes*.

On SNA. *Mycelium* consisting of hyaline, smooth, branched, septate, 1.5–2.5 µm diam hyphae. *Conidiophores* unbranched, erect, straight, medium brown, thick-walled, (1–)3–7-septate, up to 170 µm tall, 2.5–3 µm diam; basal cell 3–5 µm diam. *Conidiogenous cells* terminal, integrated, subcylindrical, 30–75 µm long, pale brown, forming a rachis with pimple-shaped denticles less than 1 µm long, 0.5 µm diam, slightly thickened. *Conidia* solitary, aseptate, pale brown, thin-walled, smooth, guttulate, with or without wing-like gelatinous sheath in middle, ellipsoid to fusoid, (7–)8–9(–10) × (2.5–)3(–3.5) µm; hilum unthickened, nor darkened, 0.5 µm diam.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 25 mm diam after 7 d at 25 °C. On MEA surface honey to isabelline, reverse isabelline; on PDA surface and reverse luteous; on OA surface umber.

Typus. NETHERLANDS, Zeeland Province, Oranjestad, on *Sambucus nigra* (*Adoxaceae*), 17 Sept. 2020, A.L. van Iperen, HPC 3510 (holotype CBS H-24886, culture ex-type CPC 39959 = CBS 148444, ITS and LSU sequences GenBank OK664707.1 and OK663746.1, MycoBank MB 841789).

Notes — *Myrmecridium* was established by Arzanlou et al. (2007) for ramichloridium-like fungi with solitary conidiophores having pimple-shaped denticles along the medium brown rachis, and subhyaline conidia that frequently have a wing-like gelatinous sheath. *Myrmecridium sambuci* is closely related to *M. phragmiticola* known from *Phragmites australis* in Ukraine (conidia ellipsoid to fusoid, (7–)8–9 × (2.5–)3 µm, conidiophores 2–4-septate, up to 70 µm tall, 3–3.5 µm diam; basal cell 4–6 µm diam; Crous et al. 2020b), but can be distinguished based on its taller and narrower conidiophores.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Myrmecridium phragmiticola* (strain CPC 36367, GenBank NR_170826.1; Identities = 541/551 (98 %), three gaps (0 %)), *Myrmecridium schulzeri* (strain NTOU 4421, GenBank MZ422994.1; Identities = 541/555 (97 %), three gaps (0 %)), and *Myrmecridium fluviae* (strain CNUFC-YR61-2, GenBank KX839679.1; Identities = 456/471 (97 %), four gaps (0 %)). Closest hits using the **LSU** sequence are *Myrmecridium phragmiticola* (strain CPC 36367, GenBank NG_074444.1; Identities = 799/802 (99 %), one gap (0 %)), *Myrmecridium schulzeri* (strain CBS 188.96, GenBank EU041829.1; Identities = 794/802 (99 %), one gap (0 %)), and *Myrmecridium banksiae* (strain CBS 132536, GenBank NG_042684.1; Identities = 793/802 (99 %), one gap (0 %)).

Colour illustrations. Zeeland Province, the Netherlands. Conidiophores on SNA; conidiophores with conidiogenous cells giving rise to conidia; conidia with mucoid sheath. Scale bars = 10 µm.

Myrmecridium is based on *M. sambuci* (basionym: *Psilobotrys schulzeri*, isolated from a trunk of *Carpinus betulus*, Croatia). Unfortunately, neither De Hoog (1977) nor Arzanlou et al. (2007) could locate the type specimen, which is presumably lost. Since its description, however, numerous isolates have been assigned to this species name, which cluster phylogenetically throughout the genus. To facilitate a more stable taxonomy, we therefore designate the reference strain used by Arzanlou et al. (2007) as neotype below.

***Myrmecridium schulzeri* (Sacc.) Arzanlou et al., Stud. Mycol. 58: 84. 2007**

Basionym. *Psilobotrys schulzeri* Sacc., Hedwigia 23: 126. 1884.
For additional synonyms see Arzanlou et al. (2007).

Typus. NETHERLANDS, Lienden, isolated from *Triticum aestivum* root, C.L. de Graaff (CBS 325.74, preserved as metabolically inactive culture, designated as neotype here, MBT 10003688, culture ex-neotype CBS 325.74 = JCM 7234).

Myrmecridium aquaticum (conidia 3-septate, 14–16 × 4–6 µm, on submerged decaying wood, China; Luo et al. 2019), is morphologically similar to, and represents the older name for *Neomyrmecridium guizhouense* (conidia (2–)3-septate, 8.9–12.7 × 2.8–4.8 µm, on submerged decaying wood, China; Hyde et al. 2020), and phylogenetically, they also cluster in the same clade.

***Neomyrmecridium aquaticum* (Z.L. Luo et al.) Crous, comb. nov.** MycoBank MB 841858


Basionym: *Myrmecridium aquaticum* Z.L. Luo et al., Fungal Diversity 99: 501. 2019.

Synonym: *Neomyrmecridium guizhouense* N.G. Liu et al., Fungal Diversity 100: 187. 2020.

Supplementary material

FP1293 Phylogenetic tree.

Neofabraea salicina



Fungal Planet 1294 – 24 December 2021

***Neofabraea salicina* Crous, sp. nov.**

Etymology. Name refers to the host genus *Salix* from which it was isolated.

Classification — *Dermateaceae*, *Helotiales*, *Leotiomyces*.

Sporulating poorly on SNA. *Conidiomata* poorly developed, stromatic, acervular. *Conidiophores* simple to branched, smooth, hyaline, septate. *Conidiogenous cells* integrated, phialidic, smooth, hyaline, cylindrical to narrowly ampulliform, giving rise to macro- and microconidia, 10–15 × 2.5–3.5 µm. *Macroconidia* cylindrical-fusoid to ellipsoid, aseptate, hyaline, smooth-walled, with basal scar, (10–)15–17(–25) × (4–)5–6(–6.5) µm. *Microconidia* cylindrical, apex obtuse, base truncate, aseptate, hyaline, smooth-walled, (5–)7–8 × (2–)2.5–3 µm.


Culture characteristics — Colonies erumpent, spreading, surface folded, with sparse aerial mycelium and smooth, lobate margin, reaching 10 mm diam after 7 d at 25 °C. On MEA surface dirty white, reverse ochreous; on PDA surface and reverse dirty white; on OA surface sienna.

Typus. NETHERLANDS, Gelderland Province, Beuningen, river Waal, on *Salix* sp. (*Salicaceae*), 17 Oct. 2020, A.L. van Iperen, HPC 3485 (holotype CBS H-24887, culture ex-type CPC 39973 = CBS 148445, ITS, LSU, *rpb2*, *tef1* (first part) and *tub2* sequences GenBank OK664708.1, OK663747.1, OK651161.1, OK651184.1 and OK651205.1, MycoBank MB 841790).

Notes — Although *Phlyctema vagabunda* (syn. *Neofabraea alba*) is a known pathogen of *Salix* spp., we are not aware of *Neofabraea* spp. on this host. *Neofabraea salicina* is closely related to species such as *N. brasiliensis* (macroconidia 12–22 × 2.5–3.7 µm; Crous et al. 2015), *N. kienholzii* (macroconidia 12–17.5 × 2.5–3.5 µm; Spotts et al. 2009) *N. perennans* (macroconidia 12–25 × 3–6 µm; Spotts et al. 2009) and *N. malicorticis* (macroconidia 16–31 × 4.5–6 µm; Chen et al. 2016), but can easily be distinguished based on its DNA phylogeny.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Neofabraea perennans* (strain CBS 453.64,

GenBank KR859089.1; Identities = 536/541 (99 %), no gaps), *Neofabraea malicorticis* (strain CBS 102863, GenBank KR859085.1; Identities = 536/541 (99 %), no gaps), and *Neofabraea inaequalis* (strain CBS 326.75, GenBank NR_155470.1; Identities = 535/541 (99 %), no gaps). Closest hits using the LSU sequence are *Neofabraea malicorticis* (strain CBS 122030, GenBank NG_068983.1; Identities = 761/764 (99 %), two gaps (0 %)), *Neofabraea kienholzii* (strain KARE1951, GenBank MK160254.1; Identities = 761/764 (99 %), two gaps (0 %)), and *Neofabraea perennans* (strain PPO-45431, GenBank KX768551.1; Identities = 761/764 (99 %), two gaps (0 %)). Closest hits using the *rpb2* sequence had highest similarity to *Neofabraea malicorticis* (strain CBS 102863, GenBank KR859325.1; Identities = 722/735 (98 %), no gaps), *Neofabraea perennans* (strain GC57, GenBank MN729470.1; Identities = 721/735 (98 %), no gaps), and *Neofabraea inaequalis* (strain CBS 326.75, GenBank KR859321.1; Identities = 711/735 (97 %), no gaps). Closest hits using the *tef1* (first part) sequence had highest similarity to *Neofabraea perennans* (strain CBS 139.41, GenBank KX982710.1; Identities = 510/531 (96 %), seven gaps (1 %)), *Neofabraea malicorticis* (strain CBS 102863, GenBank KX982708.1; Identities = 494/534 (93 %), 23 gaps (4 %)), and *Neofabraea kienholzii* (strain CBS 355.72, GenBank KX982716.1; Identities = 248/269 (92 %), four gaps (1 %)). Closest hits using the *tub2* sequence had highest similarity to *Neofabraea eucalyptorum* (strain CBS 146634, GenBank MT375121.1; Identities = 527/578 (91 %), six gaps (1 %)), *Phlyctema phoenicis* (strain CPC 29372, GenBank KY173611.1; Identities = 314/384 (82 %), 13 gaps (3 %)), and *Coleophoma eucalyptorum* (strain CBS 131314, GenBank KU728606.1; Identities = 453/589 (77 %), 45 gaps (7 %)).



The first of five equally most parsimonious trees obtained from a phylogenetic analysis of the *Neofabraea rpb2* nucleotide alignment. The tree was rooted to *Pezicula acericola* (culture CBS 245.97; GenBank KF376213.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and **bold** text. GenBank accession (superscript) and/or culture collection/voucher numbers (in **bold** face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 17 strains including the outgroup; 736 characters including alignment gaps analysed: 546 constant, 99 variable and parsimony-uninformative and 91 parsimony-informative. Tree statistics: Tree Length = 268, Consistency Index = 0.832, Retention Index = 0.858, Rescaled Consistency Index = 0.714. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Colour illustrations. Gelderland Province, Beuningen, river Waal. Conidiophores and conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Polyscytalum vaccinii



Fungal Planet 1295 – 24 December 2021

***Polyscytalum vaccinii* Crous, sp. nov.**

Etymology. Name refers to the host genus *Vaccinium* from which it was isolated.

Classification — *Phlogicylindriaceae*, *Xylariales*, *Sordariomycetes*.

Mycelium consisting of pale brown, smooth, septate, branched, 1.5–2 µm diam hyphae. **Conidiophores** erect, flexuous, dark brown, thick-walled, septate, up to 200 µm tall, 2.5–3 µm diam. **Ramoconidia** giving rise to branched chains of cylindrical conidia, pale brown, smooth-walled, subcylindrical, 0(–1)-septate, (12–)15–18(–20) × (2–)2.5 µm, with 1–4 flat-tipped scars, 1–1.5 µm diam. **Conidia** hyaline to pale olivaceous, cylindrical, 0(–1)-septate, guttulate, (10–)13–16(–20) × 2(–2.5) µm.

Culture characteristics — Colonies flat, spreading, with moderate aerial mycelium and feathery, lobate margin, reaching 10 mm diam after 7 d at 25 °C. On MEA surface and reverse umber, with diffuse sienna pigment; on PDA surface and reverse umber; on OA surface umber with diffuse umber pigment.


Typus. NETHERLANDS, Limburg Province, Weertbos, on *Vaccinium myrtillus* (*Ericaceae*), 6 Sept. 2020, A.L. van Iperen, HPC 3463 (holotype CBS H-24888, culture ex-type CPC 39935 = CBS 148446, ITS and LSU sequences GenBank OK664709.1 and OK663748.1, MycoBank MB 841791).

Notes — *Polyscytalum* was treated by Crous et al. (2018b), with several species placed in *Cylindrium*. *Polyscytalum vaccinii* is phylogenetically distinct from taxa presently known in the genus, and morphologically distinguishable based on its


conidial dimensions. It is similar but distinct from *P. pinicola* (conidiophores up to 80 µm tall, conidia (13–)14–15(–16) × 2 µm; Crous et al. 2020c) based on a combination of conidiophore and conidium characteristics. It also needs to be compared to *P. pini-canariensis* (conidia (18–)22–26(–48) × 3(–3.5) µm; Crous et al. 2020c), *P. fecundissimum* (conidia 13–18 × 2 mm) and *P. neofecundissimum* (conidia (12–)14–17(–20) × 2(–3) µm) (Crous et al. 2018b), which have overlapping conidial dimensions, but are phylogenetically distinct.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Anungitea eucalyptigena* (strain CPC 28762, GenBank NR_164411.1; Identities = 404/417 (97 %), two gaps (0 %)), *Polyscytalum neofecundissimum* (strain CBS 143390, GenBank NR_158959.1; Identities = 567/589 (96 %), 11 gaps (1 %)), and *Anungitea nullicana* (strain CBS 143406, GenBank NR_156391.1; Identities = 401/417 (96 %), two gaps (0 %)). Closest hits using the **LSU** sequence are *Polyscytalum neofecundissimum* (strain CBS 143390, GenBank NG_066207.1; Identities = 812/817 (99 %), two gaps (0 %)), *Polyscytalum pinicola* (strain CPC 36759, GenBank NG_074425.1; Identities = 798/805 (99 %), one gap (1 %)), and *Polyscytalum pini-canariensis* (strain CBS 146819, GenBank NG_074496.1; Identities = 805/813 (99 %), one gap (0 %)).

The first of 16 equally most parsimonious trees obtained from a phylogenetic analysis of the *Polyscytalum* / *Phlogicylindrium* ITS nucleotide alignment. The tree was rooted to *Sarcostroma grevilleae* (culture CBS 101.71; GenBank MH553952.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and bold text. GenBank accession (superscript) and/or culture collection/voucher numbers (in bold face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 26 strains including the outgroup; 590 characters including alignment gaps analysed: 321 constant, 75 variable and parsimony-uninformative and 194 parsimony-informative. Tree statistics: Tree Length = 572, Consistency Index = 0.682, Retention Index = 0.772, Rescaled Consistency Index = 0.526. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).



Colour illustrations. *Vaccinium myrtillus*. Conidiophores and conidigenous cells giving rise to conidial chains; conidia. Scale bars = 10 µm.



Fungal Planet 1296 – 24 December 2021

***Ramularia pararhabdospora* Crous, sp. nov.**

Etymology. Name refers to its morphological similarity to *Ramularia rhabdospora*.

Classification — *Mycosphaerellaceae*, *Mycosphaerellales*, *Dothideomycetes*.

Leaf spots amphigenous, subcircular, medium brown with raised margin, 3–7 mm diam. *Conidiomata* fasciculate, amphigenous, stroma absent, arising from stomata in moderately dense fascicles. *Conidiophores* 0–2-septate, subcylindrical, hyaline, smooth-walled, 15–55 × 4–5 µm. *Conidiogenous cells* integrated, subcylindrical, straight to geniculate-sinuous, hyaline, smooth-walled, 15–25 × 3–4 µm; scars thickened, darkened, refractive, 1.5–2 µm diam. *Conidia* solitary, hyaline, smooth-walled, subcylindrical, tapering from middle to subobtuse apex, base obconically rounded, with truncate hilum, thickened, darkened and refractive, 1.5–2 µm diam, 3(–5)-septate, (38–)45–55(–75) × (3–)3.5–4 µm. In culture conidia are up to 100 µm long, flexuous, and undergo microcyclic conidiation.

Culture characteristics — Colonies erumpent, spreading, with sparse to moderate aerial mycelium and smooth, lobate margin, reaching 5 mm diam after 7 d at 25 °C. On MEA surface saffron, reverse ochrous; on PDA surface and reverse grey olivaceous in centre, pale luteous in outer region; on OA surface smoke grey.

Typus. NETHERLANDS, Gelderland Province, Beuningen, river Waal, on *Plantago lanceolata* (*Plantaginaceae*), 17 Oct. 2020, A.L. van Iperen, HPC 3473 (holotype CBS H-24889, culture ex-type CPC 39983 = CBS 148447, ITS, LSU, *actA*, *gapdh*, *his3*, *rpb2* and *tef1* (first part) sequences GenBank OK664710.1, OK663749.1, OK651124.1, OK651147.1, OK651148.1, OK651162.1 and OK651185.1, MycoBank MB 841792).

Notes — Two species are known from this host in Europe. *Ramularia rhabdospora* is common on *Plantago lanceolata*, and has catenate conidia that are (10–)15–40(–50) × 3–7 µm, 0–3(–4)-septate. *Ramularia plantaginis* has catenate, ellipsoid-ovoid to cylindrical conidia, 10–35 × 3–6 µm, 0–3-septate (Braun 1998), thus different from those of *R. pararhabdospora*. *Ramularia rhabdospora* and *R. pararhabdospora* are phylogenetically related and it is possible that more cryptic species exist within *R. rhabdospora*.

Colour illustrations. Gelderland Province, Beuningen, river Waal. Leaf spot; conidiophores and conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Ramularia hieracii-umbellati* (strain CPC 10691 GenBank KX287429.1; Identities = 518/522 (99 %), no gaps), *Ramularia rhabdospora* (strain CBS 118415, GenBank KX287492.1; Identities = 518/525 (99 %), no gaps), and *Ramularia kriegiana* (strain CPC 10827, GenBank KX287461.1; Identities = 509/516 (99 %), one gap (0 %)). Closest hits using the **LSU** sequence are *Ramularia veronicicola* (strain CBS 113981, GenBank KX287241.1; Identities = 804/804 (100 %), no gaps), *Ramularia rhabdospora* (strain CBS 312.92, GenBank KX287193.1; Identities = 804/804 (100 %), no gaps), and *Ramularia bosniaca* (strain CBS 123880, GenBank KX287045.1; Identities = 804/804 (100 %), no gaps). Closest hits using the **actA** sequence had highest similarity to *Ramularia rhabdospora* (strain CBS 118415, GenBank KX287774.1; Identities = 552/568 (97 %), one gap (0 %)), *Ramularia inaequalis* (strain CPC 25742, GenBank KP894336.1; Identities = 538/582 (92 %), 12 gaps (2 %)), and *Ramularia armoraciae* (strain CBS 241.90, GenBank KX287613.1; Identities = 528/573 (92 %), six gaps (1 %)). Closest hits using the **gapdh** sequence had highest similarity to *Ramularia rhabdospora* (strain CBS 118415, GenBank KX288332.1; Identities = 500/522 (96 %), no gaps), *Ramularia digitalis-ambiguae* (strain CBS 434.67, GenBank KX288247.1; Identities = 424/462 (92 %), five gaps (1 %)), and *Ramularia hieracii-umbellati* (strain CPC 10789, GenBank KX288272.1; Identities = 411/455 (90 %), five gaps (1 %)). Closest hits using the **his3** sequence had highest similarity to *Ramularia rhabdospora* (strain CBS 312.92, GenBank KX288940.1; Identities = 330/342 (96 %), no gaps), *Ramularia gaultheriae* (strain CBS 299.80, GenBank KX288859.1; Identities = 324/343 (94 %), two gaps (0 %)), and *Ramularia tricherae* (strain CBS 108995, GenBank KX288968.1; Identities = 322/342 (94 %), no gaps). Closest hits using the **rpb2** sequence had highest similarity to *Ramularia rhabdospora* (strain CBS 312.92, GenBank KX288651.1; Identities = 758/781 (97 %), no gaps), *Ramularia digitalis-ambiguae* (strain CBS 297.37, GenBank KX288567.1; Identities = 721/781 (92 %), no gaps), and *Ramularia geraniicola* (strain CPC 25912, GenBank KX288574.1; Identities = 716/778 (92 %), no gaps). Closest hits using the **tef1** (first part) sequence had highest similarity to *Ramularia rhabdospora* (strain CBS 118415, GenBank KX288052.1; Identities = 294/310 (95 %), one gap (0 %)), *Ramularia abscondita* (strain CBS 114727, GenBank KX287864.1; Identities = 325/357 (91 %), seven gaps (1 %)), and *Ramularia hieracii-umbellati* (strain CPC 10789, GenBank KX287993.1; Identities = 323/356 (91 %), four gaps (1 %)).

Supplementary material

FP1296 Phylogenetic tree.



Fungal Planet 1297 – 24 December 2021

***Cyphellophora sambuci* Crous, sp. nov.**

Etymology. Name refers to the host genus *Sambucus* from which it was isolated.

Classification — *Cyphellophoraceae*, *Chaetothyriales*, *Eurotiomycetes*.

Mycelium consisting of pale brown, smooth, guttulate, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** reduced to solitary conidiogenous cells on hyphae. **Conidiogenous cells** phialidic, flared, pale brown, 1–3 × 1–3 µm, giving rise to dimorphic conidia. **Macroconidia** subcylindrical to narrowly obclavate, guttulate, straight to slightly bent, apex subobtuse, base obconically truncate, (1–)3-septate, constricted at septa, hilum truncate, 1 µm diam, not darkened nor thickened, (15–)18–22(–28) × (1.5–)2 µm. **Microconidia** solitary, aseptate, pale brown, obovoid to ellipsoid, 2–5 × 1.5–2.5 µm.

Culture characteristics — Colonies erumpent, sectored, with moderate aerial mycelium and smooth, lobate margin, reaching 7 mm diam after 7 d at 25 °C. On MEA and PDA surface pale olivaceous grey, margin iron grey, reverse iron grey; on OA surface iron grey.

Typus. NETHERLANDS, South Holland Province, Goeree-Overflakkee, De Kwade Hoek, on *Sambucus nigra* (*Adoxaceae*), 13 Sept. 2020, A.L. van Iperen, HPC 3507 (holotype CBS H-24890, culture ex-type CPC 39957 = CBS 148448, ITS, LSU and *tub2* sequences GenBank OK664711.1, OK663750.1 and OK651206.1, MycoBank MB 841793).


Notes — *Cyphellophora* is characterised by having elongated, curved, one- to multiseptate conidia, and well-developed, flared phialides, which frequently develop directly on fungal hyphae (Crous et al. 2009). Species have been isolated from human skin and nail infections (De Hoog et al. 2000, Réblová et al. 2013), soil, plants, water and other substrates (Feng et al. 2014, Crous et al. 2019b, 2020a). *Cyphellophora sambuci* is phylogenetically closely related to *C. phyllostachydis* (conidia 0–3 septate, 25–30(–45) × 1.5–2 µm; Gao et al. 2015) and

C. europaea (conidia aseptate, 1.8–2.5 µm diam; De Hoog et al. 2000), but can be distinguished based on its conidial dimensions.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Cyphellophora europaea* (strain CF27, GenBank KY038650.1; Identities = 469/493 (95 %), seven gaps (1 %)), *Cyphellophora oxyspora* (strain CBS 416.89, GenBank MH862181.1; Identities = 491/533 (92 %), ten gaps (1 %)), and *Cyphellophora phyllostachydis* (strain HMAS 245769, GenBank NR_158393.1; Identities = 563/613 (92 %), 25 gaps (4 %)). Closest hits using the **LSU** sequence are *Cyphellophora europaea* (strain CBS 101466, GenBank KC455259.1; Identities = 807/814 (99 %), one gap (0 %)), *Cyphellophora phyllostachydis* (strain HLHNZWYZ08, GenBank KP122933.1; Identities = 806/814 (99 %), one gap (0 %)), and *Cyphellophora musae* (strain GLMMZZ4, GenBank KP122934.1; Identities = 754/763 (99 %), one gap (0 %)). Closest hits using the **tub2** sequence had highest similarity to *Cyphellophora phyllostachydis* (strain HLHNZWYZ08, GenBank KP122929.1; Identities = 284/343 (83 %), seven gaps (2 %)), *Cyphellophora europaea* (strain CBS 101466, GenBank KC455229.1; Identities = 317/393 (81 %), 12 gaps (3 %)), and *Cyphellophora ludoviensis* (as *Cyphellophora* sp. RG-2016; strain CMRP1317, GenBank KX583749.1; Identities = 283/357 (79 %), 24 gaps (6 %)).

The first of 10 equally most parsimonious trees obtained from a phylogenetic analysis of the *Cyphellophora* ITS nucleotide alignment. The tree was rooted to *Exophiala salmonis* (culture CBS 157.67; GenBank MH858932.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and **bold** text. GenBank accession (superscript) and/or culture collection/voucher numbers (in **bold** face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 31 strains including the outgroup; 681 characters including alignment gaps analysed: 328 constant, 99 variable and parsimony-uninformative and 254 parsimony-informative. Tree statistics: Tree Length = 1379, Consistency Index = 0.520, Retention Index = 0.623, Rescaled Consistency Index = 0.324. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Colour illustrations. *Sambucus nigra* at De Kwade Hoek, South Holland Province. Conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.



10

Paradissoconium narthecii



Fungal Planet 1298 – 24 December 2021

Paradissoconium* Crous & Boers, gen. nov.Etymology.* Name refers to its morphological similarity to *Dissoconium*.Classification — *Dissoconiaceae*, *Mycosphaerellales*, *Dothideomycetes*.*Mycelium* consisting of hyaline, smooth, branched, septate hyphae. *Conidiophores* solitary, unbranched, erect, subcylindrical to subulate, with truncate apex, straight to gently curved, smooth, pale brown, septate. *Conidiogenous cells* terminal, integrated, loci slightly darkened, proliferating sympodially, andrejuvenating percurrently. *Primary conidia* solitary, straight, constricted at median septum, smooth, hyaline, ellipsoid, apex obtuse, base obconically truncate, hilum pronounced, darkened. *Secondary conidia* adjacent to primary conidia, hyaline, smooth, aseptate, ellipsoid, tapering to truncate hilum; anastomosing with primary conidia after active discharge.*Type species.* *Paradissoconium narthecii* Crous & Boers
MycoBank MB 841794.***Paradissoconium narthecii* Crous & Boers, sp. nov.***Etymology.* Name refers to the host genus *Narthecium* from which it was isolated.*Ascospores* could not be linked to specific ascomata on host tissue, as several taxa were involved, including species with pseudothecia and thyrothecia. *Ascospores* smooth-walled, fusoid-ellipsoid, widest just above septum, germinating with flexuous germ tubes from both polar ends, spore becoming brown and verruculose, 6 µm diam, constricted at septum, not distorted. *Mycelium* consisting of hyaline, smooth, branched, septate, 2–3 µm diam hyphae. *Conidiophores* solitary, unbranched, erect, subcylindrical to subulate, with truncate apex, straight to gently curved, smooth, pale brown, 0–2-septate, 15–100 × 3.5–4.5 µm. *Conidiogenous cells* terminal, integrated, 15–45 × 3–4 µm, loci slightly darkened, 1–2 µm diam, cicatrized, at times subdenticulate, proliferating sympodially, and rejuvenating percurrently. *Primary conidia* (12–)14–16(–18) × (5–)6–7 µm, solitary, straight, constricted at median septum, smooth, hyaline, ellipsoid, apex obtuse, base obconically truncate, hilum pronounced, darkened, 2–2.5 µm diam. *Secondary conidia* 6–9 × 5–6 µm, adjacent to primary conidia, hyaline, smooth, aseptate, ellipsoid, tapering to truncate hilum, 2–2.5 µm; anastomosing with primary conidia after active discharge.

Culture characteristics — Colonies erumpent, spreading, with moderate aerial mycelium and feathery, lobate margin, reaching 6 mm diam after 7 d at 25 °C. On MEA, PDA and OA surface and reverse olivaceous grey.

Typus. NETHERLANDS, Drenthe Province, Dwingelderveld National Park, 52.829188, 6.432495, on dead leaves of *Narthecium ossifragum* (*Nartheciaceae*), 4 July 2021, J. Boers, HPC 3653 (holotype CBS H-24891, cultures ex-type CPC 41970, 41971 = CBS 148449, ITS, LSU, *actA* and *rpb1* sequences GenBank OK664712.1, OK663751.1, OK651125.1 and OK651151.1, MycoBank MB 841795).*Additional materials examined.* NETHERLANDS, Drenthe Province, Dwingelderveld National Park, 52.829188, 6.432495, on dead leaves of *N. ossifragum*, 4 July 2021, J. Boers, HPC 3653, single ascospore cultures CPC 41976, 41977.Notes — The distinction between *Dissoconium* and *Uwebraunia* was discussed by Li et al. (2012). Both genera have solitary conidiophores with actively discharging aseptate secondary and 1-septate primary conidia that anastomose upondischarge onto the agar surface. *Uwebraunia* and *Dissoconium* have pseudothecial sexual morphs (Crous et al. 2004, Hyde et al. 2020). *Paradissoconium* has a typical dissoconium-like asexual morph that develops in culture, and is morphologically similar to *Dissoconium* s.str. Thus far *Paradissoconium* is monotypic but appears commonly on dead leaves of *Narthecium ossifragum*, where it is associated with thyrothecia. Although we tried on several locations to link the asexual and sexual morph, all attempts were unsuccessful, as too many diverse sexual morphs were involved. *Paradissoconium* is located basal in *Dissoconiaceae*.Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to '*Dissoconium* sp. GS-2011' (strain MYQWGNL19, GenBank JF701607.1; Identities = 483/491 (98 %), four gaps (0 %)), *Ramichloridium punctatum* (strain NTOU 4892, GenBank MK448255.1; Identities = 465/509 (91 %), 14 gaps (2 %)), and *Dissoconium proteae* (strain CBS 122900, GenBank NR_156213.1; Identities = 469/514 (91 %), 17 gaps (3 %)). Closest hits using the **LSU** sequence are *Ramichloridium eucleeae* (strain CBS 138000, GenBank NG_058086.1; Identities = 814/828 (98 %), two gaps (0 %)), *Ramichloridium indicum* (strain CBS 171.96, GenBank EU041856.1; Identities = 811/827 (98 %), no gaps), and *Ramichloridium apiculatum* (strain CPC 12310, GenBank GU214687.1; Identities = 809/828 (98 %), two gaps (0 %)). Closest hits using the **actA** sequence had highest similarity to *Uwebraunia communis* (as *Mycosphaerella communis*; strain X1023, GenBank EU514299.1; Identities = 490/569 (86 %), 29 gaps (5 %)), *Pseudophaeophleospora phormii* (strain CBS 144606, GenBank MK442643.1; Identities = 427/480 (89 %), 16 gaps (3 %)), and *Zasmidium commune* (strain CBS 142528, GenBank KY979857.1; Identities = 366/392 (93 %), one gap (0 %)). Distant hits obtained using the **rpb1** sequence had highest similarity to *Nothophaeocryptopus gaeumannii* (strain CBS 267.37, GenBank GU357770.1; Identities = 513/723 (71 %), 34 gaps (4 %)), *Zymoseptoria tritici* (strain ST99CH_3D1, GenBank LT854276.1; Identities = 493/717 (69 %), 27 gaps (3 %)), and *Rasutoria tsugae* (strain ratstk, GenBank GU371809.1; Identities = 489/720 (68 %), 31 gaps (4 %)).**Supplementary material****FP1298** Phylogenetic tree.*Colour illustrations.* *Narthecium ossifragum*. Conidiophores on SNA with primary and secondary conidia; primary and secondary conidia that discharge together and then anastomose. Scale bars = 10 µm.

Paramyrothecium lathyri



Fungal Planet 1299 – 24 December 2021

***Paramyrothecium lathyri* Crous & Bulgakov, sp. nov.**

Etymology. Name refers to the host genus *Lathyrus* from which it was isolated.

Classification — *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*.

Spots on living leaves and stems amphigenous, subcircular, 5–14 mm diam, fair brown to grey, with clear border. *Conidiomata* sporodochial, stromatic, cupulate, superficial, scattered, oval, 200–300 µm diam, with white, setose fringe, surrounding an olivaceous green mucoid conidial mass. *Stroma* well developed, of hyaline to subhyaline *textura angularis*. *Setae* arising from stroma, thick-walled, 3–10-septate, flexuous, up to 300 µm long, 3–4 µm diam, tapering to rounded apex. *Conidiophores* arising from basal stroma, consisting of penicillate, branched conidiophores; stipes unbranched, hyaline, septate, smooth-walled, 5–10 × 2–3.5 µm; primary branches aseptate, smooth-walled, 5–10 × 2–3 µm; secondary branches aseptate, smooth-walled, 5–8 × 2–3 µm; terminal whorl of 3–6 *conidiogenous cells* cylindrical to subcylindrical, phialidic, hyaline, smooth-walled, straight to slightly curved, 5–10 × 2–3 µm, with periclinal thickening and minute collarettes. *Conidia* aseptate, subhyaline, smooth-walled, cylindrical, ends rounded, guttulate, (8–)9(–10) × 2(–2.5) µm.

Culture characteristics — Colonies erumpent, spreading, surface folded, with moderate aerial mycelium and smooth, lobate margin, reaching 50 mm diam after 7 d at 25 °C. On MEA surface pale luteous, reverse ochreous; on PDA surface and reverse pale luteous; on OA surface saffron.

Typus. Russia, Rostov region, Krasnosulinsky district, state natural wildlife area 'Gomensky', edge of artificial forest, on *Lathyrus tuberosus* (*Fabaceae*), 10 July 2020, T.S. Bulgakov, HPC 3370 = PC 084 (holotype CBS H-24793, isotype LE F-332408, culture ex-type CPC 39712 = CBS 148248, ITS, LSU, *cmdA*, *rpb2* and *tef1* (first part) sequences GenBank OK664713.1, OK663752.1, OK651141.1, OK651163.1 and OK651186.1, MycoBank MB 841796).

Colour illustrations. State natural wildlife area 'Gomensky', Russia. Symptomatic leaf of *Lathyrus tuberosus*. Conidiomata with setae on PNA; conidioma with setae on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 300 µm (conidiomata), 10 µm (all others).

Notes — *Paramyrothecium* is distinguished from *Myrothecium* by having septate, thin-walled setae surrounding its sporodochia. Morphologically *P. lathyri* is distinct from the other species presently known, based on a combination of its conidial morphology and multiseptate setae (Lombard et al. 2016, Crous et al. 2021a). *Paramyrothecium lathyri* is located basal in *Paramyrothecium*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Myrothecium lachae* (strain 13-1913, GenBank MH001945.1; Identities = 538/563 (96 %), eight gaps (1 %)), *Paramyrothecium parvum* (strain CBS 257.35, GenBank NR_145076.1; Identities = 556/582 (96 %), seven gaps (1 %)), and *Paramyrothecium roridum* (as *Myrothecium roridum*; strain BBA 62764, GenBank AJ301993.1; Identities = 563/590 (95 %), seven gaps (1 %)). Closest hits using the **LSU** sequence are *Paramyrothecium roridum* (strain CBS 357.89, GenBank NG_069343.1; Identities = 816/820 (99 %), one gap (0 %)), *Paramyrothecium breviseta* (strain CBS 544.75, GenBank NG_069340.1; Identities = 816/820 (99 %), one gap (0 %)), and *Paramyrothecium foliicola* (strain Bas4_m2, GenBank MH807539.1; Identities = 816/820 (99 %), one gap (0 %)). Closest hits using the **cmdA** sequence had highest similarity to *Paramyrothecium pituitipietianum* (strain CPC 38688, GenBank MW173100.1; Identities = 528/631 (84 %), 23 gaps (3 %)), *Paramyrothecium foliicola* (strain QB3, GenBank MT415358.1; Identities = 500/601 (83 %), 20 gaps (3 %)), and *Paramyrothecium tellicola* (strain CBS 478.91, GenBank KU846272.1; Identities = 497/602 (83 %), 24 gaps (3 %)). Closest hits using the **rpb2** sequence had highest similarity to *Paramyrothecium sinense* (strain ZSY21, GenBank MH818825.1; Identities = 526/593 (89 %), no gaps), *Paramyrothecium roridum* (as *Myrothecium roridum*; strain UAMH 1369, GenBank DQ676588.1; Identities = 568/642 (88 %), no gaps), and *Paramyrothecium nigrum* (strain CBS 116537, GenBank KU846357.1; Identities = 637/721 (88 %), no gaps). Closest hits using the **tef1** (first part) sequence had highest similarity to *Neomyrothecium humicola* (strain CBS 310.96, GenBank KU846527.1; Identities = 211/233 (91 %), six gaps (2 %)), *Gregatothecium humicola* (strain CBS 205.96, GenBank KU846402.1; Identities = 168/174 (97 %), two gaps (1 %)), and *Albifimbria verrucaria* (strain 207C9, GenBank KU845942.1; Identities = 201/224 (90 %), seven gaps (3 %)).

Supplementary material

FP1299 Phylogenetic tree.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Timur S. Bulgakov, Department of Plant Protection, Federal Research Centre the Subtropical Scientific Centre of the Russian Academy of Sciences, Yana Fabritsiusa street 2/28, 354002 Sochi, Krasnodar region, Russia; e-mail: fungi-on-don@yandex.ru

Colpoma junipericola



Fungal Planet 1300 – 24 December 2021

***Colpoma junipericola* Crous & Bulgakov, sp. nov.**

Etymology. Name refers to the host genus *Juniperus* from which it was isolated.

Classification — *Rhytismataceae*, *Rhytismatales*, *Leotiomyces*.

Ascomata amphigenous on leaf litter, lens-shaped, immersed in epidermis, black, up to 1 mm long, 300–400 µm diam, opening by longitudinal slit. **Asci** cylindrical-clavate, short-stalked, 8-spored, not staining in Melzer's reagent, 70–95 × 8–9 µm. **Ascospores** filiform, tapering more prominently to base, hyaline, aseptate, guttulate, at times with gelatinous sheath, (25–)30–35(–40) × 1.5–2 µm. **Paraphyses** filiform, extending above asci, hyaline, septate, guttulate, apex clavate, 2–2.5 µm diam, up to 100 µm tall, 1.5–2 µm wide, covered in mucoid layer. **Conidiomata** developing in culture, acervular, 150–200 µm diam, pale brown. **Conidiophores** reduced to conidiogenous cells lining inner cavity, ampulliform to subcylindrical, straight, hyaline, smooth-walled, 6–15 × 2.5–3 µm, phialidic with periclinal thickening. **Conidia** aseptate, hyaline, smooth-walled, subcylindrical, straight with obtuse ends, 5–7 × 1.5–2 µm.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and feathery, lobate margin, reaching 40 mm diam after 7 d at 25 °C. On MEA surface saffron, reverse peach; on PDA surface and reverse saffron; on OA surface pale luteous..

Typus. RUSSIA, Rostov region, Shakhty, private garden, on needles of *Juniperus sabina* (*Cupressaceae*), 22 May 2020, T.S. Bulgakov, HPC 3350 = PC 118 (holotype CBS H-24792, isotype LE F-332409, culture ex-type CPC 39702 = CBS 148247, ITS and LSU sequences GenBank OK664714.1 and OK663753.1, MycoBank MB 841797).

Notes — The *Rhytismataceae* occurring on *Juniperus* was reviewed by Hou et al. (2005). *Colpoma junipericola* is related to taxa that have been described in *Colpoma* and *Coccoomyces*. Both genera are highly polyphyletic, and thus we tentatively describe the new taxon on *Colpoma*. It is distinct from *Colpoma juniperi* which has ascomata that open via irregular splits, ascospores filiform, aseptate, 40–50 × 1–2 µm, enclosed in mucoid sheath, and paraphyses that are curled or coiled (but not swollen) (Minter 1996), and *Coccoomyces juniperi*, which has ascospores 40–50 × 1–2 µm, filiform, often with branched paraphyses, and intracortical apothecia, opening via irregular splits (Sherwood 1980).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Colpoma quercinum* (strain C, GenBank AJ293879.1; Identities = 446/479 (93 %), five gaps (1 %)), *Coccoomyces strobili* (strain NB-645C, GenBank MH457134.1; Identities = 470/511 (92 %), 10 gaps (1 %)), and *Trybliopsis magnesi* (strain DAOMC 252096, GenBank NR_164611.1; Identities = 503/559 (90 %), 13 gaps (2 %)). Closest hits using the **LSU** sequence are *Colpoma quercinum* (strain COL, GenBank MT707243.1; Identities = 740/759 (97 %), no gaps), *Sporomega degenerans* (voucher Lantz 367 (UPS), GenBank HM140567.1; Identities = 737/759 (97 %), no gaps), and *Colpoma ledi* (voucher Lantz 379 (UPS), GenBank HM140512.1; Identities = 734/759 (97 %), no gaps).

Colour illustrations. *Juniperus sabina*. Needles with ascomata; conidiomata on PNA; asci with paraphyses; ascus; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material**FP1300** Phylogenetic tree.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Timur S. Bulgakov, Department of Plant Protection, Federal Research Centre the Subtropical Scientific Centre of the Russian Academy of Sciences, Yana Fabritsiusa street 2/28, 354002 Sochi, Krasnodar region, Russia; e-mail: fungi-on-don@yandex.ru

Fungal Planet 1301 & 1302 – 24 December 2021

***Neochalara* Crous, gen. nov.**

Etymology. Name refers to the genus *Chalara*, to which it is morphologically similar.

Classification — *Pezizellaceae*, *Rhytismatales*, *Leotiomyces*.

Mycelium consisting of hyaline, smooth-walled, branched, septate hyphae. *Conidiophores* arising from bulbous base, or as extension of hyphae, frequently aggregated, cylindrical,

***Neochalara spiraeae* Crous, sp. nov.**

Etymology. Name refers to the host genus *Spiraea* from which it was isolated.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 3–4 µm diam hyphae. *Conidiophores* arising from bulbous base, or as extension of hyphae, frequently aggregated, cylindrical, brown, erect, thick-walled, unbranched, straight to slightly bent, 1–4-septate, sometimes with the basal cell inflated, with terminal conidiogenous cell, 100–170 × 6–8 µm. *Conidiogenous cells* phialidic, brown, smooth-walled, consisting of a subcylindrical venter that gradually merges into a pale brown cylindrical collarette; venter 30–40 µm long, 7–10 µm wide, neck collarette 30–50 µm long, 4–5 µm wide, ratio of mean lengths of collarette and venter 1 : 1. *Conidia* in long, unbranched chains, cylindrical with both ends truncate with marginal frill, medianly 1-septate, hyaline, guttulate, smooth-walled, (11–)12–14(–16) × (3–)4(–4.5) µm, av. 13 × 4.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 30 mm diam after 7 d at 25 °C. On MEA surface luteous, reverse ochreous; on PDA surface and reverse ochreous; on OA surface ochreous.

***Sporidesmium spiraeae* Crous, sp. nov.**

Etymology. Name refers to the host genus *Spiraea* from which it was isolated.

Classification — *Sporidesmiaceae*, *Sporidesmiales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth, branched, septate, 1.5–2 µm diam hyphae. *Conidiophores* solitary or aggregated in clusters, erect, dark brown, subcylindrical, unbranched, 1–4-septate, thick-walled, 15–40 × 5–7 µm. *Conidiogenous cells* terminal, integrated, subcylindrical, blastic, medium brown, smooth-walled, 10–20 × 5–6 µm. *Conidia* solitary, obclavate, straight to flexuous, apex subobtuse, base obconically truncate, (4–)6–10-distoseptate, septa with central pore, wall smooth, pale brown, basal cell dark brown, hilum truncate, 4–5 µm diam, (45–)70–85(–100) × (8–)9(–10) µm.

Colour illustrations. *Spiraea japonica* at Belmonte Botanical Garden. Left column: *Neochalara spiraeae*. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidial chains. Right column: *Sporidesmium spiraeae*. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

brown, erect, thick-walled, unbranched, septate, with terminal conidiogenous cells. *Conidiogenous cells* phialidic, brown, smooth-walled, consisting of a venter that gradually merges into a cylindrical collarette that is pale brown. *Conidia* in long, unbranched chains, cylindrical with both ends truncate with marginal frill, medianly 1-septate, hyaline, guttulate, smooth-walled.

Type species. *Neochalara spiraeae* Crous
MycoBank MB 841798.

Typus. NETHERLANDS, Gelderland Province, Wageningen, Belmonte Botanical Garden, on *Spiraea japonica* (*Rosaceae*), 28 July 2020, P.W. Crous, HPC 3291 (holotype CBS H-24865, culture ex-type CPC 39565 = CBS 148332, ITS and LSU sequences GenBank OK664715.1 and OK663754.1, MycoBank MB 841799).

Notes — Although morphologically resembling members of the genus *Chalara*, the newly described species cannot be accommodated in any of the known *Chalara* species. Members of the genus *Chalara* with 1-septate conidia either have conidia with rounded ends, or a rounded apex and truncate base (Nag Raj & Kendrick 1976, McKenzie et al. 2002). However, *N. spiraeae* is distinct in having conidia with both ends being truncate. Furthermore, the distinct phylogenetic position also supports the taxonomical novelty. The present genus, *Neochalara*, appears to be allied to orbiliopsis-like taxa, although data to support this potential link presently remains inconclusive.

(notes *Neochalara spiraeae* continues on
Supplementary material page FP1301 & 1302)

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1300.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 8 mm diam after 7 d at 25 °C. On MEA surface dirty white, reverse pale luteous; on PDA surface and reverse dirty white; OA surface dirty white.

Typus. NETHERLANDS, Gelderland Province, Wageningen, Belmonte Botanical Garden, on *Spiraea japonica* (*Rosaceae*), 28 July 2020, P.W. Crous, HPC 3291 (holotype CBS H-24837, culture ex-type CPC 39766 = CBS 148298, ITS, LSU and *rpb2* sequences GenBank OK664716.1, OK663755.1 and OK651164.1, MycoBank MB 841800).

Notes — Although the type species of *Sporidesmium* (*S. atrum*) is not known from DNA data, *Sporidesmium* has been shown to be polyphyletic (Liu et al. 2019). No species of *Sporidesmium* (syn. *Ellisembia*) is known from *Spiraea*, and based on its phylogeny, as well as the morphological key provided by Wu & Zhuang (2005), *S. spiraeae* appears distinct.

(notes *Sporidesmium spiraeae* continues on
Supplementary material page FP1301 & 1302)

Supplementary material

FP1302 Phylogenetic tree.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Ondřej Koukol, Department of Botany, Charles University, Faculty of Science, Benátská 2, 128 01 Prague 2, Czech Republic; e-mail: ondrej.koukol@natur.cuni.cz

Toxicocladosporium glendoranum



Fungal Planet 1303 – 24 December 2021

***Toxicocladosporium glendoranum* Crous & Jurjević, sp. nov.**

Etymology. Name refers to the city where this fungus was collected, Glendora, California.

Classification — *Cladosporiaceae*, *Cladosporiales*, *Dothideomycetes*.

Conidiophores erect, arising from globose cells in superficial mycelium, 1–8-septate, up to 80 µm tall, 2.5–3.5 µm diam, brown, smooth-walled, mostly unbranched. **Conidiogenous cells** terminal and lateral, integrated, brown, smooth-walled, subcylindrical, 7–11 × 2–2.5 µm, with 1–3 apical loci, darkened, thickened, not refractive, 0.5 µm diam. **Conidia** occurring in branched chains, pale brown, smooth-walled, fusoid-ellipsoid; primary ramoconidia fusoid to subcylindrical, 0–1-septate, 7–20 × 2.5–3 µm; hila thickened and darkened, not refractive, 1 µm diam. Secondary ramoconidia aseptate, 5–10 × 2–2.5 µm; intercalary conidia in long chains (–12), fusoid-ellipsoid, (4–)5–6 × 1.5–2 µm; terminal conidia fusoid-ellipsoid, 3–4 × 1.5–2 µm; hila darkened, thickened, not refractive, 0.5 µm diam.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 7–10 mm diam after 7 d at 25 °C. On MEA, PDA, Czapek yeast agar (CYA) and OA surface greenish olivaceous, reverse iron grey.

Typus. USA, California, Glendora, from air in a brick room, Oct. 2019, Z. Jurjević, 5421 (holotype CBS H-24802, culture ex-type CPC 40763 = CBS 148256, ITS, LSU, *actA* and *tef1* (first part) sequences GenBank OK664717.1, OK663756.1, OK651126.1 and OK651187.1, MycoBank MB 841801).

Notes — *Toxicocladosporium glendoranum* clusters basal to *Toxicocladosporium* but is also closely related to several *Cladosporium* spp. (Crous et al. 2016, Bezerra et al. 2017, Bensch et al. 2018). Morphologically it is characterised by having rather short conidiophores, and branched conidial chains with smooth, primarily aseptate conidia. Phylogenetically it is distinct from all species presently known in *Toxicocladosporium*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Cladosporium halotolerans* (strain CBS 127014, GenBank MH864391.1; Identities = 516/550 (94 %), eight gaps (1 %)), *Cladosporium endophyticum* (strain MFLUCC 17-0599, GenBank NR_158360.1; Identities = 516/550 (94 %), eight gaps (1 %)), and *Cladosporium parahalotolerans* (strain CPC 22376, GenBank MF473155.1; Identities = 516/550 (94 %), eight gaps (1 %)). Closest hits using the **LSU** sequence are *Toxicocladosporium ficiniae* (strain CBS 136406, GenBank NG_058054.1; Identities = 792/810 (98 %), no gaps), *Toxicocladosporium hominis* (strain CBS 140694, GenBank MH878169.1; Identities = 773/794 (97 %), two gaps (0 %)), and *Toxicocladosporium irritans* (strain CBS 185.58, GenBank MH869283.1; Identities = 789/811 (97 %), two gaps (0 %)). Closest hits using the **actA** sequence had highest similarity to *Cladosporium velox* (strain CPC 18450, GenBank KT600654.1; Identities = 536/606 (88 %), 32 gaps (5 %)) and *Cladosporium inversicolor* (strain CPC 18238, GenBank KT600588.1; Identities = 530/602 (88 %), 26 gaps (4 %)). No significant hits were obtained when the **tef1** sequence was used in blastn and megablast searches.

Colour illustrations. Brick wall in Glendora, California. Conidiophores and conidiogenous cells giving rise to conidia; conidial chains. Scale bars = 10 µm.

Supplementary material

FP1303 Phylogenetic tree.



Fungal Planet 1304 – 24 December 2021

***Sporidesmiella juncicola* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Junewangiaceae*, *Incertae sedis*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** solitary, erect, macronematous, straight to somewhat flexuous, brown, smooth- and thick-walled, base swollen, 8–12 µm diam, 2–3-septate, 70–100 × 4–5 µm. **Conidiogenous cells** integrated, terminal, subcylindrical, scars inconspicuous, proliferating percurrently with delayed succession, conidia remaining attached via a marginal frill, rejuvenating sympodially, and appearing as polyblastic (sensu Luo et al. 2019; but in fact its percurrent, as can be seen from the marginal frill attached to the conidial hila), pale brown, smooth-walled, 20–45 × 4–5 µm. **Conidia** solitary, dry, subcylindrical to obovoid, medium brown, smooth-walled, guttulate, apex obtuse, base 2.5–3 µm diam, (3–)4-distoseptate, septa with central pore, (20–)28–32(–35) × (8–)9–10 µm.

Culture characteristics — Colonies erumpent, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 7 mm diam after 7 d at 25 °C. On MEA, PDA and OA surface saffron, reverse ochreous.

Typus. NETHERLANDS, Utrecht Province, Bornia, near Driebergen, 8 m a.s.l., N52°02'23" E05°17'56", on dead culm of *Juncus effusus* (*Juncaceae*), 25 Jan. 2021, *E.R. Osieck*, HPC 3583 = WI-15/#4211 (holotype CBS H-24850, culture ex-type CPC 41075 = CBS 148311, ITS, LSU and *rpb2* sequences GenBank OK664718.1, OK663757.1 and OK651165.1, MycoBank MB 841802).

Additional material examined. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'44" E05°09'40", on dead culm of *Juncus inflexus*, 4 Feb. 2021, *E.R. Osieck*, HPC 3590 = WI-21/#4205, CBS H-24824, culture CPC 41109 = CBS 148278, ITS, LSU, *rpb2* and *tef1* (first part) sequences GenBank OK664719.1, OK663758.1, OK651166.1 and OK651188.1.

Colour illustrations. *Juncus effusus* at Nieuw Wulven, near Houten. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.


Notes — The most common species of *Sporidesmiella* occurring in Europe (on various substrates) is *S. hyalosperma* (conidia brown, smooth, thick-walled, 4-distoseptate, (17–)22–25(–29) × 9–12(–13) µm, Kirk 1982; (19–)20–22(–24) × (9–)10(–11) µm, Crous et al. 2020c). Phylogenetically *S. juncicola* is closely related to *S. aquatica* (conidia 51–59 × 8–22, 3–4-distoseptate, on submerged wood, Luo et al. 2019), from which it is morphologically distinct based on its conidium dimensions.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence of CPC 41075 had highest similarity to *Sporidesmiella aquatica* (voucher MFLU 18-1602, GenBank NR_168811.1; Identities = 483/496 (97 %), five gaps (1 %)), *Sporidesmiella hyalosperma* (strain CPC 37552, GenBank MT223845.1; Identities = 507/573 (88 %), 24 gaps (4 %)), and *Sporidesmiella obovoidia* (strain MFLUCC 17-2372, GenBank NR_172446.1; Identities = 504/579 (87 %), 37 gaps (6 %)). The ITS sequences of CPC 41075 and CPC 41109 differ with a single indel (500/501 (99 %), one gap (0 %)). Closest hits using the **LSU** sequence of CPC 41075 are *Sporidesmiella aquatica* (as *Sporidesmiella* sp. ZLL-2019a; strain S-777, GenBank MK849843.1; Identities = 763/763 (100 %), no gaps), *Sporidesmiella obovoidia* (strain MFLUCC 17-2372, GenBank NG_075412.1; Identities = 780/803 (97 %), one gap (0 %)), and *Sporidesmiella hyalosperma* (strain S-1518, GenBank MK849842.1; Identities = 773/797 (97 %), one gap (0 %)). The LSU sequences of CPC 41075 and CPC 41109 differ with a single indel (799/800 (99 %), one gap (0 %)). Closest hits using the **rpb2** sequence of CPC 41075 had highest similarity to *Sporidesmiella aquatica* (voucher MFLU 18-2331, GenBank MN124524.1; Identities = 638/671 (95 %), no gaps), *Sporidesmiella novae-zelandiae* (voucher MFLU 18-1604, GenBank MN124526.1; Identities = 591/674 (88 %), no gaps), and *Sporidesmiella hyalosperma* (strain MFLUCC 18-1013, GenBank MW504070.1; Identities = 584/674 (87 %), no gaps). The *rpb2* sequences of CPC 41075 and CPC 41109 are identical (674/674 nucleotides). No significant hits were obtained when the **tef1** (first part) sequence of CPC 41109 was used in blastn and megablast searches.

Supplementary material

FP1304 Phylogenetic tree.

Xenopyricularia junci
& *Zaanenomyces quadripartis*



Fungal Planet 1305 & 1306 – 24 December 2021

Xenopyricularia junci Crous & Osieck, *sp. nov.*

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Pyriculariaceae*, *Magnaporthales*, *Sordariomycetes*.

Mycelium consisting of hyaline to subhyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. *Conidiophores* arising from superficial mycelium, erect, flexuous, dark brown, thick-walled, up to 25-septate, base somewhat swollen, 5–7 µm diam, pale brown, unbranched, up to 800 µm tall, 3.5–4.5 µm wide. *Conidiogenous cells* integrated, terminal and intercalary, forming a rachis of pimple-like denticles, 0.5–1 × 0.5–1 µm, slightly darkened, up to 100 µm long, pale brown, smooth-walled, 4.5–6 µm wide. *Conidia* in dry clusters, solitary, pale brown, guttulate, 2-septate, with middle cell darker brown, (18–)

27–30(–35) × (5–)6(–6.5) µm, pyriform to narrowly fusoid-ellipsoid, apex subobtuse, tapering toward truncate hilum, 1.5–2 µm diam, somewhat darkened.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, even margin, reaching 25 mm diam after 2 wk at 25 °C. On MEA, PDA and OA surface and reverse dirty white.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'45" E05°10'34", on dead culm of *Juncus effusus* (*Juncaceae*), 8 Jan. 2021, E.R. Osieck, HPC 3563 = WI-09/#4195 (holotype CBS H-24847, culture ex-type CPC 40968 = CBS 148308, ITS, LSU, *actA*, *cmdA* and *rpb1* sequences GenBank OK664720.1, OK663759.1, OK651127.1, OK651142.1 and OK651152.1, MycoBank MB 841803).

(notes *Xenopyricularia junci* on Supplementary material page FP1305 & 1306)

Supplementary material

FP1305 Phylogenetic tree.

Zaenenomyces Crous & Osieck, *gen. nov.*

Etymology. Named for Mieke Zaenen LL.M., who was the Director General of the Royal Dutch Academy of Arts and Sciences (Feb. 2015 – July 2021), and overall responsible for the day-to-day management of the Academy. During this period, she also oversaw the expansion of the research infrastructure and rebranding of the Centraalbureau voor Schimmelfungalcultures to the present Westerdijk Fungal Biodiversity Institute (WI).

Classification — *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate hyphae. *Conidiophores* solitary, arising from superficial mycelium, erect, subcylindrical, unbranched, brown, smooth-

walled, septate; base swollen or with rhizoids, conidiophore with or without brown nodulose swellings, thick-walled. *Conidiogenous cells* terminal, integrated, forming a rachis with pimple-like denticles, slightly thickened and darkened, not refractive. *Conidia* solitary, forming a dry mass at apex, narrowly obclavate, straight to flexuous, apex subobtuse, base obconically truncate, septate, hyaline, smooth-walled, guttulate, hilum thickened and darkened.

Type species. *Zaenenomyces quadripartis* Crous & Osieck
MycoBank MB 841804.

For *Zaenenomyces* (see also FP 1307 & 1313)

Zaenenomyces quadripartis Crous & Osieck, *sp. nov.*

Etymology. Name refers to the expansion of the WI research infrastructure (D-wing; *quadripartis* ('the fourth part'), during the Westerdijk year (2017), celebrating Johanna Westerdijk, who was the first female professor in the Netherlands (10 February 1917).

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. *Conidiophores* solitary, arising from superficial mycelium, erect, subcylindrical, unbranched, brown, smooth-walled, 1–4-septate; base swollen or with rhizoids, conidiophore with or without brown nodulose swellings, thick-walled, 15–100 × 2.5–3.5 µm. *Conidiogenous cells* terminal, integrated, forming a rachis with pimple-like denticles, slightly thickened and darkened, not refractive, 0.5 µm diam, 10–35 × 2.5–3.5 µm. *Conidia* solitary, forming a dry mass at apex, narrowly obclavate, straight to flexuous, apex subobtuse, base obconically truncate, (1–)4-septate, hyaline, smooth-walled, guttulate, hilum thickened and darkened, 0.5–1 µm

diam, (12–)23–30(–35) × (2.5–)3 µm; conidia becoming prominently constricted at septa with age.

Culture characteristics — Colonies erumpent, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 3 mm diam after 7 d at 25 °C. On MEA surface and reverse sienna; on PDA surface and reverse pale luteous; on OA surface pale luteous.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'45" E05°10'34", on dead culm of *Juncus effusus* (*Juncaceae*), 8 Jan. 2021, E.R. Osieck, HPC 3563 = WI-09/#4195 (holotype CBS H-24849, culture ex-type CPC 41073 = CBS 148310, ITS and LSU sequences GenBank OK664721.1 and OK663760.1, MycoBank MB 841805).

Additional material examined. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, on dead culm of *Juncus effusus* (*Juncaceae*), 8 Jan. 2021, E.R. Osieck, HPC 3565 = WI-11, CBS H-24818, culture CPC 41013 = CBS 148272, ITS and LSU sequences GenBank OK664722.1 and OK663761.1.


(notes *Zaenenomyces quadripartis* on Supplementary material page FP1305 & 1306)

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1290.

Colour illustrations. *Juncus effusus* at Nieuw Wulven, near Houten. Left column: *Xenopyricularia junci*. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Right column: *Zaenenomyces quadripartis*. Conidiophores developing on SNA; conidiogenous cells giving rise to conidia. Scale bars = 10 µm.

Zaanenomyces moderatricis-academiae



Fungal Planet 1307 – 24 December 2021

***Zaanenomyces moderatricis-academiae* Crous & Osieck, sp. nov.**

Etymology. Name refers to the role Mieke Zaanen LL.M. occupied in the Netherlands (Feb. 2015 – July 2021); '*moderatrix*' = female director, and '*academia*' = academy.

Classification — *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*.

Mycelium consisting of branched, septate, pale brown, smooth-walled, 1.5–2 µm diam hyphae. *Conidiophores* solitary, erect, subcylindrical, pale brown, smooth-walled, straight, unbranched, 1–2-septate, 7–30 × 2–2.5 µm. *Conidiogenous cells* integrated, terminal, pale brown, smooth-walled, holoblastic, at times sympodial, 5–15 × 2–2.5 µm; scars somewhat darkened, non-refractive. *Conidia* solitary, straight to slightly curved, narrowly obclavate to subcylindrical, widest in lowest third, pale brown, smooth-walled, guttulate, apex subobtuse, base truncate, 1.5–2 µm diam, not darkened, 4–10-septate, (32–)40–52(–57) × 2.5–3 µm.

Culture characteristics — Colonies erumpent, spreading, with sparse aerial mycelium and feathery, lobate margin, reaching 7 mm diam after 7 d at 25 °C. On MEA, PDA and OA surface and reverse olivaceous grey.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'44" E05°09'40", on dead culm of *Juncus inflexus* (*Juncaceae*), 4 Feb. 2021, *E.R. Osieck*, HPC 3590 = WI-21/#4205 (holotype CBS H-24854, culture ex-type CPC 41273 = CBS 148315, ITS, LSU and *rpb2* sequences GenBank OK664723.1, OK663762.1 and OK651167.1, MycoBank MB 841806).

Notes — *Zaanenomyces moderatricis-academiae* represents a second species of the genus *Zaanenomyces*, also collected from dead culms of *Juncus* spp. in the Netherlands (see FP 1306). The ex-type strain was cultured from minute, dark, setose ascomata (material too depauperate for description), suggesting that it has a sexual morph. Unfortunately, only the asexual morph developed in culture. *Zaanenomyces moderatricis-academiae* is distinguished from *Z. quadripartis* based on its conidiophore and conidium morphology; it is also phylogenetically distinct.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Acanthostigma perpusillum* (strain UAMH 7237, GenBank AY916492.1; Identities = 442/506 (87 %), 16 gaps (3 %)), *Camporesiomyces vaccinii* (strain CBS 216.90, GenBank NR_156202.1; Identities = 432/505 (86 %), 13 gaps (2 %)), and *Helicosporium flavisporum* (strain MFLUCC 17-2020, GenBank NR_160372.1; Identities = 409/472 (87 %), 28 gaps (5 %)). Closest hits using the **LSU** sequence are *Helicosporium luteosporum* (strain MFLUCC 16-0226, GenBank NG_059773.1; Identities = 765/793 (96 %), no gaps), *Helicosporium setiferum* (as *Helicosporium* sp. YZL-2018c; strain MFLUCC 17-2007, GenBank MH558863.1; Identities = 763/793 (96 %), no gaps), and *Helicosporium flavisporum* (strain MFLUCC 17-2020, GenBank NG_066235.1; Identities = 761/793 (96 %), no gaps). No significant hits were obtained when the **rpb2** was used in blastn and megablast searches.

Colour illustrations. *Juncus inflexus* growing at Nieuw Wulven, near Houten. Conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1290.



Fungal Planet 1308 – 24 December 2021

***Tetracoccosporium pseudoaerium* Crous & Jurjević, sp. nov.**

Etymology. Name refers to its morphological similarity to *Tetracocco-sporium aerium*.

Classification — *Sympoventuriaceae*, *Venturiales*, *Dothideo-mycetes*.

Mycelium consisting of pale brown, smooth-walled, septate, branched, 2.5–3 µm diam hyphae. **Conidiophores** terminal and lateral on hyphae, erect, subcylindrical, pale brown, smooth-walled, 1–2-septate, with 1–3 terminal conidiogenous cells, 7–25 × 2.5–3 µm. **Conidiogenous cells** pale brown, smooth, doliiform to subcylindrical, monoblastic, tapering toward apex, 1 µm diam, 3–10 × 2–3 µm. **Conidia** solitary, dry, acrogenous, spherical, medium brown, verruculose, becoming warty, divided by septa at right angles into four sections, becoming swollen with age and constricted at septa, 10–13 µm wide, 7–8 µm high.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 10 mm diam after 7 d at 25 °C. On MEA, PDA, Czapek yeast agar (CYA) and OA surface and reverse umber. On CYA reaching 14 mm diam after 7 d at 37 °C.

Typus. USA, Delaware, Wilmington, from air sample in house, Jan. 2010, Ž. Jurjević 5594 (holotype CBS H-24822, culture ex-type CPC 41065 = CBS 148276, ITS and LSU sequences GenBank OK664724.1 and OK663763.1, MycoBank MB 841807).

Notes — *Tetracoccosporium aerium* (ex-type CBS 493.92) was described to have conidia that are dark brown, 4-celled, 8–12 × 6–8 µm, constricted at septa. As is the case with CBS 148276, it was also isolated from air, but collected in India (Misra & Srivastava 1976). Morphologically the two isolates have overlapping conidial dimensions, but are phylogenetically distinct, and thus the isolate from the USA is herewith described as a new species.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Tetracoccosporium aerium* (strain CBS 493.92, GenBank NR_160189.1; Identities = 484/526 (92 %), 19 gaps (3 %)), *Ochroconis bacilliformis* (strain CBS 100442, GenBank NR_155566.1; Identities = 269/295 (91 %), 18 gaps (6 %)), and *Ochroconis musae* (strain CBS 145061, GenBank MK442605.1; Identities = 271/301 (90 %), 12 gaps (3 %)). Closest hits using the **LSU** sequence are *Tetracoccosporium aerium* (strain CBS 493.92, GenBank MH874035.1; Identities = 796/812 (98 %), three gaps (0 %)), *Melnikomyces vietnamensis* (strain CBS 136209, GenBank NG_058087.1; Identities = 780/813 (96 %), one gap (0 %)), and *Melnikomyces thailandicus* (strain CBS 145767, GenBank NG_068685.1; Identities = 741/774 (96 %), one gap (0 %)).

Colour illustrations. House at Wilmington, Delaware, USA. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1290.



Fungal Planet 1309 – 24 December 2021

***Myrmecridium junci* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Myrmecridiaceae*, *Myrmecridiales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 2–3 µm diam hyphae. **Conidiophores** solitary, erect, subcylindrical, medium brown, smooth-walled, unbranched, straight to flexuous, septate, 45–200 × 3–4 µm. **Conidiogenous cells** integrated, terminal, and intercalary, subcylindrical, pale brown, smooth-walled, forming a rachis in upper part with numerous pimple-like denticles, 0.5–1 × 0.5 µm, somewhat refractive; conidiophore has whorls of conidia along its length, 7–80 × 3–3.5 µm. **Conidia** aseptate, solitary, obovoid, hyaline, pale olivaceous with age, apex obtuse, tapering to truncate hilum, 1 µm diam, aseptate, with mucoid sheath surrounding median region, 1–2 µm diam, (8–)9–10(–11) × 2.5–3 µm.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 15 mm diam after 7 d at 25 °C. On MEA surface orange, reverse scarlet; on PDA surface and reverse pale luteous; on OA surface orange.

Typus. NETHERLANDS, Utrecht Province, Bornia, near Driebergen, 8 m a.s.l., N52°02'23" E05°17'56", on dead culm of *Juncus effusus* (*Juncaceae*), 25 Jan. 2021, *E.R. Osieck*, HPC 3583 = WI-15 (holotype CBS H-24820, culture ex-type CPC 41032 = CBS 148274, ITS and LSU sequences GenBank OK664725.1 and OK663764.1, MycoBank MB 841808).

Notes — *Myrmecridium junci* has conidia of similar dimensions to those of *M. sambuci* ((7–)8–9(–10) × (2.5–)3(–3.5) µm), but is distinct in that conidia are arranged in distinct whorls along the length of the conidiophore, which is not the case with *M. sambuci*. The species is distinct from all other *Myrmecridium* species known from DNA data.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Myrmecridium schulzeri* (strain CSB_F134, GenBank KU574706.1; Identities = 522/531 (98 %), four gaps (0 %)), *Myrmecridium flexuosum* (strain CBS 398.76, GenBank NR_146238.1; Identities = 511/522 (98 %), three gaps (0 %)), and *Myrmecridium fluviae* (strain CNUFC YR61-1, GenBank NR_164555.1; Identities = 461/471 (98 %), one gap (0 %)). Closest hits using the **LSU** sequence are *Myrmecridium schulzeri* (strain CBS 114996, GenBank EU041835.1; Identities = 780/790 (99 %), no gaps), *Myrmecridium banksiae* (strain CBS 132536, GenBank NG_042684.1; Identities = 779/790 (99 %), no gaps), and *Myrmecridium spartii* (strain CPC 24953, GenBank KR611902.1; Identities = 778/790 (98 %), no gaps).

Colour illustrations. *Juncus effusus* at Bornia, near Driebergen. Conidiophores on SNA; conidiophores and conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.


Supplementary material

See the phylogenetic tree provided with the supplementary material FP1293.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Eduard R. Osieck, Jkvr. C.M. van Asch van Wijcklaan 19, 3972 ST Driebergen-Rijnsburg, Netherlands; e-mail: panurus@ziggo.nl

Caeliomyces tampanus



Fungal Planet 1310 – 24 December 2021

Caeliomyces* Crous & Jurjević, gen. nov.Etymology.* Name refers to the fact that it was isolated from air 'caeli'.Classification — *Teratosphaeriaceae*, *Mycosphaerellales*, *Dothideomycetes*.*Conidiomata* solitary, pycnidial, brown, globose, with central ostiole; wall of 3–6 layers of brown *textura angularis*. *Conidio-**phores* reduced to conidiogenous cells lining the inner cavity, hyaline, smooth-walled, ampulliform, phialidic. *Conidia* solitary, aseptate, ellipsoid, ends obtuse, green-brown, smooth-walled.*Type species.* *Caeliomyces tampanus* Crous & Jurjević
Mycobank MB 841809.***Caeliomyces tampanus* Crous & Jurjević, sp. nov.***Etymology.* Name refers to the collection site, Tampa, Florida.*Conidiomata* solitary, pycnidial, brown, globose, 20–80 µm diam, with central ostiole, 3–6 µm diam; wall of 3–6 layers of brown *textura angularis*. *Conidiophores* reduced to conidiogenous cells lining the inner cavity, hyaline, smooth-walled, ampulliform, 3–6 × 2–3 µm, phialidic. *Conidia* solitary, aseptate, ellipsoid, ends obtuse, green-brown, smooth-walled, 2–4 × 1.5–2 µm.

Culture characteristics — Colonies flat, spreading, with sparse to moderate aerial mycelium and smooth, lobate margin, reaching 10–20 mm diam after 2 wk at 25 °C. On MEA surface and reverse olivaceous grey; on PDA and OA surface and reverse iron grey.

Typus. USA, Florida, Tampa, from office dust, Oct. 2018, Ž. Jurjević 5589 (holotype CBS H-24821, culture ex-type CPC 41060 = CBS 148275, ITS, LSU, *cmdA*, *tef1* (first part) and *tub2* sequences GenBank OK664726.1, OK663765.1, OK651143.1, OK651189.1 and OK651207.1, MycoBank MB 841810).Notes — *Caeliomyces* is phylogenetically closely related to coelomycetous genera such as *Caatingomyces* (Hyde et al. 2019) and *Readeriella* (Crous et al. 2019c). All three genera have pycnidial conidiomata, phialidic conidiogenous cells, and aseptate, pigmented conidia. However, *Readeriella* has conidia that tend to vary in shape, and have a thicker wall, darker pigmentation, and percurrently proliferating conidiogenous cells, while *Caatingomyces* has septate, branched conidiophores with terminal and intercalary conidiogenous cells.Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Readeriella deanei* (strain CBS 134746, GenBank MH866077.1; Identities = 416/448 (93 %), ten gaps (2 %)), *Caatingomyces brasiliensis* (as *Capnodiales* sp. JB-2019a; strain URM 7916, GenBank MH929437.1; Identities = 460/497 (93 %), 11 gaps (2 %)), and *Readeriella angustia* (strain CBS 124997, GenBank NR_156531.1; Identities = 491/533 (92 %), 14 gaps (2 %)). Closest hits using the LSU sequence are *Readeriella angustia* (strain CBS 124998, GenBank NG_057843.1; Identities = 801/817 (98 %), four gaps (0 %)), *Caatingomyces brasiliensis* (as *Capnodiales* sp. JB-2019a; strain URM 7916, GenBank MH929439.1; Identities = 799/815 (98 %), five gaps (0 %)), and *Readeriella dimorphospora* (strain CPC 25379, GenBank KX306792.1; Identities = 763/779 (98 %), four gaps (0 %)). Closest hits using the *cmdA* sequence had highest similarity to *Readeriella tasmanica* (strain CPC 13631, GenBank KF902686.1; Identities = 283/311 (91 %), no gaps), *Readeriella dendritica* (strain CBS 120032, GenBank KF902674.1; Identities = 282/310 (91 %), no gaps), and *Readeriella novaezealandiae* (strain CBS 114357, GenBank KF902683.1; Identities = 280/309 (91 %), no gaps). No significant hits were obtained when the *tef1* sequence was used in blastn and megablast searches. Closest hits using the *tub2* sequence had highest similarity to *Readeriella ellipsoidea* (strain CBS 142525, GenBank KY979925.1; Identities = 302/362 (83 %), 24 gaps (6 %)), *Readeriella lehmannii* (strain CPC 28935, GenBank KY173617.1; Identities = 301/361 (83 %), 22 gaps (6 %)), and *Readeriella menaiensis* (strain CBS 125003, GenBank KF442490.1; Identities = 291/352 (83 %), 17 gaps (4 %)).*Colour illustrations.* Office at Tampa, Florida. *Conidiomata* on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 80 µm (*conidiomata*), 10 µm (all others).**Supplementary material**

See the phylogenetic tree provided with the supplementary material FP1298.



Fungal Planet 1311 – 24 December 2021

***Microcera physciae* Crous & Boers, sp. nov.**

Etymology. Name refers to the host genus *Physcia* from which it was isolated.

Classification — *Nectriaceae*, *Hypocreales*, *Sordariomycetes*.

Associated with discoloured thalli of *Physcia tenella*. Mycelium consisting of hyaline, smooth-walled, branched, septate, 2–3 µm diam hyphae. Conidiomata sporodochial, pale cream on SNA, consisting of tightly aggregated conidiophores that are reduced to conidiogenous cells or with a short stipe, giving rise to 1–4 conidiogenous cells. Conidiogenous cells subulate, tapering towards apex, flexuous, monophialidic with periclinal thickening and minute non-flared collarette, 10–20 × 2–3 µm. Conidia hyaline, smooth-walled, guttulate, (1–)3-septate, distinctly curved and twisted, apex attenuated, subobtuse, dorsiventrally curved, basal cell attenuated to subobtuse, non foot-shaped to slightly notched basal cell, (20–)24–26(–33) × 3(–3.5) µm.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 15 mm diam after 7 d at 25 °C. On MEA surface orange, reverse orange with patches of ochreous; on PDA surface and reverse orange; on OA surface ochreous with orange to creamy sporodochia.

Typus. NETHERLANDS, Drenthe Province, Dwingeloo graveyard, 52.831376, 6.365404, on *Physcia tenella* (*Physciaceae*), 10 Mar. 2021, J. Boers, HPC 3602 (holotype CBS H-24829, culture ex-type CPC 41284 = CBS 148283, ITS, LSU, *rpb1*, *rpb2*, *tef1* (first part) and *tub2* sequences GenBank OK664727.1, OK663766.1, OK651153.1, OK651168.1, OK651190.1 and OK651208.1, MycoBank MB 841811).

Additional material examined. NETHERLANDS, Drenthe Province, Dwingeloo, 52.835821, 6.364449, on *Physcia tenella*, 10 Feb. 2021, J. Boers, HPC 3585, CBS H-24834, culture CPC 41038 = CBS 148288, ITS, LSU, *rpb1*, *rpb2*, *tef1* (first part) and *tub2* sequences GenBank OK664728.1, OK663767.1, OK651154.1, OK651169.1, OK651191.1 and OK651209.1.

Notes — Species of *Microcera* are usually pathogens of scale insects (Gräfenhan et al. 2011, Crous et al. 2021b). It is thus surprising to find a species associated with discoloured thalli of *Physcia tenella*. Although no insects were observed at the time of collection, it could well be that this element was overlooked, and further observations would be needed to resolve the ecology of *M. physciae*. The present collections should be compared with *Fusarium sampaii*, which was described from various lichens including *Physcia* '*leptalea*'. The latter species was discussed by Hawksworth (1979), who concluded that the *Fusarium* name should be rejected since the studied material was based on discordant elements. *Microcera physciae* is related to sequences identified as *M. coccophila*.

Colour illustrations. *Physcia tenella*. Sporodochia on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence of CPC 41284 had highest similarity to *Microcera larvarum* (strain ICMP 5444, GenBank MT107902.1; Identities = 486/498 (98 %), three gaps (0 %)), *Cosmospora aurantiicola* (strain F-251,305, GenBank EU860061.1; Identities = 486/498 (98 %), three gaps (0 %)), and *Microcera rubra* (strain CBS 638.76, GenBank MH861019.1; Identities = 485/498 (97 %), three gaps (0 %)). The ITS sequence of CPC 41038 differs with two nucleotides from that of CPC 41284 (495/497 (99 %), no gaps). Closest hits using the LSU sequence of CPC 41284 are *Microcera rubra* (strain CBS 638.76, GenBank NG_058100.1; Identities = 803/814 (99 %), no gaps), *Microcera larvarum* (strain CBS 738.79, GenBank KM231701.1; Identities = 803/814 (99 %), no gaps), and *Microcera coccophila* (strain MAFF 241482, GenBank KC291787.1; Identities = 780/793 (98 %), no gaps). The LSU sequence of CPC 41038 is identical to that of CPC 41284 (813/813 (100 %), no gaps). Closest hits using the *rpb1* sequence of CPC 41284 had highest similarity to *Microcera coccophila* (strain MAFF 241482, GenBank KC291895.1; Identities = 629/688 (91 %), one gap (0 %)), *Microcera larvarum* (strain A.R. 4580, GenBank KC291894.1; Identities = 627/688 (91 %), one gap (0 %)), and *Microcera rubra* (strain CBS 638.76, GenBank KM232253.1; Identities = 658/730 (90 %), one gap (0 %)). The *rpb1* sequences of CPC 41038 and 41230 differ with two and 29 nucleotides from that of CPC 41284 (740/742 (99 %), one gap, and 709/738 (96 %), one gap, respectively). Closest hits using the *rpb2* sequence of CPC 41284 had highest similarity to *Microcera* sp. (strain NRRL 26790, GenBank JX171636.1; Identities = 843/850 (99 %), no gaps), *Microcera larvarum* (strain NRRL 20473, GenBank JX171587.1; Identities = 764/848 (90 %), no gaps), and *Microcera rubra* (strain CBS 638.76, GenBank KM232388.1; Identities = 761/845 (90 %), no gaps). The *rpb2* sequence of CPC 41038 differs with five nucleotides from that of CPC 41284 (870/875 (99 %), no gaps). Closest hits using the *tef1* (first part) sequence of CPC 41284 had highest similarity to '*Fusarium* sp.' (strain NRRL 20473, GenBank JF740695.1; Identities = 397/479 (83 %), 19 gaps (3 %)), *Microcera larvarum* (strain CBS 738.79, GenBank KM231957.1; Identities = 388/479 (81 %), 26 gaps (5 %)), and *Microcera coccophila* (strain ZJUP0179, GenBank MN614420.1; Identities = 346/426 (81 %), 21 gaps (4 %)). The *tef1* (first part) sequence of CPC 41038 differs with 15 nucleotides from that of CPC 41284 (455/470 (97 %), one gap). Closest hits using the *tub2* sequence of CPC 41284 had highest similarity to '*Fusarium* sp.' (strain F-267,620, GenBank EU860029.1; Identities = 521/526 (99 %), one gap (0 %)), *Microcera larvarum* (as *Fusarium larvarum* var. *larvarum*; strain F-266,784, GenBank EU860024.1; Identities = 455/522 (87 %), five gaps (0 %)), and *Microcera rubra* (as *Fusarium larvarum* var. *rubrum*; strain F-257,517, GenBank EU860021.1; Identities = 454/522 (87 %), five gaps (0 %)). The *tub2* sequence of CPC 41038 differs with three nucleotides from that of CPC 41284 (343/346 (99 %), no gaps).

Supplementary material

FP1311 Phylogenetic tree.

Myrmecridium dactylidis
& *Zaanenomyces versatilis*



Fungal Planet 1312 & 1313 – 24 December 2021

***Myrmecridium dactylidis* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Dactylis* from which it was isolated.

Classification — *Myrmecridiaceae*, *Myrmecridiales*, *Sordariomycetes*.

Mycelium consisting of branched, septate, hyaline, smooth-walled, 1–2 µm diam hyphae. **Conidiophores** arising from aerial hyphae, unbranched, straight, medium brown, thick-walled, 1–4-septate, up to 100 µm tall, 3–4 µm diam; basal cell not inflated. **Conidiogenous cells** integrated, cylindrical, 20–35 × 3.5–4 µm, pale brown, forming a rachis with scattered pimple-shaped denticles less than 1 µm long and ± 0.5 µm wide, apically pointed, pigmented, slightly thickened. **Conidia** solitary, 0–1(–2)-septate, subhyaline, thin-walled, smooth, guttulate, surrounded by a wing-like gelatinous sheath, ± 0.5 µm thick, ellipsoid to obovoid or fusoid, (6–)8–10(–11.5) × (3–)3.5(–4) µm, tapering to a subtruncate hilum; hilum un-pigmented, not darkened.

Culture characteristics — Colonies flat, spreading, with sparse to moderate aerial mycelium and smooth, lobate margin, reaching 12 mm diam after 2 wk at 25 °C. On MEA, PDA and OA surface and reverse orange.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°03'04" E05°09'48", on dead culms of *Dactylis glomerata*

(*Poaceae*), 4 Feb. 2021, E.R. Osieck, HPC 3601 = WI-27/#4222 (holotype CBS H-24827, culture ex-type CPC 41247 = CBS 148281, ITS and LSU sequences GenBank OK664729.1 and OK663768.1, MycoBank MB 841812).

Notes — *Myrmecridium dactylidis* is closely related to *M. phragmites* (conidia 0–1-septate, (6.5–)7–8(–9) × (2.5–)3(–3.5) µm; Crous et al. 2011), but distinct in that it has larger conidia that can be up to 2-septate.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Myrmecridium phragmitis* (strain CBS 131311, GenBank NR_137782.1; Identities = 542/552 (98 %), four gaps (0 %)), *Myrmecridium schulzeri* (strain 9Y-G52, GenBank MT138583.1; Identities = 514/527 (98 %), two gaps (0 %)), and *Myrmecridium phragmiticola* (strain CPC 36367, GenBank NR_170826.1; Identities = 527/549 (96 %), four gaps (0 %)). Closest hits using the **LSU** sequence are *Myrmecridium phragmitis* (strain CBS 131311, GenBank NG_057948.1; Identities = 796/798 (99 %), no gaps), *Myrmecridium schulzeri* (strain DUCC4137, GenBank MH819185.1; Identities = 784/798 (98 %), no gaps), and *Myrmecridium banksiae* (strain CBS 132536, GenBank NG_042684.1; Identities = 784/798 (98 %), no gaps).

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1293.

***Zaanenomyces versatilis* Crous & Osieck, sp. nov.**

Etymology. Name refers to the versatile role Mieke Zaanen LL.M. occupied as Director General of the Royal Dutch Academy of Arts and Sciences (Feb. 2015 – July 2021).

Classification — *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*.

Mycelium consisting of pale brown, smooth-walled, branched, septate, 2–2.5 µm diam hyphae. **Conidiophores** solitary, erect, subcylindrical, straight to geniculous-sinuuous, unbranched, 0–5-septate, 5–60 × 2.5–3.5 µm. **Conidiogenous cells** terminal, integrated, pale brown, smooth-walled, subcylindrical with apical rachis of denticles, cylindrical, 1–2 × 1–1.5 µm; scars not thickened nor darkened. **Conidia** solitary, in dry clusters, subcylindrical, widest in middle, tapering to subobtuse apex and truncate hilum, 1.5–2 µm diam, pale brown, smooth, guttulate, straight to slightly flexuous, (3–)7–10(–12)-septate, (16–)43–50(–55) × (2.5–)3(–3.5) µm.

Culture characteristics — Colonies erumpent, spreading, surface folded, with moderate aerial mycelium and smooth, lobate margin, reaching 6 mm diam after 7 d at 25 °C. On MEA, PDA and OA surface umber, reverse chestnut.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'44" E05°09'40", on dead culm of *Juncus inflexus* (*Juncaceae*), 4 Feb. 2021, E.R. Osieck, HPC 3590 = WI-21/#4205 (holotype CBS H-24851,

culture ex-type CPC 41224 = CBS 148312, ITS and LSU sequences GenBank OK664730.1 and OK663769.1, MycoBank MB 841813).

Notes — *Zaanenomyces versatilis* represents the third species in the genus (see FP 1306 & 1307) collected on *Juncus* in the Netherlands, distinct from *Z. quadripartis* (conidia (1–)4-septate, (12–)23–30(–35) × (2.5–)3 µm), and *Z. moderatricis-academiae* (conidia 4–10-septate, (32–)40–52(–57) × 2.5–3 µm).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Acanthostigma perpusillum* (strain UAMH 7237, GenBank AY916492.1; Identities = 504/568 (89 %), 16 gaps (2 %)), *Camporesiomyces vaccinii* (strain CBS 216.90, GenBank NR_156202.1; Identities = 498/568 (88 %), 15 gaps (2 %)), and *Tubeufia cerea* (strain BCC 8125, GenBank AY916491.1; Identities = 493/568 (87 %), 30 gaps (5 %)). The ITS sequences of CPC 41273 (*Zaanenomyces juncicola*) and CPC 41224 are dissimilar (484/505 (96 %), six gaps (1 %)). Closest hits using the **LSU** sequence are *Helicosporium luteosporum* (strain MFLU 16-2871, GenBank NG_059773.1; Identities = 794/818 (97 %), no gaps), *Helicosporium flavisporum* (voucher MFLU 17-1105, GenBank NG_066235.1; Identities = 790/818 (97 %), no gaps), and *Helicosporium vesicarium* (as *Helicosporium* sp. YZL-2018e; strain MFLUCC 17-1795, GenBank MH558864.1; Identities = 790/818 (97 %), no gaps). The LSU sequences of CPC 41273 (*Zaanenomyces juncicola*) and CPC 41224 are dissimilar (789/793 (99 %), no gaps).

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1290.

Colour illustrations. *Juncus inflexus* at Nieuw Wulven, near Houten. Left column: *Myrmecridium dactylidis*. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Right column: *Zaanenomyces versatilis*. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Eduard R. Osieck, Jkvr. C.M. van Asch van Wijcklaan 19, 3972 ST Driebergen-Rijsenburg, Netherlands; e-mail: panurus@ziggo.nl



Fungal Planet 1314 – 24 December 2021

***Myrmecridium juncicola* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Myrmecridiaceae*, *Myrmecridiales*, *Sordariomycetes*.

Mycelium consisting of pale brown, smooth-walled, branched, septate, 2.5–3 µm diam hyphae. **Conidiophores** solitary, erect, unbranched, subcylindrical, flexuous, brown, smooth- and thick-walled, 1–10-septate, up to 170 µm tall, 4–5 µm wide, arising from superficial mycelium. **Conidiogenous cells** integrated, terminal, subcylindrical, pale brown, smooth-walled, 20–35 × 3.5–4 µm, forming a rachis with numerous pimple-like denticles, 0.5–1 × 0.5 µm, slightly darkened, not reflective. **Conidia** solitary, fusoid, pale brown, smooth-walled, guttulate with mucoid sheath in mid region (up to 2 µm diam), apex subobtuse, tapering to truncate hilum, 0.5 µm diam, slightly darkened, 0–1-septate, (12–)14–16(–18) × (2.5–)3(–3.5) µm.

Culture characteristics — Colonies flat, spreading, with sparse to moderate aerial mycelium and smooth, lobate margin, reaching 20 mm diam after 7 d at 25 °C. On MEA surface and reverse ochreous; on PDA surface and reverse saffron; on OA surface saffron with patches of umber.

Typus. NETHERLANDS, Overijssel Province, Engbertsdijkswenen, near Vriezenveen, 13 m a.s.l., N52°26'53" E06°40'02", on dead culm of *Juncus effusus* (*Juncaceae*), 9 Mar. 2021, *E.R. Osieck*, HPC 3612 = WI-31/#4227 (holotype CBS H-24855, culture ex-type CPC 41313 = CBS 148316, ITS and LSU sequences GenBank OK664731.1 and OK663770.1, MycoBank MB 841814).

Additional materials examined. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'45" E05°10'34", on dead culm of *J. effusus*, 8 Jan. 2021, *E.R. Osieck*, HPC 3565 = WI-11/#4195, CBS H-24813, culture CPC 40831 = CBS 148267, ITS and LSU sequences GenBank OK664733.1 and OK663772.1; *ibid.*, CBS H-24812, culture CPC 40829 = CBS 148266, ITS and LSU sequences GenBank OK664732.1 and OK663771.1.

Colour illustrations. *Juncus effusus* at Engbertsdijkswenen, near Vriezenveen, Overijssel Province. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Notes — *Myrmecridium juncicola* needs to be compared to *M. junci* (conidia (8–)9–10(–11) × 2.5–3 µm, arranged in whorls; described elsewhere in this paper), but has larger conidia, also when compared to other species treated here, like *M. dactylidis*, and *M. sambuci*. Phylogenetically, the species is related to *M. dactylidis*, and *M. phragmitis*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence of CPC 41313 had highest similarity to *Myrmecridium schulzeri* (strain CSB_F134, GenBank KU574706.1; Identities = 484/505 (96 %), seven gaps (1 %)), *Myrmecridium phragmiticola* (strain CPC 36367, GenBank NR_170826.1; Identities = 480/505 (95 %), seven gaps (1 %)), and *Myrmecridium flexuosum* (strain CBS 398.76, GenBank NR_146238.1; Identities = 473/501 (94 %), ten gaps (1 %)). The ITS sequences of CPC 40829 and 40831 differ with one nucleotide each from that of CPC 41313 (499/500 (99 %), no gaps, and 499/500 (99 %), no gaps, respectively). Closest hits using the **LSU** sequence of CPC 41313 are *Myrmecridium banksiae* (strain CBS 132536, GenBank NG_042684.1; Identities = 804/813 (99 %), two gaps (0 %)), *Myrmecridium spartii* (strain CPC 24953, GenBank KR611902.1; Identities = 803/813 (99 %), two gaps (0 %)), and *Myrmecridium schulzeri* (strain CBS 114996, GenBank EU041835.1; Identities = 803/813 (99 %), two gaps (0 %)). The LSU sequences of CPC 40829 and 40831 are identical to that of CPC 41313 (811/811 (100 %), no gaps, and 788/788 (100 %), no gaps, respectively).

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1293.



Fungal Planet 1315 – 24 December 2021

***Sarocladium junci* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Sarocladiaceae*, *Hypocreales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** arising from superficial mycelium, erect, flexuous, rarely branched, subcylindrical with apical taper, hyaline, smooth-walled, 1–3-septate, 15–60 × 2–3 µm. **Conidiogenous cells** integrated, terminal rarely intercalary, 15–30 × 2–3 µm, monophialidic, with periclinal thickening; conidia aggregating in mucoid mass. **Conidia** solitary, hyaline, smooth-walled, aseptate, subcylindrical, tapering to subobtusate ends, (3.5–)4–5 × 1.5–2 µm. **Chlamydospores** or **sexual morph** not observed.

Culture characteristics — Colonies flat, spreading, surface folded, with sparse aerial mycelium and smooth, lobate margin, reaching 35 mm diam after 7 d at 25 °C. On MEA surface dirty white, reverse pale luteous; on PDA and OA surface and reverse dirty white.


Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'44" E05°09'40", on dead culm of *Juncus inflexus* (*Juncaceae*), 4 Feb. 2021, E.R. Osieck, HPC 3590 = WI-21/#4205 (holotype CBS H-24823, culture ex-type CPC 41107 = CBS 148277, ITS, LSU, *actA*, *rpb2* and *tub2* sequences GenBank OK664734.1, OK663773.1, OK651128.1, OK651170.1 and OK651210.1, MycoBank MB 841815).


Notes — *Sarocladium* is an acremonium-like genus that contains several important plant and human pathogens (Summerbell et al. 2011, Giraldo et al. 2015). *Sarocladium junci* is phylogenetically and morphologically distinct from known species in the genus, including its close relative *S. spirale* (conidia 2.5–4 × 1–2 µm, colonies pale yellowish on MEA and PDA; Ou et al. 2020).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Sarocladium spirale* (strain BCRC FU31117, GenBank NR_165864.1; Identities = 510/517 (99 %), one gap (0 %)), *Sarocladium hominis* (strain UTHSC 04-1034, GenBank NR_155779.1; Identities = 544/560 (97 %), five gaps (0 %)), and *Sarocladium kiliense* (strain R3PS(A), GenBank MK752433.1; Identities = 550/572 (96 %), four gaps (0 %)). Closest hits using the **LSU** sequence are *Sarocladium oryzae* (strain CBS 180.74, GenBank NG_057866.1; Identities = 801/814 (98 %), no gaps), *Sarocladium kiliense* (strain CGMCC3.19018, GenBank MH380001.1; Identities = 802/816 (98 %), two gaps (0 %)), and *Sarocladium strictum* (strain CBS 376.701, GenBank MH871473.1; Identities = 802/816 (98 %), two gaps (0 %)). Closest hits using the **actA** sequence had highest similarity to *Sarocladium kiliense* (strain CBS 400.52, GenBank KM231258.1; Identities = 594/658 (90 %), seven gaps (1 %)), *Sarocladium sasijaorum* (strain NL19100007, GenBank MW890032.1; Identities = 584/655 (89 %), 13 gaps (1 %)), and *Sarocladium dejongiae* (strain CBS 144929, GenBank MK069412.1; Identities = 578/647 (89 %), 12 gaps (1 %)). Closest hits using the **rpb2** sequence had highest similarity to *Sarocladium kiliense* (strain CBS 400.52, GenBank KM232425.1; Identities = 717/842 (85 %), six gaps (0 %)), *Furcaterigmium furcatum* (as *Acremonium furcatum*; strain CBS 145322, GenBank LR594790.1; Identities = 344/437 (79 %), ten gaps (2 %)), and *Flavocillium bifurcatum* (strain YFCC 6101, GenBank MN576897.1; Identities = 408/528 (77 %), 20 gaps (3 %)). Closest hits using the **tub2** sequence had highest similarity to *Sarocladium spirale* (strain 3-22, GenBank LC464483.1; Identities = 562/598 (94 %), four gaps (0 %)), *Sarocladium kiliense* (strain CBS 400.52, GenBank KM232119.1; Identities = 522/607 (86 %), 32 gaps (5 %)), and *Sarocladium strictum* (strain AS5144, GenBank AY435440.1; Identities = 514/608 (85 %), 32 gaps (5 %)).

The first of two equally most parsimonious trees obtained from a phylogenetic analysis of the *Sarocladium tub2* nucleotide alignment. The tree was rooted to *Parasarocladium wereldwysianum* (culture NL_19095011; GenBank MW890143.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and bold text. GenBank accession (superscript) and/or culture collection/voucher numbers (in bold face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 17 strains including the outgroup; 647 characters including alignment gaps analysed: 293 constant, 147 variable and parsimony-uninformative and 207 parsimony-informative. Tree statistics: Tree Length = 620, Consistency Index = 0.831, Retention Index = 0.865, Rescaled Consistency Index = 0.718. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Colour illustrations. *Juncus inflexus* at Nieuw Wulven, near Houten. Conidiophores on SNA; conidiophores giving rise to conidia; conidia. Scale bars = 10 µm.





Fungal Planet 1316 – 24 December 2021

***Myrmecridium juncigenum* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Myrmecridiaceae*, *Myrmecridiales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2.5 µm diam hyphae, frequently forming hyphal coils. **Conidiophores** solitary, erect, arising from superficial mycelium, subcylindrical, straight to geniculate-sinuous, brown, smooth-walled, unbranched, 1–2-septate, 25–60 × 3–4 µm. **Conidiogenous cells** integrated, terminal, pale brown, smooth-walled, subcylindrical, forming a rachis of pimple-like subdenticulate loci, 0.5 µm diam, 25–45 × 3–4 µm. **Conidia** solitary, in dry clusters, subcylindrical, pale brown, guttulate, tapering towards subobtuse apex and truncate hilum, 0.5 µm diam, slightly darkened, middle region with mucoid sheath, 1–2 µm diam, 3-septate, (17–)20–22(–24) × (2.5–)3 µm.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 15 mm diam after 7 d at 25 °C. On MEA, PDA and OA surface and reverse orange.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'45" E05°10'34", on dead culm of *Juncus effusus* (*Juncaceae*), 8 Jan. 2021, *E.R. Osieck*, HPC 3564 = WI-10/#4195 (holotype CBS H-24814, culture ex-type CPC 40970 = CBS 148268, ITS and LSU sequences GenBank OK664735.1 and OK663774.1, MycoBank MB 841816).

Notes — *Myrmecridium juncigenum* (conidia 3-septate, (17–)20–22(–24) × (2.5–)3 µm) is morphologically distinct from other species known from this host like *M. juncicola* (conidia 0–1-septate, (12–)14–16(–18) × (2.5–)3(–3.5) µm) and *M. junci* (conidia aseptate, (8–)9–10(–11) × 2.5–3 µm, arranged in whorls). Furthermore, it is also phylogenetically distinct from other species known in the genus.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Myrmecridium hiemale* (strain A64, GenBank MF614998.1; Identities = 506/515 (98 %), one gap (0 %)), *Myrmecridium spartii* (strain CBS 140006, GenBank NR_155376.1; Identities = 533/544 (98 %), four gaps (0 %)), and *Myrmecridium schulzeri* (strain CK1477, GenBank MH474102.1; Identities = 517/528 (98 %), four gaps (0 %)). Closest hits using the **LSU** sequence are *Myrmecridium spartii* (strain CPC 24953, GenBank KR611902.1; Identities = 811/813 (99 %), no gaps), *Myrmecridium banksiae* (strain CBS 132536, GenBank NG_042684.1; Identities = 814/817 (99 %), one gap (0 %)), and *Myrmecridium schulzeri* (strain CBS 114996, GenBank EU041835.1; Identities = 813/817 (99 %), one gap (0 %)).


Colour illustrations. *Juncus effusus* at Nieuw Wulven, near Houten. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia with mucoid sheath. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1293.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Eduard R. Osieck, Jkvr. C.M. van Asch van Wijcklaan 19, 3972 ST Driebergen-Rijsenburg, Netherlands; e-mail: panurus@ziggo.nl



Fungal Planet 1317 & 1318 – 24 December 2021

***Cylindromonium everniae* Crous & Boers, sp. nov.**

Etymology. Name refers to the host genus *Evernia* from which it was isolated.

Classification — *Nectriaceae*, *Hypocreales*, *Sordariomycetes*.


Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** solitary, erect, flexuous, hyaline, smooth-walled, subcylindrical, unbranched, 1(–2)-septate, 40–100 × 2.5–3 µm. **Conidiogenous cells** integrated, terminal, subcylindrical with apical taper, hyaline, smooth-walled, 35–55 × 2.5–3 µm, monophialidic with periclinal thickening and non-flared collarette, 1.5–2 µm diam. **Conidia** solitary, aseptate, aggregating in mucoid mass, hyaline, smooth-walled, guttulate, subcylindrical, tapering to subobtuse apex and truncate, protruding hilum, 1 µm diam, (8–)12–14(–19) × (3–)3.5(–4) µm.

The first of two equally most parsimonious trees obtained from a phylogenetic analysis of the *Cylindromonium* ITS nucleotide alignment. The tree was rooted to *Neocosmospora citricola* (culture CBS 146513; GenBank NR_172994.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and **bold** text. GenBank accession (superscript) and/or culture collection/voucher numbers (in **bold** face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: eight strains including the outgroup; 527 characters including alignment gaps analysed: 361 constant, 109 variable and parsimony-uninformative and 57 parsimony-informative. Tree statistics: Tree Length = 240, Consistency Index = 0.896, Retention Index = 0.725, Rescaled Consistency Index = 0.650. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Culture characteristics — Colonies flat, spreading, surface folded, with sparse aerial mycelium and smooth, lobate margin, reaching 15 mm diam after 2 wk at 25 °C. On MEA surface pale luteous, reverse ochreous; on PDA surface pale luteous, reverse plate luteous with diffuse luteous pigment; on OA surface pale luteous.

Typus. NETHERLANDS, Drenthe Province, Dwingeloo graveyard, on *Evernia prunastri* (*Parmeliaceae*), Mar. 2021, J. Boers, HPC 3557 (holotype CBS H-24801, culture ex-type CPC 40760 = CBS 148255, ITS, LSU, *actA*, *his3*, *rpb2* and *tef1* (first part) sequences GenBank OK664736.1, OK663775.1, OK651129.1, OK651149.1, OK651171.1 and OK651192.1, MycoBank MB 841817).

(notes *Cylindromonium everniae* on Supplementary material page FP1317 & 1318)

**Supplementary material**

Also see the phylogenetic tree provided with the supplementary material FP1284.

***Rhodoveronaea everniae* Crous & Boers, sp. nov.**

Etymology. Name refers to the host genus *Evernia* from which it was isolated.

Classification — *Rhamphoriaceae*, *Rhamphoriales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** erect, arising from superficial mycelium, subcylindrical, branched or not, red-brown, thick-walled, warty, 1–6-septate, straight to geniculate-sinuuous, 25–100 × 3–4 µm. **Conidiogenous cells** integrated, terminal and intercalary, pale brown, smooth to warty, 15–40 × 3–4 µm, forming a rachis of minute pimple-like subdenticulate loci, 0.5 µm diam. **Conidia** solitary, pale brown, thin- and smooth-walled, ellipsoid to obovoid, 0–3-septate, (6.5–)9–11(–12) × (3–)4(–4.5) µm, with a protruding hilum, 1 µm diam, and minute marginal frill.

Colour illustrations. *Evernia prunastri* at Dwingeloo graveyard, Drenthe Province. Left column: *Cylindromonium everniae*. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Right column: *Rhodoveronaea everniae*. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Culture characteristics — Colonies erumpent, spreading, with moderate aerial mycelium and smooth, lobate margin, reaching 4 mm diam after 7 d at 25 °C. On MEA surface saffron, reverse pale luteous; on PDA surface and reverse isabelline; on OA surface ochreous.

Typus. NETHERLANDS, Drenthe Province, Dwingeloo graveyard, 52.831376, 6.365404, on *Evernia prunastri* (*Parmeliaceae*), 6 Jan. 2021, J. Boers, HPC 3557 (holotype CBS H-24848, culture ex-type CPC 41005 = CBS 148309, ITS, LSU and *rpb2* sequences GenBank OK664737.1, OK663776.1 and OK651172.1, MycoBank MB 841818).

(notes *Rhodoveronaea everniae* on Supplementary material page FP1317 & 1318)

Supplementary material

FP1318 Phylogenetic tree.

Kiflimonium junci



Fungal Planet 1319 – 24 December 2021

Kiflimonium junci Crous & Osieck, *sp. nov.*

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** erect, hyaline, smooth-walled, subcylindrical, arising from superficial mycelium, base slightly swollen, frequently branched, giving rise to 2–3 conidiogenous cells, 1–3-septate, 60–110 × 2.5–4 µm. **Conidiogenous cells** hyaline, smooth-walled, flexuous, subcylindrical with apical taper, monophialidic, with non-flared apical collarette, 1–1.5 µm diam, 50–100 × 2–3 µm. **Conidia** solitary, aggregating in mucoid mass, hyaline, smooth-walled, guttulate, reniform, apex obtuse to subobtuse, tapering to truncate hilum, 0.5 µm diam, (3.5–)5–6(–7) × (2–)2.5(–3) µm.

Culture characteristics — Colonies flat, spreading, surface folded, with sparse to moderate aerial mycelium and smooth, lobate margin, reaching 20 mm diam after 2 wk at 25 °C. On MEA surface ochreous, reverse umber; on PDA surface and reverse pale luteous; on OA surface pale luteous.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'44" E05°09'40", on dead culm of *Juncus inflexus* (*Juncaceae*), 4 Feb. 2021, *E.R. Osieck*, HPC 3590 = WI-21/#4205 (holotype CBS H-24825, cultures ex-type CPC 41222 = CBS 148279, CPC 41223 = CBS 148280, ITS, LSU and *actA* sequences GenBank OK664738.1, OK663777.1 and OK651130.1, MycoBank MB 841819).

Notes — *Kiflimonium* (based on *K. curvulum*) was established by Summerbell et al. (2018) to accommodate a species with 0–1-septate, curved conidia (4–6.7 × 1.4–2.1 µm) (Gams 1971). Although the genus was monotypic, Summerbell et al. (2018) referred to several undescribed species deposited in the CBS collection, which will be named in a future study. *Kiflimonium junci* is closely related to *K. curvulum*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence of CPC 41222 had highest similarity to *Kiflimonium curvulum* (strain CBS 825.70, GenBank MH859964.1; Identities = 492/549 (90 %), 21 gaps (3 %)), *Achroistachys aurantisporea* (strain CBS 187.73, GenBank KU845803.1; Identities = 479/568 (84 %), 39 gaps (6 %)), and *Neonectria ditissima* (strain CBS 226.31, GenBank JF735309.1; Identities = 481/571 (84 %), 39 gaps (6 %)). The ITS sequences of CPC 41222 and CPC 41223 are identical (544/544 nucleotides, no gaps). Closest hits using the LSU sequence of CPC 41222 are *Kiflimonium curvulum* (strain CBS 384.70C, GenBank MH871504.1; Identities = 801/814 (98 %), no gaps), *Paramyrothecium roridum* (as *Myrothecium roridum*; strain ATCC 16297, GenBank AY489708.1; Identities = 770/815 (94 %), six gaps (0 %)), and *Lasionectria martiniensis* (strain CBS 129746, GenBank KP899121.1; Identities = 770/816 (94 %), five gaps (0 %)). Closest hits using the *actA* sequence of CPC 41222 had highest similarity to *Sarocladium dejongiae* (strain CBS 144929, GenBank MK069412.1; Identities = 469/514 (91 %), seven gaps (1 %)), *Sarocladium sasijaorum* (strain NL19100007, GenBank MW890032.1; Identities = 556/643 (86 %), 21 gaps (3 %)), and *Sarocladium kiliense* (strain CBS 400.52, GenBank KM231258.1; Identities = 555/651 (85 %), 31 gaps (4 %)).

Colour illustrations. *Juncus inflexus* at Nieuw Wulven, near Houten. Conidiophores on SNA; conidiophores and conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1284.



Fungal Planet 1320 – 24 December 2021

Varicosporellopsis americana* Crous & Jurjević, sp. nov.Etymology.* Name refers to the country where it was collected, USA.*Classification* — *Nectriaceae*, *Hypocreales*, *Sordariomycetes*.

Mycelium consisting of hyaline, branched, septate, smooth, 3–5 µm diam hyphae, lacking chlamydospores, and frequently forming hyphal coils. *Conidiophores* solitary, erect, branched at base, 0–2-septate, or reduced to conidiogenous cells, 40–70 × 3–4 µm, giving rise to lateral branches or conidiogenous cells. *Conidiogenous cells* subcylindrical with slight apical taper, hyaline, smooth, 30–50 × 3–4 µm, apex phialidic with flared collarette, 2–3 mm diam, giving rise to clusters of slimy conidia. *Conidia* solitary, hyaline, smooth, guttulate, ellipsoid, aseptate, straight to curved, apex subobtusate, base tapered to a truncate hilum, 1–1.5 mm diam, (10–)12–16(–24) × (3–)4(–4.5) µm.

Culture characteristics — Colonies flat, spreading, with sparse to moderate aerial mycelium and smooth, lobate margin, reaching 30 mm diam after 2 wk at 25 °C. On MEA, PDA and OA surface and reverse saffron.

Typus. USA, California, San Francisco, from sludge in water reservoir, Apr. 2020, Ž. Jurjević 5496 (holotype CBS H-24803, culture ex-type CPC 40767 = CBS 148257, ITS, LSU, *actA* and *tub2* sequences GenBank OK664739.1, OK663778.1, OK651131.1, and OK651211.1, MycoBank MB 841820); *ibid.*, CBS H-24804, culture CPC 40768 = CBS 148258, ITS, LSU, *actA* and *tub2* sequences GenBank OK664740.1, OK663779.1, OK651132.1, and OK651212.1.

Notes — The genus *Varicosporellopsis* is monotypic, based on *V. aquatilis*, which was described from submerged wood collected in freshwater in southwestern France (conidia 6–11 × 2.8–3.2 µm; Lechat & Fournier 2016), and later also reported from soil in the Netherlands (conidia (6–)11–13(–15) × (3–)4(–4.5) µm; Crous et al. 2018b). *Varicosporellopsis americana* is distinct in that it has larger conidia. *Varicosporellopsis americana* forms a lineage sister to *V. aquatilis*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence of CPC 40767 had highest similarity to *Varicosporellopsis aquatilis* (strain CBS 140158, GenBank NR_171219.1; Identities = 506/530 (95 %), five gaps (0 %)), *Paracremonium variiforme* (strain LC5832, GenBank KU746693.1; Identities = 473/498 (95 %), eight gaps (1 %)), and *Paracremonium binnewijzendii* (strain CBS 143277, GenBank NR_157491.1; Identities = 533/564 (95 %), 13 gaps (2 %)). The ITS sequences of CPC 40767 and 40768 are identical (551/551 nucleotides, no gaps). Closest hits using the **LSU** sequence of CPC 40767 are *Varicosporellopsis aquatilis* (strain CBS 140158, GenBank NG_058207.1; Identities = 802/815 (98 %), no gaps), *Paracremonium pembeum* (strain P169, GenBank MT252038.1; Identities = 797/814 (98 %), no gaps), and *Paracremonium variiforme* (strain LC5832, GenBank KU746739.1; Identities = 796/813 (98 %), one gap (0 %)). The LSU sequences of CPC 40767 and 40768 are identical (807/807 nucleotides, no gaps). Closest hits using the **actA** sequence of CPC 40767 had highest similarity to *Paracremonium contagium* (strain CBS 110348, GenBank KM231240.1; Identities = 502/567 (89 %), 18 gaps (3 %)), *Paracremonium inflatum* (strain CBS 485.77, GenBank KM231238.1; Identities = 487/548 (89 %), 11 gaps (2 %)), and *Xenoacremonium recifei* (strain CBS 137.35, GenBank KM231241.1; Identities = 425/470 (90 %), five gaps (1 %)). The *actA* sequences of CPC 40767 and 40768 are identical (565/565 nucleotides, no gaps). Closest hits using the **tub2** sequence of CPC 40767 had highest similarity to *Varicosporellopsis aquatilis* (strain CBS 143509, GenBank MH108052.1; Identities = 509/542 (94 %), 12 gaps (2 %)), *Paracremonium inflatum* (strain CBS 485.77, GenBank KM232101.1; Identities = 480/536 (90 %), 21 gaps (3 %)), and *Paracremonium bendijkiorum* (strain NL19_24005, GenBank MW890139.1; Identities = 484/545 (89 %), 22 gaps (4 %)). The *tub2* sequences of CPC 40767 and 40768 are identical (535/535 nucleotides, no gaps).

Colour illustrations. Water reservoir with sludge. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1284.



Fungal Planet 1321 – 24 December 2021

***Cylindrium desperesense* Crous & Jurjević, sp. nov.**

Etymology. Name refers to the collection site, Des Peres, Missouri, USA.

Classification — *Cylindriaceae*, *Hypocreales*, *Sordariomycetes*.

Conidiomata sporodochial, crystalline to saffron, 50–300 µm diam, lacking chlamydospores. *Conidiophores* subcylindrical, smooth-walled, hyaline, branched, septate, up to 40 µm tall, 3–4 µm wide. *Conidiogenous cells* hyaline, smooth-walled, terminal and lateral, 5–10 × 1.5–2.5 µm; proliferating sympodially, scars flattened, not thickened nor darkened. *Conidia* hyaline, smooth-walled, aseptate, thick-walled, with two large guttules, granular, fusoid-ellipsoid, straight to slightly dorsiventrally curved, tapering towards both ends, apex bluntly rounded, somewhat truncate, 1–1.5 µm diam; hilum truncate, 1.5–2 µm diam, (12–)14–16(–22) × 4(–5) µm.


Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 40 mm diam after 2 wk at 25 °C. On MEA surface and reverse orange; on PDA and OA surface and reverse saffron.

Typus. USA, Missouri, Des Peres, from air in a store, May 2020, Ž. Jurjević 5505 (holotype CBS H-24805, culture ex-type CPC 40769 = CBS 148259, ITS, LSU, *actA*, *cmdA*, *rpb2*, *tef1* (part 1) and *tub2* sequences GenBank OK664741.1, OK663780.1, OK651133.1, OK651144.1, OK651173.1, OK651193.1 and OK651213.1, MycoBank MB 841821).

Notes — *Cylindrium* accommodates species with sporodochia, hyaline to pale brown, subcylindrical conidiophores, conidiogenous cells with sympodial proliferation, and aseptate, hyaline conidia (usually in chains) (Lombard et al. 2015, Crous et al. 2018b). *Cylindrium desperesense* represents a new, distinct species based on its conidial morphology and phylogeny.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Cylindrium corymbiae* (strain CPC 35637, GenBank MT223792.1; Identities = 518/560 (93 %), 23 gaps (4 %)), *Cylindrium grande* (strain CBS 145655, GenBank NR_165557.1; Identities = 520/563 (92 %), 24 gaps (4 %)), and *Cylindrium elongatum* (strain CBS 685.83A, GenBank KM231852.1; Identities = 514/561 (92 %), 23 gaps (4 %)). Closest hits using the LSU sequence are *Pseudoidriella syzygii* (strain CBS 131307, GenBank JQ044441.1; Identities = 813/817 (99 %), no gaps), *Cylindrium grande* (strain CBS 145655, GenBank NG_068656.1; Identities = 799/805 (99 %), no gaps), and *Tristataperidium microsporum* (strain MFLUCC 15-0413, GenBank KT696539.1; Identities = 730/736 (99 %), no gaps). Closest hits using the *actA* sequence had highest similarity to *Cylindrium grande* (strain CBS 145578, GenBank MK876456.1; Identities = 599/693 (86 %), 30 gaps (4 %)), *Cylindrium corymbiae* (strain

CBS 146087, GenBank MT223750.1; Identities = 481/526 (91 %), seven gaps (1 %)), and *Cylindrium aeruginosum* (strain CBS 693.83, GenBank KM231265.1; Identities = 497/550 (90 %), 13 gaps (2 %)). Closest hits using the *cmdA* sequence had highest similarity to *Cylindrium grande* (strain CBS 145578, GenBank MK876468.1; Identities = 483/620 (78 %), 38 gaps (6 %)), *Cylindrium corymbiae* (strain CBS 146087, GenBank MT223763.1; Identities = 484/622 (78 %), 41 gaps (6 %)), and *Cylindrium elongatum* (strain CBS 115974, GenBank KM231449.1; Identities = 350/444 (79 %), 30 gaps (6 %)). Closest hits using the *rpb2* sequence had highest similarity to *Cylindrium grande* (strain CPC 35403, GenBank MK876481.1; Identities = 668/764 (87 %), no gaps), and *Cylindrium corymbiae* (strain CBS 146087, GenBank MT223679.1; Identities = 661/764 (87 %), no gaps). Closest hits using the *tef1* (first part) sequence had highest similarity to *Cylindrium corymbiae* (strain CBS 146087, GenBank MT223712.1; Identities = 269/304 (88 %), 16 gaps (5 %)), *Cylindrium elongatum* (strain CBS 685.83A, GenBank KM231988.1; Identities = 258/288 (90 %), 15 gaps (5 %)), and *Cylindrium grande* (strain CPC 35403, GenBank MK876495.1; Identities = 255/286 (89 %), ten gaps (3 %)). Closest hits using the *tub2* sequence had highest similarity to *Cylindrium elongatum* (strain CBS 685.83A, GenBank KM232122.1; Identities = 534/683 (78 %), 48 gaps (7 %)), and *Cylindrium corymbiae* (strain CBS 146087, GenBank MT223732.1; Identities = 519/674 (77 %), 41 gaps (6 %)).



The first of 36 equally most parsimonious trees obtained from a phylogenetic analysis of the *Cylindrium* ITS nucleotide alignment. The tree was rooted to *Phlogicylindrium tereticornis* (culture CBS 143168; GenBank NR_156660.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured block and bold text. GenBank accession (superscript) and/or culture collection/voucher numbers (in bold face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 18 strains including the outgroup; 533 characters including alignment gaps analysed: 314 constant, 130 variable and parsimony-uninformative and 89 parsimony-informative. Tree statistics: Tree Length = 366, Consistency Index = 0.790, Retention Index = 0.709, Rescaled Consistency Index = 0.560. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Colour illustrations. Store in Des Peres, Missouri, USA. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Valsonectria portsmouthensis



Fungal Planet 1322 – 24 December 2021

***Valsonectria portsmouthensis* Crous & Jurjević, sp. nov.**

Etymology. Name refers to the location where the fungus was collected, Portsmouth, New Hampshire, USA.

Classification — *Incertae sedis*, *Hypocreales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. *Conidiophores* reduced to conidiogenous cells, solitary, arising from superficial hyphae, flexuous, subulate with apical taper, hyaline, smooth-walled, 15–45 × 2.5–3 µm, monophialidic, apex without flared collar-ette, 1 µm diam. *Conidia* solitary, aggregating in mucoid mass, hyaline, smooth-walled, guttulate, 0–1-septate, subcylindrical, apex subobtuse, tapering at base to truncate hilum, 1 µm diam, (10–)14–16(–22) × 3.5–4 µm.

Culture characteristics — Colonies flat, spreading, with moderate aerial mycelium and feathery, lobate margin, reaching 30 mm diam after 2 wk at 25 °C. On MEA surface dirty white, reverse pale luteous; on PDA and OA surface and reverse dirty white.

Typus. USA, New Hampshire, Portsmouth, from air in men's locker room, Aug. 2020, Ž. Jurjević 5534 (holotype CBS H-24806, culture ex-type CPC 40779 = CBS 148260, ITS, LSU, *actA*, *rpb2* and *tub2* sequences GenBank OK664742.1, OK663781.1, OK651134.1, OK651174.1 and OK651214.1, MycoBank MB 841822).

Notes — *Valsonectria portsmouthensis* clusters in a clade with *V. pulchella*, the type species of *Valsonectria*, which is known to have acremonium-like and stilbella-like asexual morphs (Rossman 2000). It thus seems prudent to name the present acremonium-like isolate in *Valsonectria*, where it is distinct from species presently known from culture (Summerbell et al. 2011).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Acremonium roseolum* (strain CBS 289.62, GenBank MH858153.1; Identities = 530/575 (92 %), 15 gaps (2 %)), *Acremonium hyalinulum* (strain FMR 10351, GenBank KP131521.1; Identities = 526/574 (92 %), 17 gaps (2 %)), and *Nectria bactridioides* (strain IMI 063286, GenBank AY254161.1; Identities = 518/570 (91 %), 21 gaps (3 %)). Closest hits using the **LSU** sequence are *Acremonium roseolum* (strain CBS 289.62, GenBank MH869748.1; Identities = 799/810 (99 %), no gaps), *Scopinella solani* (strain CBS 770.84, GenBank AY015632.1; Identities = 797/808 (99 %), no gaps), and *Valsonectria pulchella* (strain MUT<ITA> 4890, GenBank KP671718.1; Identities = 800/812 (99 %), two gaps (0 %)). Closest hits using the **actA** sequence had highest similarity to *Aquilomyces patris* (strain CBS 135760, GenBank KP184120.1; Identities = 556/627 (89 %), 17 gaps (2 %)), *Darksidea alpha* (strain CBS 135630, GenBank KP184090.1; Identities = 531/616 (86 %), 18 gaps (2 %)), and *Roussoella euonymi* (strain CBS 143426, GenBank MH107986.1; Identities = 448/496 (90 %), three gaps (0 %)). No significant hits were obtained when the **rpb2** and **tub2** sequences were used in blastn and megablast searches.

Colour illustrations. Men's locker room, Portsmouth, New Hampshire, USA. Conidiogenous cells on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1284.

Toxicocladosporium losalamitosense



Fungal Planet 1323 – 24 December 2021

***Toxicocladosporium losalamitosense* Crous & Jurjević, sp. nov.**

Etymology. Name refers to the place where the present fungus was collected, Los Alamitos, California.

Classification — *Cladosporiaceae*, *Cladosporiales*, *Dothideomycetes*.

Mycelium consisting of pale brown, smooth to roughened, septate, branched, 2–3 µm diam hyphae. *Conidiophores* reduced to individual conidiogenous loci on hyphae. *Conidiogenous cells* integrated, 1–2 × 1.5–2 µm; scars thickened, darkened and somewhat refractive. *Ramoconidia* 0–1-septate, pale brown, smooth-walled, subcylindrical, 8–17 × 2.5–3 µm; conidia in branched chains; *intercalary conidia* pale brown, smooth, fusoid-ellipsoid, aseptate, tapering toward both ends, (6–)9–11 × 2–2.5 µm; *small terminal conidia* aseptate, pale brown, smooth, fusoid-ellipsoid, 6–9 × 2(–2.5) µm; scars thickened, darkened, and somewhat refractive, 1–1.5 µm diam. Submerged hyphae forming chains of *chlamydospores*, becoming dark brown, smooth- and thick-walled, muriformly septate, 6–15 µm diam.

Culture characteristics — Colonies erumpent, spreading, surface folded, with sparse aerial mycelium and feathery, lobate margin, reaching 5 mm diam after 7 d at 25 °C. On MEA surface isabelline, reverse sepia; on PDA surface and reverse sepia; on OA surface sepia.

Typus. USA, California, Los Alamitos, from air in a classroom, Oct. 2020, Ž. Jurjević 5558 (holotype CBS H-24807, culture ex-type CPC 40792 = CBS 148261, ITS, LSU, *actA*, *rpb2* and *tub2* sequences GenBank OK664743.1, OK663782.1, OK651135.1, OK651175.1 and OK651215.1, MycoBank MB 841823).

Colour illustrations. Classroom at Los Alamitos, California. Chlamydospores on SNA; conidiogenous cell giving rise to conidia; conidial chains. Scale bars = 10 µm.


Notes — *Toxicocladosporium losalamitosense* is related to *T. ficiniae*. It is morphologically and phylogenetically distinct from all known species (Bezerra et al. 2017).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Toxicocladosporium hominis* (as *Toxicocladosporium* sp.; strain UTHSC DI-13-172, GenBank LN834444.1; Identities = 467/481 (97 %), one gap (0 %)), *Toxicocladosporium banksiae* (strain CBS 128215, GenBank NR_152322.1; Identities = 532/550 (97 %), four gaps (0 %)), and *Toxicocladosporium irritans* (voucher KoLRI_EL006197, GenBank MN844802.1; Identities = 466/482 (97 %), three gaps (0 %)). Closest hits using the **LSU** sequence are *Toxicocladosporium ficiniae* (strain CBS 136406, GenBank NG_058054.1; Identities = 785/797 (98 %), two gaps (0 %)), *Toxicocladosporium irritans* (strain CBS 185.58, GenBank MH869283.1; Identities = 781/797 (98 %), two gaps (0 %)), and *Toxicocladosporium strelitziae* (strain CBS 132535, GenBank NG_042687.1; Identities = 781/797 (98 %), two gaps (0 %)). Closest hits using the **actA** sequence had highest similarity to *Cladosporium inversicolor* (strain CPC 18238, GenBank KT600588.1; Identities = 390/412 (95 %), no gaps), *Cladosporium sinuosum* (strain CPC 18365, GenBank KT600643.1; Identities = 403/433 (93 %), no gaps), and *Cladosporium velox* (strain CPC 18450, GenBank KT600654.1; Identities = 387/410 (94 %), no gaps). Closest hits using the **rpb2** sequence had highest similarity to *Toxicocladosporium strelitziae* (strain CPC 19762, GenBank LT799788.1; Identities = 712/858 (83 %), 18 gaps (2 %)), *Toxicocladosporium ficiniae* (strain CPC 21283, GenBank LT799780.1; Identities = 683/834 (82 %), 14 gaps (1 %)), and *Toxicocladosporium hominis* (strain FMR 13297, GenBank LT799781.1; Identities = 677/831 (81 %), 18 gaps (2 %)). Closest hits using the **tub2** sequence had highest similarity to *Toxicocladosporium velox* (strain CBS 124159, GenBank KY706609.1; Identities = 282/345 (82 %), 12 gaps (3 %)), *Toxicocladosporium chlamydosporum* (strain CBS 124157, GenBank KY706598.1; Identities = 279/341 (82 %), 12 gaps (3 %)), and *Toxicocladosporium irritans* (strain CBS 185.58, GenBank KY706601.1; Identities = 281/346 (81 %), 16 gaps (4 %)).

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1303.

Phialoseptomonium junci
& *Alfaria junci*



Fungal Planet 1324 & 1325 – 24 December 2021

***Phialoseptomonium junci* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Nectriaceae*, *Hypocreales*, *Sordariomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1.5–2 µm diam hyphae. **Conidiophores** erect, flexuous, unbranched, arising directly from superficial mycelium, cylindrical, 2–3-septate, 200–300 × 2.5–3.5 µm, base pale brown, smooth-walled, becoming hyaline toward apex. **Conidiogenous cells** apical, integrated, subcylindrical, phialidic, with minute non-flared collarette, 60–120 × 1.5–2 µm. **Conidia** solitary, aggregating in a mucoid mass, hyaline, smooth-walled, granular, guttulate, fusoid, straight, 3-septate, apex obtuse, base truncate, 1–1.5 µm diam, (20–)25–28(–32) × (3.5–)4–4.5(–5) µm.

Culture characteristics — Colonies flat, spreading, with moderate aerial mycelium and feathery, lobate margin, reaching 30 mm diam after 2 wk at 25 °C. On MEA surface saffron, reverse ochrous; on PDA and OA surface and reverse saffron.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'45" E05°10'34", on dead culm of *Juncus effusus* (*Juncaceae*), 8 Jan. 2021, E.R. Osieck, HPC 3564 = WI-10/#4195 (holotype CBS H-24846, culture ex-type CPC 40819 = CBS 148307, ITS and LSU sequences GenBank OK664744.1 and OK663783.1, MycoBank MB 841824).

***Alfaria junci* Crous & Osieck, sp. nov.**

Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*.

Conidiomata sporodochial, stromatic, superficial, cupulate, scattered to gregarious, oval in outline, 100–300 µm diam, with surrounding setae. **Stroma** well-developed, of hyaline *textura angularis*. **Marginal hyphae** arising from stroma, straight to curved, irregularly branched. **Setae** arising from stroma, thick and smooth-walled, unbranched, pale brown, septate, terminating to subobtuse apex, 80–120 × 4–5 µm. **Conidiophores** subcylindrical, hyaline, smooth-walled, penicillately branched. **Conidiogenous cells** phialidic, hyaline, smooth-walled, subcylindrical, tapering at apex with periclinal thickening, 5–20 × 2–2.5 µm. **Conidia** aseptate, subcylindrical, pale olivaceous, aggregating in mucoid mass, guttulate, apex subobtuse, base truncate, (10–)11–12(–13) × 2(–2.5) µm.

Culture characteristics — Colonies flat, spreading, surface folded, with moderate aerial mycelium and smooth, regular margin, covering dish after 2 wk at 25 °C. On MEA, PDA and OA surface dirty white, reverse luteous.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°02'45" E05°10'34", on dead culm of *Juncus effusus* (*Juncaceae*), 8 Jan. 2021, E.R. Osieck, HPC 3564 = WI-10/#4195 (holotype CBS H-24811, culture ex-type CPC 40821 = CBS 148265, ITS and LSU sequences GenBank OK664745.1 and OK663784.1, MycoBank MB 841825).

Colour illustrations. *Juncus effusus* at Nieuw Wulven, near Houten. Left column: *Phialoseptomonium junci*. Flexuous conidiophores on SNA giving rise to conidia; conidia. Right column: *Alfaria junci*. Conidiomata on PNA; conidiophores and conidiogenous cells giving rise to conidia; seta; conidiogenous cells; conidia. Scale bars = 10 µm.

Notes — *Phialoseptomonium* is an acremonium-like genus characterised by long, flexuous conidiophores, and hyaline, septate, fusoid conidia. *Phialoseptomonium junci* is closely related to *P. eucalypti* (conidia 1-septate, (16–)19–21(–23) × 3(–3.5) µm; Crous et al. 2019a), but distinct in that it has longer, 3-septate conidia.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Phialoseptomonium eucalypti* (strain CBS 145542, GenBank NR_165572.1; Identities = 558/592 (94 %), ten gaps (1 %)), *Cylindromonium lichenicola* (strain CBS 188.70, GenBank MH859549.1; Identities = 545/596 (91 %), 14 gaps (2 %)), and *Cylindromonium alloxyli* (strain CBS 146806, GenBank NR_171760.1; Identities = 532/587 (91 %), 16 gaps (2 %)). Closest hits using the **LSU** sequence are *Cylindromonium lichenicola* (strain CBS 303.70, GenBank MH871429.1; Identities = 800/819 (98 %), four gaps (0 %)), *Phialoseptomonium eucalypti* (strain CBS 145542, GenBank NG_067890.1; Identities = 781/800 (98 %), three gaps (0 %)), and *Cylindromonium lichenicola* (strain CBS 415.70A, GenBank MH871536.1; Identities = 804/824 (98 %), three gaps (0 %)).

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1284.

Notes — The sexual genus *Alfaria* (Crous et al. 2014a) clusters with myrothecium-like asexual morphs (Lombard et al. 2016). The present collection is phylogenetically related to isolates that have in the past also been identified as species of *Amerosporium* (*A. atrum* CBS 151.69, *A. platense* CBS 658.73), suggesting that these isolates also represent species of *Alfaria*. Johnston & Gamundi (2000) linked *Amerosporium patellarioidesi*, the type species of *Amerosporium* to the sexual morph *Zoellneria* (*Chaetomellaceae*, *Helotiales*). *Amerosporium*, however, remains a rather confused genus in need to revision. *Alfaria junci* is phylogenetically distinct from other species presently accepted in *Alfaria*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Amerosporium atrum* (strain CBS 166.65, GenBank MH858530.1; Identities = 559/563 (99 %), two gaps (0 %)), *Myrothecium gramineum* (strain XF36, GenBank KJ780796.1; Identities = 551/557 (99 %), two gaps (0 %)), and *Xepicula leucotricha* (strain CK775, GenBank MH474462.1; Identities = 448/456 (98 %), four gaps (0 %)). Closest hits using the **LSU** sequence are *Alfaria dandenongensis* (strain CBS 143399, GenBank NG_069537.1; Identities = 785/785 (100 %), no gaps), *Amerosporium atrum* (strain CBS 151.69, GenBank MH877704.1; Identities = 785/785 (100 %), no gaps), and *Amerosporium platense* (strain CBS 658.73, GenBank MH872519.1; Identities = 785/785 (100 %), no gaps).

Supplementary material

FP1325 Phylogenetic tree.



Fungal Planet 1326 – 24 December 2021

***Xenoacrodoniaceae* Crous, fam. nov.**

Classification — *Xenoacrodoniaceae*, *Hypocreales*, *Sordariomycetes*.

Etymology. Name is based on the genus *Xenoacrodonium*.

Mycelium consisting of hyaline, smooth-walled, branched, septate hyphae. *Conidiophores* reduced to conidiogenous cells arising directly from hyphae, hyaline, smooth- and thin-walled, subulate, straight to flexuous, proliferating sympodially, forming a rachis in upper part, with multiple subdenticulate loci, slightly

subulate, straight to flexuous, proliferating sympodially, forming a rachis in upper part, with multiple subdenticulate loci, slightly thickened and refractive, not darkened. *Conidia* solitary, hyaline, aseptate, smooth- and thin-walled, guttulate, ellipsoid, with obtuse apex; hilum slightly thickened, not darkened.

Type genus. *Xenoacrodonium* Crous
MycoBank MB 841843.

***Xenoacrodonium* Crous, gen. nov.**

Etymology. Name refers to its morphological similarity to *Acrodonium*.

Mycelium consisting of hyaline, smooth-walled, branched, septate hyphae. *Conidiophores* reduced to conidiogenous cells arising directly from hyphae, hyaline, smooth- and thin-walled, subulate, straight to flexuous, proliferating sympodially, forming a rachis in upper part, with multiple subdenticulate loci, slightly

thickened and refractive, not darkened. *Conidia* solitary, hyaline, aseptate, smooth- and thin-walled, guttulate, ellipsoid, with obtuse apex; hilum slightly thickened, not darkened.

Type genus. *Xenoacrodonium juglandis* Crous
MycoBank MB 841826.

***Xenoacrodonium juglandis* Crous, sp. nov.**

Etymology. Name refers to the host genus *Juglans* from which it was isolated.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 1–1.5 µm diam hyphae. *Conidiophores* reduced to conidiogenous cells arising directly from hyphae, hyaline, smooth- and thin-walled, subulate, straight to flexuous, proliferating sympodially, forming a rachis in upper part, 20–40 × 2–3 µm, with multiple subdenticulate loci, slightly thickened and refractive, not darkened. *Conidia* solitary, hyaline, aseptate, smooth- and thin-walled, guttulate, ellipsoid, with obtuse apex, (4–)5(–6) × 2(–2.5) µm; hilum slightly thickened, not darkened, 0.5 µm diam.

Culture characteristics — Colonies erumpent, spreading, surface folded, with moderate aerial mycelium and feathery, lobate margin, reaching 12 mm diam after 2 wk at 25 °C. On MEA surface pale luteous, reverse ochreous; on PDA and OA surface and reverse pale luteous.

Typus. NETHERLANDS, Gelderland Province, Beuningen, river Waal, on *Juglans regia* (*Juglandaceae*), 17 Oct. 2020, A.L. van Iperen, HPC 3489 (holotype CBS H-24840, culture ex-type CPC 40016 = CBS 148301, ITS, LSU, *actA*, *cmdA*, *his3*, *rpb2* and *tef1* (first part) sequences GenBank OK664746.1, OK663785.1, OK651136.1, OK651145.1, OK651150.1, OK651176.1 and OK651194.1, MycoBank MB 841827).

Colour illustrations. *Juglans regia* at the river Waal, Beuningen. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Notes — *Acrodonium* was shown to be a genus in *Teratosphaeriaceae*, with several *acrodonium*-like species belonging to other orders (Videira et al. 2016). *Xenoacrodonium* is presently a monotypic genus in a new family, *Xenoacrodoniaceae*, clustering with isolate CBS 580.67 (isolated as culture contaminant, Houston, Texas, and identified as *Acrodonium salmoneum*), which appears to represent a second species in the genus.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to '*Acrodonium salmoneum*' (strain ANER3, GenBank MK370695.1; Identities = 466/511 (91 %), 17 gaps (3 %)), *Stachybotrys reniformis* (strain CBS 136198, GenBank KU846740.1; Identities = 494/575 (86 %), 29 gaps (5 %)), and *Stachybotrys nephrospora* (strain ATCC 18839, GenBank AF081476.2; Identities = 494/575 (86 %), 29 gaps (5 %)). Closest hits using the LSU sequence are '*Acrodonium salmoneum*' (strain CBS 580.67, GenBank MH870773.1; Identities = 793/800 (99 %), no gaps), *Sarocladium brachiariae* (as *Sarocladium* sp. XBL-2015; strain HND5, GenBank KP715271.1; Identities = 762/800 (95 %), no gaps), and *Sarocladium implicatum* (strain CBS 125892, GenBank MH875549.1; Identities = 763/802 (95 %), two gaps (0 %)). Only distant hits with members of *Hypocreales* were obtained when the *actA*, *cmdA*, *his3*, *rpb2* and the *tef1* (first part) sequences were used in blastn and megablast searches.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1284.



Fungal Planet 1327 – 24 December 2021

***Sporidesmiella pini* Crous, sp. nov.**

Etymology. Name refers to the host genus *Pinus* from which it was isolated.

Classification — *Junewangiaceae*, *Incertae sedis*, *Sordario-mycetes*.

Mycelium consisting of pale brown, smooth, branched, septate, 2–2.5 µm diam hyphae. **Conidiophores** solitary, arising from superficial mycelium, subcylindrical, erect, unbranched, thick-walled, brown, smooth, 2–3-septate, 30–70 × 3.5–5 µm; basal cell T-shaped or globose, 6–8 µm diam. **Conidiogenous cells** integrated, terminal, brown- and smooth-walled, subcylindrical, 11–25 × 3–4 µm, proliferating percurrently. **Conidia** solitary, arranged in a rosette due to delayed succession, obovoid, brown-, smooth- and thick-walled, (3–)4-distoseptate, with central pore in septum, (14–)18–23(–27) × (8–)9–10(–12) µm; hilum truncate, 2–2.5 µm diam, with marginal frill.

Culture characteristics — Colonies erumpent, spreading, with moderate aerial mycelium and smooth, lobate margin, reaching 17 mm diam after 2 wk at 25 °C. On MEA surface dirty white, reverse ochreous; on PDA surface and reverse red; on OA surface saffron.

Typus. NETHERLANDS, Utrecht Province, Soest, de Zoom, on *Pinus sylvestris* (*Pinaceae*), 1 Nov. 2020, A.L. van Iperen, HPC 3492 (holotype CBS H-24841, culture ex-type CPC 40067 = CBS 148302, ITS, LSU and *rpb2* sequences GenBank OK664747.1, OK663786.1 and OK651177.1, MycoBank MB 841828).

Notes — *Sporidesmiella* was established by Kirk (1982), based on *S. claviformis*. *Sporidesmiella pini* is closely related to *S. obovoidia* (conidia (3–)4-distoseptate, 20–25 × 9–11.5 µm; Dong et al. 2021) and *S. hyalosperma* (conidia brown, smooth, thick-walled, 4-distoseptate, (17–)22–25(–29) × 9–12(–13) µm, Kirk 1982; (19–)20–22(–24) × (9–)10(–11) µm, Crous et al. 2020c), but is distinct based on its conidial morphology and phylogeny.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Sporidesmiella obovoidia* (strain MFLUCC 17-2372, GenBank NR_172446.1; Identities = 548/571 (96 %), 12 gaps (2 %)), *Sporidesmiella hyalosperma* (strain CPC 37552, GenBank MT223845.1; Identities = 546/569 (96 %), nine gaps (1 %)), and *Sporidesmiella novae-zelandiae* (strain S-1256, GenBank MK828693.1; Identities = 475/533 (89 %), 25 gaps (4 %)). Closest hits using the **LSU** sequence are *Sporidesmiella obovoidia* (strain MFLUCC 17-2372, GenBank NG_075412.1; Identities = 786/790 (99 %), one gap (0 %)), *Sporidesmiella hyalosperma* (strain S-1518, GenBank MK849842.1; Identities = 792/799 (99 %), one gap (0 %)), and *Sporidesmiella novae-zelandiae* (strain S-1256, GenBank MK849845.1; Identities = 784/799 (98 %), two gaps (0 %)). Closest hits using the **rpb2** sequence had highest similarity to *Sporidesmiella hyalosperma* (strain MFLUCC 18-1013, GenBank MW504070.1; Identities = 774/832 (93 %), no gaps), *Sporidesmiella novae-zelandiae* (voucher MFLU 18-2332, GenBank MN124525.1; Identities = 733/832 (88 %), no gaps), and *Sporidesmiella aquatica* (voucher MFLU 18-2331, GenBank MN124524.1; Identities = 638/735 (87 %), two gaps (0 %)).

Colour illustrations. *Pinus sylvestris* at Soest. Conidiophores on SNA; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1304.



Fungal Planet 1328 – 24 December 2021

***Paraxerochrysium* Crous & Decock, gen. nov.**

Etymology. Name refers to its close relationship to the genus *Xerochrysium*.

Classification — *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*.

Mycelium consisting of hyaline, smooth-walled, branched, septate hyphae. Reproductive structures consisting of solitary

or short chains of chlamydoconidia, borne by retrogressive sympodial formation from hyphal tips, or as intercalary chlamydoconidia. *Chlamydoconidia* spherical, thick- and smooth-walled, guttulate, hyaline.

Type species. *Paraxerochrysium coryli* Crous & Decock
Mycobank MB 841829.

***Paraxerochrysium coryli* Crous & Decock, sp. nov.**

Etymology. Name refers to the host genus *Corylus* from which it was isolated.

Mycelium consisting of hyaline, smooth-walled, branched, septate, 3–4(–7) µm diam hyphae. Reproductive structures consisting of solitary or short chains of chlamydoconidia, borne by retrogressive sympodial formation from hyphal tips, or as intercalary chlamydoconidia. *Chlamydoconidia* spherical, (7–)10–15(–20) µm diam, thick- and smooth-walled, guttulate, hyaline.

Culture characteristics — Colonies erumpent, spreading, surface folded, with sparse aerial mycelium and feathery, lobate margin, reaching 5 mm diam after 2 wk at 25 °C. On MEA surface dirty white, reverse pale luteous; on PDA and OA surface and reverse dirty white.

Typus. BELGIUM, Louvain-la-Neuve, from dry hazelnut, *Corylus avellana* (*Betulaceae*), Feb. 2021, C. Decock, 0234-8 (holotype CBS H-24853, culture ex-type CPC 41272 = CBS 148314 = MUCL 58103, ITS, LSU, *rpb2* and *tub2* sequences GenBank OK664748.1, OK663787.1, OK651178.1 and OK651216.1, MycoBank MB 841830).

Notes — *Paraxerochrysium* is similar to *Xerochrysium*, as both genera are only known from their asexual morphs, and are characterised by having solitary or short chains of chlamydoconidia. *Paraxerochrysium* is distinct in that it lacks aleuroconidia (Pitt et al. 2013). *Paraxerochrysium* clusters with species of *Bettsia*, *Xerochrysium* and *Xeromyces*.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Xerochrysium xerophilum* (strain CBS 153.67, GenBank NR_154476.1; Identities = 514/563 (91 %), 17 gaps (3 %)), *Bettsia fastidia* (strain CBS 454.91, GenBank MH862261.1; Identities = 460/509 (90 %), 30 gaps (5 %)), and *Leiothecium ellipsoideum* (strain CBS 147.75, GenBank MH860904.1; Identities = 490/556 (88 %), 30 gaps (5 %)). Closest hits using the **LSU** sequence are *Xerochrysium dermatitidis* (strain CBS 132.31, GenBank NG_058454.1; Identities = 822/834 (99 %), two gaps (0 %)), *Monilia medoacensis* (strain CBS 222.32, GenBank MH866748.1; Identities = 821/835 (98 %), three gaps (0 %)), and *Xerochrysium xerophilum* (strain FRR 4958, GenBank KC989712.1; Identities = 820/834 (98 %), one gap (0 %)). Closest hits using the **rpb2** sequence had highest similarity to *Xerochrysium dermatitidis* (strain CBS 132.31, GenBank JN121443.1; Identities = 622/699 (89 %), no gaps), *Pseudopenicillium megasporum* (strain CBS 256.55, GenBank JN121473.1; Identities = 538/659 (82 %), no gaps), and *Aspergillus koreanus* (strain EML-GSNP1-2, GenBank KX216527.1; Identities = 416/510 (82 %), two gaps (0 %)). No significant hits were obtained when the **tub2** sequence was used in blastn and megablast searches.

Colour illustrations. Dry hazelnuts. Chlamydoconidia on SNA; chlamydoconidia. Scale bars = 10 µm.

Supplementary material

FP1328 Phylogenetic tree.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Cony A. Decock, Mycothèque de l'Université catholique de Louvain (MUCL, BCCMTM), Earth and Life Institute – ELIM – Mycology, Université catholique de Louvain, Croix du Sud 2 bte L7.05.06, B-1348 Louvain-la-Neuve, Belgium; e-mail: cony.decock@uclouvain.be



Fungal Planet 1329 – 24 December 2021

Paramicrothecium* Crous, Krimhilde Müller, Siepe, Reul & Osieck, gen. nov.Etymology.* Name refers to its similarity to *Microthecium*.Classification — *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*.

Ascomata superficial to immersed in epidermis, globose solitary, hyaline, becoming darker brown (due to maturing ascospores); wall translucent, of *textura epidermoidea* on surface, of 3–6 layers of *textura prismatica* in section, wall covered with short, stubby, thick-walled setae; ascomata initially cleistothecial, but once mature, opening via slightly papillate apex with central ostiole, and exuding a black cirrus of ascospores. *Asci* unitunicate, clavate to subcylindrical with a long stipe, (2–)4(–6)-spored, not staining in Melzer's reagent. *Ascospores* ellipsoid, medium brown, becoming dark brown, smooth- and thick-walled, with

2–3 large guttules, and a large refractive germ pore at each polar end, somewhat truncate. *Mycelium* consisting of hyaline, smooth-walled, branched, septate hyphae. *Conidiophores* reduced to conidiogenous cells that occur singly or more commonly in clusters. *Conidiogenous cells* hyaline, smooth-walled, arising from superficial mycelium, ampulliform, consisting of an obovoid venter, and cylindrical neck. *Conidia* hyaline, smooth- and thin-walled, guttulate, aseptate, shortly clavate, apex obtuse, tapering to truncate hilum, produced in short, false chains that slime down into a globoid mass.

Type species. *Paramicrothecium sambuci* Crous, Krimhilde Müller, Siepe, Reul & Osieck
Mycobank MB 841831.

Paramicrothecium sambuci* Crous, Krimhilde Müller, Siepe, Reul & Osieck, sp. nov.Etymology.* Name refers to the host genus *Sambucus* from which it was isolated.

Ascomata superficial to immersed in epidermis, globose, 200–350 µm diam, scattered, solitary, hyaline, becoming darker brown (due to maturing ascospores); wall translucent, of *textura epidermoidea* on surface, wall 10–15 µm diam, of 3–6 layers of *textura prismatica* in section, wall covered with short, stubby, thick-walled setae, hyaline, ends obtuse, surface warty, 30–100 × 3–4 µm; ascomata initially cleistothecial, but once mature, opening via slightly papillate apex with central ostiole, and exuding a black cirrus of ascospores. *Asci* unitunicate, clavate to subcylindrical with a long stipe, (25–)30–35 × (7–)8–10 µm, (2–)4(–6)-spored, not staining in Melzer's reagent. *Ascospores* ellipsoid, medium brown, becoming dark brown, smooth- and thick-walled, with 2–3 large guttules, and a large refractive germ pore at each polar end, somewhat truncate, (5–)7–9(–10) × (3.5–)5–6(–7.5) µm. *Mycelium* consisting of hyaline, smooth-walled, branched, septate, 3–4 µm diam hyphae. *Conidiophores* reduced to conidiogenous cells that occur singly or more commonly in clusters. *Conidiogenous cells* hyaline, smooth-walled, arising from superficial mycelium, ampulliform, 7–18 × 4–6 µm, consisting of an obovoid venter, 4–10 µm long, and cylindrical neck, 3–7 µm long, not flared, 2–2.5 µm diam. *Conidia* hyaline, smooth- and thin-walled, guttulate, aseptate, shortly clavate, apex obtuse, tapering to truncate hilum, 1–2 µm diam, produced in short, false chains that slime down into a globoid mass, (3–)3.5–4(–5) × (2–)2.5(–3) µm.

Culture characteristics — Colonies erumpent, spreading, with abundant aerial mycelium on MEA, moderate aerial mycelium on PDA, and sparse aerial mycelium on OA, with even, lobate margins, reaching 30 mm diam after 2 wk at 25 °C. On MEA surface dirty white, reverse ochreous; on PDA surface and reverse pale luteous; on OA surface dirty white.

Colour illustrations. Collection site in Germany, with dead *Sambucus nigra* in foreground. *Ascomata in vivo*; setae and ascospores; asci and ascospores; conidiogenous cells giving rise to conidia; conidia. Scale bars = 350 µm (ascomata), 10 µm (all others).

Typus. GERMANY, Bayern, Markredwitz (east of Bayreuth, Bayern), 600 m a.s.l., N50°00'23.9" E12°06'07.7", on dead stem base of *Sambucus nigra* (Adoxaceae), 5 Dec. 2020, M. Reul, HPC 3546 = #8494 (holotype CBS H-24845 culture ex-type CPC 40384 = CBS 148306, ITS, LSU, *rpb2* and *tub2* sequences GenBank OK664749.1, OK663788.1, OK651179.1 and OK651217.1, MycoBank MB 841832).

Additional material examined. GERMANY, Nordrhein-Westfalen, Duisburg, Ruhrdeich, near Werthacker (NRW), 35 m a.s.l., N51°26'53.68" E6°48'04.4", on dead stem of *Sambucus ebulus*, 1 Dec. 2020, K. Müller, HPC 3545 = #SI 09/2020. – NETHERLANDS, Utrecht Province, Raaphof, near Bunnik, 2 m a.s.l., N52°03'06" E5°12'41", on dead stem of *Arctium*, 24 June 2015, E.R. Osieck, (coll. #3203-9i).

Notes — *Paramicrothecium* is phylogenetically closely related but distinct from *Pseudomicrothecium* (non-ostiolate ascomata, 2-spored asci, smooth-walled ascospores with an indistinct germ pore at each end), and *Microthecium* (ascomata ostiolate or not aseptate (i.e., glabrous or surrounded by hyphae-like hairs sensu Marin-Felix et al. (2018), asci 8-spored, smooth to wrinkled, with germ pores at each end, and phialidic asexual morph) (Marin-Felix et al. 2018).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *Thielavia basicola* (strain CBS 178.82, GenBank NR_165582.1; Identities = 490/544 (90 %), 25 gaps (4 %)), *Microthecium quadrangulatum* (strain Y-049, GenBank MN562054.1; Identities = 387/439 (88 %), 18 gaps (4 %)), and *Microthecium zobellii* (strain CBS 341.73, GenBank MK926782.1; Identities = 387/439 (88 %), 18 gaps (4 %)). Closest hits using the LSU sequence are *Thielavia basicola* (strain CBS 178.82, GenBank MK926783.1; Identities = 515/526 (98 %), no gaps), *Melanospora subterranea* (= *Pseudomicrothecium subterraneum*) (voucher BJTC FAN1001, GenBank NG_060274.1; Identities = 808/830 (97 %), three gaps (0 %)), and *Melanospora subterranea* (voucher BJTC fan1001, GenBank JN247804.1; Identities = 808/830 (97 %), three gaps (0 %)). The best hit using the *rpb2* sequence had highest similarity to *Thielavia basicola* (strain CBS 178.82, GenBank MK876745.1; Identities = 728/838 (87 %), no gaps). No significant hits were obtained when the *tub2* sequence was used in blastn and megablast searches.

Supplementary material

See the phylogenetic tree provided with the supplementary material FP1318.

Pedro W. Crous, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl
Krimhilde Müller, Falkstraße 103, D-47058 Duisburg, Germany; e-mail: Krimhilde.M@t-online.de
Klaus Siepe, Geeste 133, D-46342 Velen, Germany; e-mail: k.siepe@web.de
Matthias Reul, Ostenstraße 19, D-95615 Marktredwitz, Germany; e-mail: matthias-reul@live.de
Eduard R. Osieck, Jkvr. C.M. van Asch van Wijcklaan 19, 3972 ST Driebergen-Rijsenburg, Netherlands; e-mail: panurus@ziggo.nl



Fungal Planet 1330 – 24 December 2021

***Ophioceras junci* Crous & Osieck, sp. nov.**


Etymology. Name refers to the host genus *Juncus* from which it was isolated.

Classification — *Ophioceraaceae*, *Magnaporthales*, *Sordariomycetes*.

Ascomata solitary, brown, immersed, globose to lobesided base, 100–150 µm diam, surface of *textura epidermoidea*; wall of 4–6 layers of brown *textura angularis*, with elongated, cylindrical neck, up to 300 µm long, 30–70 µm wide, brown at base, pale brown at apex, that becomes slightly swollen, filled with cylindrical, hyaline periphyses, 2–2.5 µm diam, abundant in central ostiole; neck covered in short, 5–15 × 2–2.5 µm hyphae with obtuse ends, smooth-walled, brown, becoming subhyaline towards apex. **Paraphyses** extending above asci, hyaline, smooth-walled, subcylindrical, constricted at septa, up to 90 µm tall, 4–10 µm wide. **Asci** unitunicate, stipitate with foot, subcylindrical, hyaline, straight to slightly curved, apex obtuse, apical mechanism refractive, not staining in Melzer's reagent, with eight multiseriate ascospores, 35–60 × 6–7 µm. **Ascospores** narrowly fusoid, widest in middle, tapering to acutely rounded apices, at times more rounded at base than apex, straight to curved, guttulate, 3-septate, slightly constricted at septa, hyaline, smooth-walled, (25–)27–30(–35) × 3(–3.5) µm.

Culture characteristics — Colonies flat, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 8 mm diam after 7 d at 25 °C. On MEA and OA surface and reverse buff; on PDA surface and reverse hazel in middle, but in outer region.

Typus. NETHERLANDS, Utrecht Province, Nieuw Wulven, near Houten, 1.5 m a.s.l., N52°03'45" E05°10'33", on dead culm of *Juncus effusus* (*Juncaceae*), 24 June 2021, E.R. Osieck, HPC 3659 = WI-41/#4274 (holotype CBS H-24892, culture ex-type CPC 42234 = CBS 148450, ITS, LSU, *actA* and *rpb1* sequences GenBank OK664750.1, OK663789.1, OK651137.1 and OK651155.1; *ibid.*, culture CPC 42235 = CBS 148451, ITS, LSU, *actA* and *rpb1* sequences GenBank OK664751.1, OK663790.1, OK651138.1 and OK651156.1, MycoBank MB 841833).



Colour illustrations. *Juncus effusus* at Nieuw Wulven, near Houten. Ascomatal necks *in vivo*; broken ascoma; asci with ascospores; ascospores. Scale bars = 70 µm (ascomatal necks), 10 µm (all others).

Notes — *Ophioceras* was recently treated by Jiang et al. (2021). Based on the species presently accepted, *O. junci* should be compared with *O. castillensis* (ascospores fusoid, hyaline to pale brown, 3-septate, 29–40 × 4–5.5 µm, ascomata arranged in large clusters, known from bark or wood in Costa Rica, Nicaragua, and Puerto Rico; Jiang et al. 2021, no ITS sequence available), from which it differs in ascomatal arrangement, and having smaller ascospores. Phylogenetically, it forms a sister lineage to *O. aquaticum*. This represents the first *Ophioceras* species described from *Juncaceae* as host (most occur on wood, but five species are known from monocotyledons like *Juncus*) and the second from Europe (*O. leptosporum* on herbs and herbaceous debris; Shearer et al. 1999).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *Ophioceras aquaticum* (strain IFRDCC 3091, GenBank NR_165842.1; Identities = 447/515 (87 %), 23 gaps (4 %)), *Ophioceras freycinetiae* (strain CBS 146781, GenBank NR_173031.1; Identities = 427/508 (84 %), 21 gaps (4 %)), *Pestalotiopsis mangifolia* (voucher INBio 820B, GenBank KU204659.1; Identities = 426/508 (84 %), 22 gaps (4 %)) and *Ophioceras leptosporum* (strain CPC 39147, GenBank MW883435.1; Identities = 421/506 (83 %), 19 gaps (3 %)). The ITS sequences of CPC 42234 and 42235 are identical (498/498 nucleotides, no gaps). Closest hits using the **LSU** sequence are *Ophioceras aquaticum* (strain IFRDCC 3091, GenBank NG_067778.1; Identities = 445/466 (95 %), two gaps (0 %)), *Ophioceras commune* (strain BCC3328, GenBank DQ341503.1; Identities = 439/470 (93 %), two gaps (0 %)), and *Ophioceras hongkongense* (strain HKUCC3624, GenBank DQ341509.1; Identities = 442/475 (93 %), six gaps (1 %)). The LSU sequences of CPC 42234 and 42235 are identical (377/377 nucleotides, no gaps). No significant hits were obtained when the **actA** sequence was used in blastn and megablast searches. The **actA** sequences of CPC 42234 and 42235 are identical (381/381 nucleotides, no gaps). Closest hits using the **rpb1** sequence had highest similarity to *Ophioceras dolichostomum* (strain CBS 114926, GenBank JX134731.1; Identities = 459/574 (80 %), 12 gaps (2 %)), *Ophioceras commune* (strain M91, GenBank JX134729.1; Identities = 455/572 (80 %), 11 gaps (1 %)), and *Ophioceras leptosporum* (strain CBS 894.70, GenBank JX134732.1; Identities = 394/523 (75 %), 14 gaps (2 %)). The **rpb1** sequences of CPC 42234 and 42235 are identical (541/541 nucleotides, no gaps).

The first of 26 equally most parsimonious trees obtained from a phylogenetic analysis of the *Ophioceras* ITS nucleotide alignment. The tree was rooted to *Harzia macrospora* (culture CBS 122807; GenBank NR_161008.1) and the scale bar indicates the number of changes. Parsimony bootstrap support values from 1000 replicates and > 79 % are shown at the nodes and the treated species is highlighted with a coloured box and **bold** text. GenBank accession (superscript) and/or culture collection/voucher numbers (in **bold** face when having a type status) are indicated for all species. Branches present in the strict consensus tree are thickened. Alignment statistics: 20 strains including the outgroup; 594 characters including alignment gaps analysed: 197 constant, 147 variable and parsimony-uninformative and 250 parsimony-informative. Tree statistics: Tree Length = 791, Consistency Index = 0.800, Retention Index = 0.865, Rescaled Consistency Index = 0.692. The alignment and tree were deposited in figshare.com (10.6084/m9.figshare.16918582).

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Eduard R. Osieck, Jkvr. C.M. van Asch van Wijklaan 19, 3972 ST Driebergen-Rijsenburg, Netherlands; e-mail: panurus@ziggo.nl



Fungal Planet 1331 – 24 December 2021

Paradinemasporium* Crous & Osieck, gen. nov.Etymology.* Name refers to its similarity to *Dinemasporium*.Classification — *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*.

Conidiomata stromatic, acervuloid, mostly solitary on host, pale brown with luteous mucoid conidial mass, setose; basal stroma of subglobose cells, excipulum absent. *Setae* interspersed, arising from basal stroma, straight, multi-septate, brown, unbranched, thick-walled, fertile at apex, with flared

collarette. *Conidiophores* lining cavity, subcylindrical, hyaline, smooth-walled, branched, septate. *Conidiogenous cells* phialidic, subcylindrical, hyaline to subhyaline, smooth-walled; apex with flared collarette. *Conidia* fusoid to subcylindrical, hyaline, smooth-walled, guttulate, 1-septate, apex with 2–3 excentric appendages; base with a single central appendage.

Type species. *Paradinemasporium junci* Crous & Osieck
MycoBank MB 841834.

Paradinemasporium junci* Crous & Osieck, sp. nov.Etymology.* Name refers to the host genus *Juncus* from which it was isolated.

Conidiomata stromatic, acervuloid, 100–500 µm diam, mostly solitary on host, to aggregated (PNA), pale brown with luteous mucoid conidial mass, setose; basal stroma of subglobose cells, lateral excipulum absent. *Setae* interspersed, arising from basal stroma, straight, multi-septate, brown, unbranched, thick-walled, 100–250 µm long, 4–6 µm wide, fertile at apex, with flared collarette. *Conidiophores* lining cavity, subcylindrical, hyaline, smooth-walled, branched, septate, 40–80 × 3–4 µm. *Conidiogenous cells* phialidic, subcylindrical, hyaline to subhyaline, smooth-walled, 15–27 × 2.5–3.5 µm; apex with flared collarette which is 3–5 µm diam. *Conidia* (12–)13–14 × (2.5–)3 µm, fusoid to subcylindrical, hyaline, smooth-walled, guttulate, medianly 1-septate, apex with 2–3 excentric appendages, 7–10 µm long; base with a single central appendage, 9–12 µm long.

Culture characteristics — Colonies flat, spreading, with moderate aerial mycelium and even, lobate margin, reaching 50 mm diam after 2 wk at 25 °C. On MEA surface cinnamon to isabelline, reverse isabelline; on PDA surface and reverse cinnamon; on OA surface cinnamon.

Typus. NETHERLANDS, Overijssel Province, Engbertsdijkswenen, near Vriezenveen, 13 m a.s.l., N52°26'53" E06°40'02", on dead culm of *Juncus effusus* (*Juncaceae*), 9 Mar. 2021, *E.R. Osieck*, HPC 3612 = WI-31/#4227 (holotype CBS H-24856, culture ex-type CPC 41315 = CBS 148317, ITS, LSU and *tef1* (second part) sequences GenBank OK664752.1, OK663791.1 and OK651199.1, MycoBank MB 841835).

Additional material examined. NETHERLANDS, Overijssel Province, Engbertsdijkswenen, near Kloosterhaar, 17 m a.s.l., N52°28'57", E06°40'02", on dead culm of *Juncus effusus*, 9 Mar. 2021, *E.R. Osieck*, HPC 3611 = WI-30/#4226, culture CPC 41343, ITS and LSU sequences GenBank OK664754.1 and OK663793.1; Utrecht Province, Utrecht, Leersum, Leersumse Plassen, 8 m a.s.l., N52°02'28" E05°26'26", on dead culm of *Juncus effusus*, 26 Jan. 2021, *E.R. Osieck*, HPC 3584 = WI-16/#4212, culture CPC 41003, ITS, LSU and *tef1* (second part) sequences GenBank OK664753.1, OK663792.1 and OK651200.1.


Colour illustrations. *Juncus effusus* at Engbertsdijkswenen, near Vriezenveen, Overijssel Province. *Conidiomata* on PNA; seta; conidiogenous cells giving rise to conidia; conidia. Scale bars = 10 µm.

Notes — *Paradinemasporium* forms a well-defined distinct clade related to *Pseudodinemasporium* and *Neopseudolachnella* (Crous et al. 2012, Hashimoto et al. 2015). It most closely resembles *Dinemasporium* in conidial morphology and appendages, but also *Neopseudolachnella* in having sporodochia that lack a well-defined lateral excipulum. *Paradinemasporium junci* resembles *D. strigosum* (common on monocotyledons) but the latter has shorter conidia and a single appendage at each end (Duan et al. 2007).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence of CPC 41315 had highest similarity to *Neopseudolachnella uniseptata* (voucher HHUF 29728, GenBank NR_154225.1; Identities = 402/437 (92 %), eight gaps (1 %)), *Neopseudolachnella magnispora* (voucher HHUF 29977, GenBank NR_154224.1; Identities = 426/469 (91 %), 16 gaps (3 %)), and *Pseudolachnella complanata* (voucher HHUF 28282, GenBank NR_154278.1; Identities = 333/368 (90 %), 12 gaps (3 %)). The ITS sequence of CPC 41315 differs two gaps (462/464 nucleotides, including two gaps) from CPC 41003 and two nucleotides and one gap from CPC 41343 (464/467 nucleotides, including one gap). Closest hits using the LSU sequence of CPC 41315 are *Neopseudolachnella magnispora* (voucher HHUF 29977, GenBank NG_059405.1; Identities = 760/774 (98 %), two gaps (0 %)), *Neopseudolachnella uniseptata* (voucher HHUF 29728, GenBank NG_059406.1; Identities = 759/773 (98 %), one gap (0 %)), and *Pseudolachnella fusiformis* (voucher HHUF 29725, GenBank AB934056.1; Identities = 758/773 (98 %), two gaps (0 %)). The LSU sequence of CPC 41315 is identical to those of CPC 41003 and CPC 41343 (both 773/773 nucleotides, no gaps). Closest hits using the *tef1* (second part) sequence of CPC 41315 had highest similarity to *Dictyochoaeta cangshanensis* (voucher MFLU 18-1614, GenBank MN194083.1; Identities = 802/845 (95 %), no gaps), *Neopseudolachnella magnispora* (voucher HHUF 29977, GenBank AB934092.1; Identities = 801/852 (94 %), no gaps), and *Dictyochoaeta submersa* (voucher MFLU 18-2321, GenBank MN194082.1; Identities = 764/814 (94 %), no gaps). The *tef1* sequence of CPC 41315 differs one nucleotide (810/811 nucleotides, no gaps) from CPC 41003.

Supplementary material**FP1331** Phylogenetic tree.

Cippumomyces mortalis



Fungal Planet 1332 – 24 December 2021

Cippumomyces* Crous, Overton & Ricci, *gen. nov.

Etymology. Name refers to the fact that it was isolated from a granite tombstone ‘*cippum*’.

Classification — *Neoantennariellaceae*, *Capnodiales*, *Dothideomycetes*.

Conidiomata aggregated in a brown stroma, visible on surface via exuding mucoid spore droplets (on OA and SNA). **Mycelium** consisting of brown, verruculose, branched, septate, hyphae, encased in mucoid sheath, forming chains of intercalary conidiomata, globose to ellipsoid, with dividing walls dissolving; wall of 2–3 layers of brown, verruculose *textura angularis*; open via irregular rupture. **Conidiogenous cells** arising from basal

layer of conidioma, brown, cylindrical, giving rise to a conidium that is hyaline, globose, encased in thick mucoid sheath; conidium content dividing into several globose conidia, hyaline, smooth, with 2, 4, 8 or multiple endoconidia; original conidial wall dissolving at maturity, but endoconidia remain encased in mucoid sheath. **Conidia** hyaline, smooth, thin-walled, ellipsoid to globose, encased in thick mucoid sheath. Mature conidia become brown, verruculose, muriformly septate, and frequently anastomose; conidia exude in a mucoid droplet.

Type species. *Cippumomyces mortalis* Crous, Overton & Ricci
Mycobank MB 841836.

Cippumomyces mortalis* Crous, Overton & Ricci, *sp. nov.

Etymology. Name refers to the association of gravestones and death = *mors*.

Conidiomata aggregated in a brown stroma, visible on surface via exuding mucoid spore droplets (on OA and SNA). **Mycelium** consisting of brown, verruculose, branched, septate, 3–6 µm diam hyphae, encased in mucoid sheath, forming chains of intercalary conidiomata, globose to ellipsoid, 30–70 µm diam, with dividing walls dissolving, forming longer conidiomata that can be up to 250 µm wide, and 30–50 µm tall; wall of 2–3 layers of brown, verruculose *textura angularis*; open via irregular rupture. **Conidiogenous cells** arising from basal layer of conidioma, brown, cylindrical, giving rise to a conidium that is hyaline, globose, encased in thick mucoid sheath; conidium content dividing into several globose endoconidia, hyaline, smooth, 5–10 µm diam, with 2, 4, 8 or multiple endoconidia; original conidial wall dissolving at maturity, but endoconidia, 3–5 × 2.5–5 µm, remain encased in mucoid sheath. **Conidia** hyaline, smooth, thin-walled, ellipsoid to globose, encased in thick mucoid sheath. Mature conidia become brown, verruculose, muriformly septate, and frequently anastomose; conidia exude in a mucoid droplet.

Culture characteristics — Colonies erumpent, spreading, with sparse aerial mycelium and smooth, lobate margin, reaching 4–6 mm diam after 7 d at 25 °C. On MEA and PDA surface and reverse isabelline; on OA surface brown vinaceous. Good growth and sporulation also observed on Acidified Rose-Bengal Agar, but the mucoid sheath tends to be less well-developed, the endoconidia less clustered, and conidiogenous cells not observed.

Typus. USA, Clinton County Pennsylvania, on tombstone, adjacent to Lock Haven University Campus, 14 Oct. 2020, G.M. Ricci (holotype CBS H-24893, culture ex-type CPC 41588 = CBS 148452, ITS, LSU, *actA*, *cmdA*, *rpb2*, *tef1* (first part) and *tub2* sequences GenBank OK664755.1, OK663794.1, OK651139.1, OK651180.1, OK651195.1 and OK651218.1; *ibid.*, culture CPC 41587 = CBS 148453, ITS, LSU, *actA*, *cmdA*, *rpb2*, *tef1* (first part) and *tub2* sequences GenBank OK664756.1, OK663795.1, OK651140.1, OK651146.1, OK651181.1, OK651196.1 and OK651219.1, MycoBank MB 841837).

Colour illustrations. Tombstones, USA. Conidiomata on MEA; conidiogenous cells giving rise to primary conidia; primary conidia giving rise to endoconidia, both with mucoid sheath. Scale bars = 70 µm (conidiomata), 10 µm (all others).

Notes — *Cippumomyces* adds yet another slow-growing yeast-like genus to the *Neoantennariellaceae*. It is characterised by forming a stroma containing conidiomata that are initially linked in chains in a mycelial network, with conidiomatal walls that dissolve at maturity, forming larger conidiomata. Primary conidia formed on conidiogenous cells are globose, and give rise to endoconidia. At first sight, the primary conidia containing propagules are rather reminiscent of asci with ascospores, but as these were found to form on stalks, that were interpreted as conidiogenous cells forming from the inner wall of conidiomata, they are regarded as primary conidia. Endoconidia vary in number and increase in number at maturity.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence of CPC 41587 had highest similarity to *Ramimonilia apicalis* (strain CBS 118327, GenBank NR_144959.1; Identities = 437/474 (92 %), seven gaps (1 %)), *Neocatenulostroma germanicum* (strain CBG, GenBank MK622897.1; Identities = 435/492 (88 %), 21 gaps (4 %)), and *Aulographina pinorum* (strain CBS 302.71, GenBank GU214622.1; Identities = 482/547 (88 %), 24 gaps (4 %)). The ITS sequences of CPC 41587 and 41588 were identical (535/535 nucleotides, no gaps). Closest hits using the **LSU** sequence of CPC 41587 are *Readeriellipsoidia fuscoporiae* (strain CBS 139900, GenBank NG_058161.1; Identities = 691/728 (95 %), five gaps (0 %)), *Readeriellipsoidia guyanensis* (as *Readeriella guyanensis*; strain CBS 117550, GenBank FJ493211.1; Identities = 691/728 (95 %), five gaps (0 %)), and *Nothophaeothea mirabilensis* (strain CBS 146980, GenBank NG_076740.1; Identities = 694/732 (95 %), nine gaps (1 %)). The LSU sequences of CPC 41587 and 41588 were identical (721/721 nucleotides, no gaps). No significant hits were obtained when the **actA**, **cmdA**, **rpb2**, **tef1** (first part) and **tub2** sequences were used in blastn and megablast searches. The **actA**, **rpb2**, **tef1** (first part) and **tub2** sequences of CPC 41587 and 41588 were identical (558/558, 817/817, 422/422 and 410/410 nucleotides, no gaps).

Supplementary material

FP1332 Phylogenetic tree.

Pedro W. Crous & Johannes Z. Groenewald, Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands; e-mail: p.crous@wi.knaw.nl & e.groenewald@wi.knaw.nl

Gianna M. Ricci, undergraduate student, 205 East Campus Science Center, Lock Haven University, Department of Biology, Lock Haven PA 17745, USA; e-mail: riccigiannam@gmail.com

Barrie E. Overton, 205 East Campus Science Center, Lock Haven University, Department of Biology, Lock Haven, PA 17745, USA; e-mail: boverton@lockhaven.edu

Amanita exilis



Fungal Planet 1333 – 24 December 2021

Amanita exilis* Loizides, Biketova, Bellanger & P.-A. Moreau, sp. nov.Etymology.* *Exilis* (Latin) = slender, referring to the species' aspect.Classification — *Amanitaceae*, *Agaricales*, *Agaricomycotina*.

Basidiomata with a schizohymenial ontogeny, initially enclosed in universal veil. *Pileus* 30–70 mm diam, at first hemispherical to convex, soon expanding to applanate and somewhat depressed in the middle, never umbonate, smooth, dry, silver grey, warm grey or bistre brown (colour chart: Anonymous 1969), sometimes faded towards the centre, usually devoid of velar remains but occasionally with a large continuous patch of white veil attached; margin markedly striate-furrowed, not translucent. *Lamellae* free to finely adnexed, moderately crowded and intercepted by sparse lamellulae, fragile, white; edges concolourous, smooth to slightly crenate. *Stipe* typically long and slender, 80–120 × 5–10 mm, exannulate, ± equidiametrical or slightly enlarged at the base, white, covered in concolourous meandering bands gradually turning pale grey or grey-brown forming a 'snakeskin' pattern. *Volva* deep and tightly adhered to the stipe (type III), membranous, fragile, rooted into the substrate, white or dirty white, sometimes with ochraceous brown stains. *Context* white, with a mild odour. *Spore deposit* white. *Basidiospores* (10–)11–14(–15) × (8.5–)9–10(–10.5) µm (av. = 12.4 × 9.6; Q = 1.11–1.57; Qm = 1.29), ovoid to broadly ellipsoid or ellipsoid, hyaline, smooth, inamyloid, with one large guttule and a short hilar appendage. *Basidia* tetrasporic or bisporic, rarely also monosporic, 34–84 × 13–18 µm, thick-walled (0.5–1 µm), broadly clavate at first but in maturity becoming elongated to pyriform and tapering to a long pedestal, filled with coarse vacuolar content refractive in 5 % KOH; sterigmata 3–7 µm long. *Marginal cells* abundant, 26–67 × 12–20 µm, thick-walled (0.5–1 µm), polymorphic, ranging from clavate to obpyriform, subcapitate, subcylindrical, or occasionally catenulate, 1–2(–3)-septate and sometimes constricted at the septa. *Clamp connections* not seen. *Subhymenium* composed of subspherical, angular or allantoid cells, 12–17 µm wide. *Pileipellis* a trichoderm of intertwined, frequently branched and thick-walled hyphae terminating in partially erect cylindrical ends, but scattered clavate or fusiform elements up to 26 µm wide also present resembling a hymeniderm in parts. *Stipitipellis* composed of tightly packed, ± parallel and thick-walled hyphae; acrophysalides rare, 17–32 µm diam. *Universal veil* filamentous, composed of tightly packed, thick-walled and sparsely septate cylindrical hyphae, 5–8 µm wide, sometimes slightly constricted at the septa, very rarely terminating in enlarged clavate, pyriform or globose, thick-walled acrophysalidic cells < 26 µm diam.

Habit, Habitat & Distribution — Known from the eastern Mediterranean basin, where it appears solitary or in small groups between October and December, sporulating on calcareous soil under *Quercus coccifera* subsp. *calliprinos*. Note that the

taxonomic status of Cypriot and Israeli oak populations is not yet entirely clear (see Vitellii et al. 2017, Hipp et al. 2020); until further clarified we follow Denk & Grimm (2010).

Typus. CYPRUS, Kelefos, on calcareous soil, in mixed *Pinus brutia* forest under *Quercus coccifera* subsp. *calliprinos* (Fagaceae), 24 Nov. 2017, M. Loizides (holotype in Herbarium of the Faculty of Pharmacy of Lille: LIP 0002174; isotype in herb. pers. M. Loizides: ML711142AH, ITS and LSU sequences GenBank MH603610 and MZ508450, respectively, MycoBank MB 841766).

Additional materials examined. *Amanita exilis*: CYPRUS, Kelefos, under *Quercus coccifera* subsp. *calliprinos*, 31 Oct. 2012, M. Loizides, ML2110103 (ITS GenBank MH603597). ISRAEL, Carmel Mount, Derech Nof HaCarmel, under *Q. coccifera* subsp. *calliprinos*, 28 Nov. 2012, Y. Cherniavsky, AB A12-058-2 (ITS sequence MZ508449). *A. lividopallescens*: ISRAEL, Carmel Mount, Derech Nof HaCarmel, under *Q. coccifera* subsp. *calliprinos*, 28 Nov. 2012, Y. Cherniavsky, AB A12-058-1 (ITS sequence GenBank MZ576431).

Notes — *Amanita exilis* is characterised by tall and slender basidiomata usually with a stipe twice as long as the pileus width, a filamentous veil and ± ellipsoid spores. This species was previously reported as '*Amanita* sp. nov. 3' in Loizides et al. (2018) and forms a well-supported clade (0.96/1) in sect. *Vaginatae*, distant from its closest neighbour *A. lividopallescens* by 15 substitutions and 8 indels (see supplementary material). The latter can sometimes occur in the same habitat and has similar spores in size, shape and quotient, but produces larger basidiomata with ochraceous brown or pale brown pileal colours, and has a spacious saccate volva (type II) with large acrophysalides up to 182 × 48 µm (Tulloss 1994, Vizzini et al. 2016). *Amanita bertaultii*, described from Mediterranean *Pinus* and *Quercus* forests based on *A. mairei* s.str. Bertault (1965) from Morocco, also has bands on the stipe and a filamentous veil (Contu 1985a, b). This enigmatic species, whose type material could not be located (probably in CAG), was later recombined by its author as a variety of *A. mairei* (Contu 2000) and was compared to *A. lividopallescens* by its expanded volva, dark grey pileal colour with ochraceous tinges and ecology under holm oak. It differs from *A. exilis* in the absence of bistre or silver-grey tinges (excluded by Contu, 1985a: '*haud argenteus!*'), and smaller spores of 10–12(–13) × 7–9 µm. Also in series *Mairei*, *A. mairei* and *A. supravolvata* have a grey pileus without umbo, ellipsoid spores and a filamentous veil, but produce more robust basidiomata strictly associated with pines, while *A. cistetorum* is associated with *Cistus* and *Halimium*, has a grey stipe always shorter than the diameter of the pileus and typically truncate lamella edges (Contu & Pacioni 1998, Hanss & Moreau 2020, Leonardi et al. 2020). *Amanita huijsmanii*, described from broadleaved woods in France, also has a filamentous veil and ellipsoid to ovoid spores measuring 12.5–14.5 × 9–10.5 µm according to Massart & Rouzeau (1989), or 9–11 × 8–9 µm according to Hanss & Moreau (2020).

(text continues on Supplementary material page FP1333)

Colour illustrations. Holotype collection area at Kelefos. Collection ML2110103; holotype collection LIP 0002174; basidiospores; stipitipellis with rare acrophysalides; hymenium. Scale bars = 50 mm (specimens *in situ*), 10 µm (basidiospores, stipitipellis and hymenium).

Supplementary material**FP1333** Phylogenetic tree.

Michael Loizides, P.O. Box 58499, 3734 Limassol, Cyprus; e-mail: michael.loizides@yahoo.com

Alona Yu. Biketova, Institute of Biochemistry, Biological Research Centre of the Eötvös Lóránd Research Network, Temesvári Blvd. 62, Szeged, H-6726, Hungary; and Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3DS, UK; e-mail: alyona.biketova@gmail.com

Jean-Michel Bellanger, CEF, CNRS, Université de Montpellier, Université Paul-Valéry Montpellier 3, EPHE, IRD, INSERM, 1919 route de Mende, F-34293 Montpellier Cedex 5, France; e-mail: jean-michel.bellanger@cefe.cnrs.fr

Pierre-Arthur Moreau, Université de Lille, Faculté de pharmacie de Lille, EA 4483, F-59000 Lille, France; e-mail: pierre-arthur.moreau@univ-lille2.fr

Aspergillus arizonicus



Fungal Planet 1334 – 24 December 2021

***Aspergillus arizonicus* Jurjević, Glässnerová, Yaguchi & Hubka, sp. nov.**

Etymology. Pertaining to Arizona, the state in the USA where the species was collected.

Classification — *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*.


Micromorphology (on malt extract agar (MEA), 25 °C after 14 d): *Conidiophores* borne from aerial hyphae; *conidial heads* radiate to columnar, greyish green, uniseriate, occasionally nodding; *stipes* hyaline, smooth, aseptate or sparsely septate, (30–)50–150(–200) × 4–7(–8.5) µm; *vesicle* subclavate, clavate or spatulate, (6–)8–15(–18) µm diam; *phialides* ampulliform, 4.5–7 µm long, covering half to two thirds of the vesicle surface; *conidia* globose to subglobose, smooth to delicately roughened, greenish grey in mass, 2.5–3 µm diam (av. = 2.6 ± 0.2). *Sexual morph* unknown.

Culture characteristics — (in darkness, 25 °C after 7 d): Colonies on MEA 52–57 mm diam, lanose to floccose, raised, greenish grey (26C2–3, colour based on Komerup & Wanschner (1967)), sporulation abundant, exudate absent, soluble pigment absent, margins entire, reverse greyish yellow (4B4). Colonies on Czapek yeast autolysate agar (CYA) 38–47 mm diam, lanose to floccose, slightly raised, radially furrowed, white, sometimes with greenish grey (26D2–E2) sectors, sporulation sparse, abundant in the greyish-green sectors when present, exudate absent, soluble pigment absent, margins delicately filiform, reverse pale yellow (3A3). Colonies on Czapek agar (CZA) 26–31 mm diam, lanose, radially furrowed, white, conidial heads absent or rare, exudate absent, soluble pigment absent, margins entire, reverse yellowish white (3A2). Colonies on CYA with 20 % sucrose (CY20S) 39–42 mm diam, lanose, raised radially furrowed, white, sometimes with greenish grey sectors, sporulation sparse, abundant in the greyish green sectors when present, exudate absent, soluble pigment absent, margins entire, reverse greyish yellow (4B5). Colonies on oatmeal agar OA 45–49 mm diam, flat, floccose with a broad submerged marginal zone, greenish grey (26B2), sporulation abundant, exudate absent, soluble pigment absent, margins entire, reverse yellowish white (2A2). Growth on MEA at 30 °C after 7 d faster than at 25 °C (> 65 mm, whole dish surface covered); at 37 °C comparable to 25 °C, 48–57 mm; at 40 °C 32–35 mm; at 42 °C 4–8 mm, no growth at 45 °C.

Typus. USA, Arizona, Tucson, indoor air of hospital, Dec. 2013, Ž. Jurjević (holotype PRM 954610, isotype PRM 954611, culture ex-type CCF 5341 = IFM 66805 = EMSL 2204 = CBS 148476, ITS, LSU, *benA*, *CaM* and *rbp2* sequences GenBank OK322364, OK321187, OK334128, OK334127 and OK334129, MycoBank MB 841359).

Notes — BLAST analyses with the sequences of the *Aspergillus* barcoding genes showed the greatest similarities with species belonging to series *Neoglabri* and *Unilaterales* (Houbraken et al. 2020). The multigene phylogeny resolved

A. arizonicus as a close relative of *A. elsenburgensis* (ser. *Neoglabri*). The latter can be differentiated by the production of ascomata in a homothallic manner and shorter conidiophores (Visagie & Houbraken 2020). The series *Neoglabri* and *Unilaterales* comprise seven asexual or heterothallic species, namely *A. brevistipitatus*, *A. conversis*, *A. marvanovae*, *A. nishimurae*, *A. tasmanicus*, *A. turcosus* and *A. unilateralis*. *Aspergillus brevistipitatus*, *A. conversis*, *A. tasmanicus* and *A. unilateralis* can be differentiated by roughened or echinulate conidia, and slower growth on MEA and CYA at 25 °C; *A. tasmanicus*, *A. turcosus* and *A. unilateralis* have shorter conidiophores that usually do not exceed 100 µm in length; *A. marvanovae*, *A. nishimurae*, *A. turcosus* and *A. unilateralis* are able to grow at 45 °C or higher temperatures in contrast to *A. arizonicus* (Hubka et al. 2013, 2017, Nováková et al. 2014).




A best scoring maximum likelihood tree based on the sequences of *benA*, *CaM* and *rbp2* showing the relationships of *A. arizonicus* to members of series *Neoglabri*, *Unilaterales* and *Fennelliarum*. The dataset contained 31 taxa and a total of 2 030 characters of which 561 were variable and 302 parsimony-informative. Partitioning scheme and substitution models for analyses were selected using PartitionFinder v. 2 (Lanfear et al. 2017): the TrNef+I+G was selected for the *benA* and *CaM* loci (partition 1) and TrN+G model for the *rbp2* gene (partition 2). The trees were constructed with IQ-TREE v. 1.4.4 (Nguyen et al. 2015). Support values at branches were obtained from 10³ standard bootstrap replicates; only support values ≥ 60 % are shown, the ex-type strains are indicated by superscript T. The tree is rooted with *A. spathulatus* NRRL 20549^T.

Colour illustrations. Hospital environment. Seven-day-old cultures of *Aspergillus arizonicus* grown at 25 °C on MEA, CYA and CZA (left to right); conidiophores; conidia. Scale bars = 10 µm.

Željko Jurjević, EMSL Analytical, Inc., 200 Route 130 North, Cinnaminson, NJ 08077 USA; e-mail: zjurjevic@emsl.com
Kateřina Glässnerová, Department of Botany, Faculty of Science, Charles University, Benátská 2, 128 01 Prague 2, Czech Republic; e-mail: Kacena.Glass@seznam.cz

Takashi Yaguchi, Medical Mycology Research Center, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba, 260-8673, Japan; e-mail: yaguchi@chiba-u.jp

Vit Hubka, Department of Botany, Faculty of Science, Charles University, Benátská 2, 128 01 Prague 2, Czech Republic, and Medical Mycology Research Center, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba, 260-8673, Japan; e-mail: vit.hubka@gmail.com



Fungal Planet 1335 – 24 December 2021

***Aureobasidium microtermitis* S. Tiwari & A. Baghela, sp. nov.**

Etymology. Name refers to the host termite genus *Microtermes* from which it was isolated.

Classification — *Sacrotheciaceae*, *Dothideales*, *Dothideomycetes*.

Conidial cells subglobose to ovoid, $2.71\text{--}5.82 \times 5.30\text{--}10.27\ \mu\text{m}$, after 7 d on potato dextrose agar (PDA). Synchronous polar budding is observed on PDA and yeast extract peptone agar (YPD). A few dark brown, short hyphae with intercalary chlamydospores observed after 10 d of growth on PDA. *Melanised chlamydospores* separate or intercalary, subglobose to globose, $6.73\text{--}10.26 \times 8.50\text{--}12.66\ \mu\text{m}$. The optimum growth temperature is $25\text{--}28\ ^\circ\text{C}$. Some growth observed at $15\ ^\circ\text{C}$ and $30\ ^\circ\text{C}$, but no growth at $4\ ^\circ\text{C}$ and $37\ ^\circ\text{C}$. Significant growth on high sugar-containing media (Harold's M40Y and M60Y) (Peterson et al. 2013). Capable of tolerating high salt concentrations ($5\text{--}15\%$ w/v NaCl).

Culture characteristics — Colonies on PDA attain 23 mm diam after 7 d growth at $25\ ^\circ\text{C}$, spreading, with irregular margins, creamy white peripherally and turning olivaceous black centrally. On yeast extract peptone (YPD) agar colonies attain 18 mm diam, circular, slightly convex, smooth, glistening, with an entire margin and a melanised ring in the centre.

Habitat & Distribution — *Aureobasidium microtermitis* was firstly isolated from the gut of the termite *Microtermes* sp. feeding on wood logs in a forest area in Rajpipla, Gujarat (India). The nectar of *Bombax ceiba* growing in Taljai Hills, Pune District, Maharashtra (India) was also another habitat for this fungus.


Typus. INDIA, Gujarat, Rajpipla district, N21.878760° E73.512894°, from the gut of a *Microtermes* sp. termite, 8 Sept. 2019, coll. S. Tiwari (holotype NFCCI 4901 preserved as metabolically inactive state at the National Fungal Culture Collection of India NFCCI, Agharkar Research Institute, Pune, India, ITS and LSU sequences GenBank MW276135 and MW276136, MycoBank MB 839078).

Additional material examined. INDIA, Maharashtra, Pune District, Taljai Hills, N18.478189° E73.843560°, in floral nectar of *Bombax ceiba*. 10 Feb. 2020, coll. A. Baghela (NFCCI 4935, ITS and LSU sequences GenBank MW692853 and MW031809).

Notes — The ascomycetous genus *Aureobasidium* is a member of the family *Sacrotheciaceae* within the class *Dothideomycetes* (Hymphries et al. 2017). Members of this genus are known for synchronous or percurrent conidiation, either on undifferentiated short denticles, intercalary or terminally on hyaline or melanised hyphae exhibiting variable culture characteristics (Zalar et al. 2008). *Aureobasidium microtermitis* was isolated from the gut of a *Microtermes* termite, and floral nectar of *Bombax ceiba*. Phylogenetic analysis using the LSU and ITS loci positions *A. microtermitis* close to *A. thailandense* but


distinguishable based on colony morphology and conidial cell size. Colonies of *A. thailandense* on PDA are reddish brown centrally, whereas *A. microtermitis* colonies are olivaceous black in the centre. The presence of melanised hyphae containing intercalary chlamydospores in the new species also sets it apart from *A. thailandense*. Conidia of *A. thailandense* are slightly larger ($3\text{--}10 \times 5\text{--}12\ \mu\text{m}$) (Peterson et al. 2013) than those of *A. microtermitis*.

Based on a megablast search of NCBI GenBank nucleotide database, the closest hits using the ITS sequence of NFCCI 4901 had highest similarity to *Aureobasidium thailandense* (strain PED7, GenBank KP860067.1; Identities = 555/566 (98 %), three gaps (0 %)), *Aureobasidium thailandense* (strain NRRL 58539, GenBank NR_147337.1; Identities = 563/580 (97 %), five gaps (0 %)), and *Aureobasidium* sp. (strain CLIB 3468, GenBank MT444031.1; Identities = 550/565 (97 %), four gaps (0 %)). The ITS sequences of NFCCI 4901 and 4935 show 100 % similarity (571/571). Closest hits using the LSU sequence of NFCCI 4901 are *Rhodotorula* sp. (strain fn_53, GenBank LC333510.1; Identities = 534/541 (99 %), no gaps), *Rhodotorula* sp. (strain YM25371, GenBank KC442285.1; Identities = 534/541 (99 %), no gaps), and *Aureobasidium* sp. (strain YM24994, GenBank KT239085.1; Identities = 533/540 (99 %), no gaps). The LSU sequences of NFCCI 4901 and 4935 show 100 % similarity (523/523).



Phylogenetic tree. The placement of *A. microtermitis* using a maximum-likelihood (ML) analysis of the combined ITS and LSU (D1/D2 domain) rRNA gene sequences employing the TIM2e+I+G4 model in IQ-TREE v. 1.6.8 (Nguyen et al. 2015). The scale bar indicates the expected number of substitutions per site. The numbers provided on branches are frequencies with which a given branch appeared in 1000 bootstrap replications. The tree was rooted with *Sydwia polyspora*. The new species proposed in the present study is highlighted and indicated in **bold** text.

Colour illustrations. Type locality – a small forest plot in Rajpipla, Gujarat, India. Host *Microtermes* sp.; colony on YPD agar; cells proliferating by polar budding; melanised chlamydospores; dark brown hyphae with intercalary chlamydospores. Scale bars = $20\ \mu\text{m}$ (budding cells), $10\ \mu\text{m}$ (hyphae and chlamydospores).



Fungal Planet 1336 – 24 December 2021

***Austroboletus yourkae* F.E. Guard, McMull.-Fish., Van Wyk, T. Lebel, & Halling, sp. nov.**

Etymology. Named for the Yourka Bush Reserve, on which this species occurs.

Classification — *Boletaceae*, *Boletales*, *Agaricomycetes*.

Pileus 2.5–5 cm broad, convex to plano-convex to plane, dry, to very slightly tacky at first, but not viscid or glutinous, tomentose, sometimes densely so, to matted tomentose, amber yellow (4B6,5,4; Kornerup & Wanscher 1983), even at margin (veil remnants apparently lacking) but with a white, barely visible sterile projection in one medium sized basidiome. **Flesh** white, unchanging. **Odour** fishy with a metallic tang, sometimes mild. **Taste** mild. **Tubes** adnexed, pale pink when young, vinaceous pink with age, bruising slightly brownish. **Stipe** 2.5–5 cm long, 0.6–1.5 cm broad, straight or curved, subclavate to equal and tapered at base, dry, coarsely alveolate-reticulate, white, developing orange stains at base in situ, slowly developing some pale brown stains at base with handling, with interior white, unchanging, pale yellow in base of mature basidiome, with white mycelium at base. **Basidiospores** 11.9–16.8 × 4.9–7 µm, (av. = 13.72 × 5.93, Q = 2.314, spores $n = 30$, specimens $n = 2$) subfusoid to ellipsoid, inequilateral in profile, hyaline in KOH, lightly dextrinoid in Melzer's, with distinct, uniform perforate-foveate ornamentation, occasionally barely ruminant. **Basidia** 30–40 × 10–12 µm, clavate, hyaline, four-sterigmate. **Tube trama** boletoid and divergent, becoming gelatinized with age, with hyphae 3–12 µm diam, hyaline in KOH and Melzer's; oleiferous elements occasionally present. **Hymenial cystidia** rare, present as cheilocystidia when young, hyaline, thin-walled, 30–45 × 7–10 µm, short subfusiform to fusiform, collapsing with age. **Pileus trama** interwoven, hyaline in KOH, with some few elements containing red brown colloidal content in Melzer's, smooth, thin-walled, 4–11 µm broad. **Pileipellis** a trichodermium of erect to suberect, tangled hyphae, hyaline in KOH, with colloidal red-brown content in Melzer's; elements cylindrical, smooth and thin-walled, 4–7 µm broad. **Stipitipellis** a hymeniform layer lining the reticulum, with long, filamentous, cylindrical, septate, smooth, thin-walled **caulocystidia** interspersed among basidiole-like elements. **Clamp connections** absent.


Habit, Habitat & Distribution — Gregarious on soil in dry sclerophyll woodland with *Corymbia intermedia*, *Eucalyptus resinifera*. So far, only known from northeast Queensland.

Colour illustrations. Sclerophyll forest of *Corymbia intermedia*, *Eucalyptus resinifera* (photo credit F. Guard). Solitary basidioma (Halling 10184, type); four basidiomata (Halling 10185); spores with DIC light microscope and with SEM; stipitipellis. Scale bars = 1 cm (basidiome habits), 20 µm (stipitipellis), 5 µm (spores).

Typus. AUSTRALIA, Queensland, Einasleigh Uplands, Yourka Bush Heritage Reserve, Junction Sandy Yard Creek Road & Valley Road, S17.98 E145.44, 600 m a.s.l., 18 Mar. 2019, R.E. Halling, Halling 10184 (holotype BRI AQ1024215, isotype NY02072697, LSU, *rbp1*, *rbp2* sequences GenBank MZ358814, MZ647699 and MZ647702, MycoBank MB 839206).

Additional material examined. AUSTRALIA, Queensland, Einasleigh Uplands, Yourka Bush Heritage Reserve, near Junction Themeda Track and Central Fire Track (3 Ways), S17.9421 E145.408, 622 m a.s.l., 19 Mar. 2019, R.E. Halling, Halling 10185 (NY02072669), LSU, *rbp1*, *rbp2* sequences GenBank MZ358815, MZ647700 and MZ647701.

Notes — In Australia, there are at least 17 species of *Austroboletus*, some of which are still undescribed (Halling et al., unpubl. data). The species that has the closest match to *A. yourkae*, in terms of morphology (spore ornamentation) and BLAST searches in GenBank for nrLSU and *rbp1*, *rbp2* gene sequences is *A. austrovirens* (Fechner et al. 2017). A RAxML analysis was conducted via Geneious v. 9.3.1 from concatenated LSU and *rbp1* data (<https://www.geneious.com>). *Austroboletus austrovirens* occurs in wetter sclerophyll habitats in northern Queensland but is easily distinguished by the green pigments on the pileus surface and stipe reticulum as well as the apricot orange pigments on the stipe surface between the reticulum ridges (Fechner et al. 2017, f. 1a).



Phylogram distinguishing *Austroboletus yourkae*. Phylogram from RAxML analysis generated via Geneious v. 9.3.1 of concatenated *rbp1* and LSU sequences for a selection of *Austroboletus* species, including some other bolete genera, with *Pulveroboletus* as outgroup. Thickened lines indicate ML support 0.90 and all bootstrap values are noted at branches. Scale bar shows substitutions per site. Novel sequences generated in this study are in **bold text**.

Frances E. Guard, Maleny, Queensland, Australia; e-mail: franguard@icloud.com
 Judson Van Wyk, Department of Plant Soil and Microbial Sciences, 1066 Bogue Street, Michigan State University, East Lansing, MI, 48824 USA; e-mail: vanwykju@msu.edu
 Teresa Lebel, State Herbarium of South Australia, Adelaide, South Australia 5000 Australia; e-mail: teresa.lebel@sa.gov.au
 Sapphire McMullan-Fisher, Royal Botanic Gardens, Victoria, Melbourne, Victoria, Australia; e-mail: ffunecol@gmail.com
 Roy E. Halling, Inst. Systematic Botany, New York Botanical Garden, 2900 Southern Blvd, Bronx, NY, USA 10458-5126; e-mail: rhalling@nybg.org

Biscogniauxia whalleyi



Fungal Planet 1337 – 24 December 2021

***Biscogniauxia whalleyi* N. Wangsawat, C. Phosri & N. Suwannasai, sp. nov.**

Etymology. Named after the British Mycologist Professor Anthony J.S. Whalley who has worked and devoted himself to studies on *Xylariales* for more than 50 years.

Classification — *Graphostromataceae*, *Xylariales*, *Sordariomycetes*.

Stromata applanate, orbicular to irregularly lobed, 0.5–3 cm long $\times$ 0.3–1 cm broad $\times$ 0.3 mm thick; outer dehiscing layer not seen, mature surface dark brick to sepia (Rayner 1970); carbonaceous immediately beneath surface and between perithecia; tissue beneath perithecia inconspicuous. **Perithecia** oblong to spherical, 0.2–0.3 mm high $\times$ 0.1–0.2 mm diam, opening centrally through individual ostioles. **Ostioles** punctate to slightly papillate by the presence of raised rim, inconspicuous, mostly evenly distributed, often plugged with greyish white substance. **Paraphyses** present. **Asci** 70–85 $\times$ 5–6 μ m, the spore-bearing parts 60–67 μ m long, the stipes 8–12 μ m long, with apical ring bluing in Melzer's iodine reagent, discoid, 0.5 μ m high $\times$ 2.5 μ m broad. **Ascospores** brown to dark brown, unicellular, ellipsoid, nearly equilateral, with broadly rounded ends, smooth, 7–8.5 $\times$ 4–5.5 μ m, with straight germ slit along the full length of the spore.

Culture characteristics — Colonies on potato dextrose agar (PDA) covering Petri dish after 3 wk at 30 °C, at first white, becoming isabelline to hazel; reverse fawn to brown. **Asexual morph** not observed.

Typus. THAILAND, Chaiyaphum, Phu Khiao Wildlife Sanctuary, on corticated wood, Aug. 2009, C. Phosri & N. Suwannasai (holotype SWUF13-85; ITS, LSU, α -actin, β -tubulin and *rpb2* sequences GenBank MW403821, MZ452416, MZ466385, MZ466386 and MZ466387, MycoBank MB 840293).

Notes — *Biscogniauxia whalleyi* is characterised by small ascospores (< 8.5 μ m long) with a straight germ slit along the spore length. Among previously known species, there are only four taxa, *B. arima*, *B. communapertura*, *B. mucigera* and *B. waitpela*, with ascospores less than 9 μ m long (Ju & Rogers 2001). Although the spore size of *B. whalleyi* is similar to *B. arima* (7.5–9 $\times$ 3–4 μ m), known from Mexico, ascospores of *B. arima* lack germ slits. *Biscogniauxia communapertura*, known from Brazil, has slightly larger spores (8–11.5 $\times$ 4.5–6 μ m) than *B. whalleyi*, and the germ slit of *B. communapertura* is straight and does not cover the entire spore length. Perithecia of *B. communapertura* are arranged in rosettes, and ascospores are discharged through a single ostiolar canal, differing from the present species. *Biscogniauxia whalleyi* is different from *B. mucigera* (8.5–10.5 $\times$ 5–5.5 μ m) in having smaller ascospores and lacking a hyaline sheath surrounding the ascospores. *Biscogniauxia waitpela* is another species possessing small ascospores, 6–7.5 $\times$ 3.5–4 μ m, but has an appendage and lacks a germ slit, which is different to *B. whalleyi*. In addition, the nucleotide sequences of five loci, ITS, LSU, α -actin, β -tubulin and *rpb2*, all confirmed *B. whalleyi* as separate from other known *Biscogniauxia* species.

Colour illustrations. Thailand, Chaiyaphum Province, Phu Khiao Wildlife Sanctuary, where the specimens were collected. From top to bottom: stroma (SWUF13-85); stroma with ostioles often plugged with greyish white substance; fungal culture on PDA; asci and ascospores with apical apparatus; ascospore with straight germ slit. Scale bars = 1 cm (fungal culture), 4 mm (upper stroma), 2 mm (lower stroma), 10 μ m (asci), 2 μ m (ascospore).

Supplementary material**FP1337** Phylogenetic tree.

Niwana Wangsawat, Department of Biology, Faculty of Science, Srinakharinwirot University, Bangkok, 10110 Thailand;

e-mail: niwana.wangsawat@swu.ac.th

Cherdchai Phosri, Biology programme, Faculty of Science, Nakhon Phanom University, Nakhon Phanom, 48000, Thailand; e-mail: Cherd.phosri@npu.ac.th

Ek Sangwichain, Department of Biology, Faculty of Science, Ramkhamhaeng University, Bangkok, 10240, Thailand; e-mail: eks@ru.ac.th

Nuttaporn Ruchikachorn, The Institute for the Promotion of Teaching Science and Technology, Bangkok, 10110, Thailand; e-mail: nruch@ipst.ac.th

Nuttika Suwannasai, Department of Microbiology, Faculty of Science, Srinakharinwirot University, Bangkok, 10110 Thailand; e-mail: nuttika@swu.ac.th



Fungal Planet 1338 – 24 December 2021

***Calocybella goethei* Angelini & Vizzini, sp. nov.**

Etymology. The epithet refers to the name of the private school (House of Goethe) in whose garden the species was found.

Classification — *Lyophyllaceae*, *Tricholomatineae*, *Agaricomycetes*.

Basidiomes small, collybioid. **Pileus** 15–25 mm diam, at first convex, then plane, mostly with a broad obtuse umbo; margin slightly-striate, thin, regular to wavy, sometimes sublobed; surface greasy, irregular (sometimes gibbous), smooth, not opaque, hygrophanous, with a somewhat silky appearance owing to a thin coating of white, micaceous pruina all over, which disappears with age, but persisting mainly around the umbo and at the pileus margin; reddish brown (6B8, 6C8, 7B8), darker in the centre, brown (7E7, 8D8), sometimes ochre yellow (4A5, 5A4) over the margin. **Lamellae** slightly spaced, adnate-emarginate, 2–3 mm broad, lamellulae I = (0) 1–3(–4), at first white, then more or less golden yellow (3A5, 4A6); edge slightly eroded and concolourous. **Stipe** 30–45 × 3–5 mm, cylindrical, but widened at the apex and pointed-rooting at the base, straight to sinuous, sometimes curved towards the base, firm, internally fistulose, olive yellowish (3B6, 3B7), white-pruinose overall, fibrillose-striate. **Context** fibrous, elastic, thin at pileus centre, purplish brown at the disc, ochre-pink in the stipe. **Odour** and **taste** strongly mealy. **Spore-print** not obtained (presumably white). **Spores** (5.0–)5.5–6.5(–7.0) × (3.5–)3.8–4.5(–4.8) µm (n = 40), av. 5.85 × 4.00 µm, Q = (1.33–)1.38–1.62(–1.67), Qm = 1.47, largely ellipsoid to ellipsoid, hyaline, slightly thick-walled, monoguttulate, cyanophilous, inamyloid, usually covered with large, obtuse, and irregular verrucae. The ornamented spores are mixed with apparently smooth spores. **Basidia** 25–28 × 7–8 µm, clavate, hyaline, siderophilous (with internal siderophilous/cyanophilous granules), 4-spored with sterigmata up to 4 µm long. **Hymenial cystidia** absent. **Hymenophoral trama** regular, consisting of parallel to slightly divergent, 4–5 µm wide hyphae. **Pileipellis** a slightly gelatinised cutis (ixocutis) composed of intertwined parallel up to 4 µm wide hyphae, with a clavate, up to 40 × 4 µm terminal element; pigment brown, parietal, cytoplasmic and encrusting. **Stipitipellis** similar to the pileipellis, of up to 6 µm wide, but non-encrusted hyphae. **Stipititrama** regular, consisting of parallel, up to 10 µm wide hyphae. **Thromboplerous hyphae** with golden yellow contents present. **Clamp connections** present everywhere, also at the base of the basidia.

Habitat & Distribution — Gregarious, in grassy areas. So far known only from the Dominican Republic.

Colour illustrations. Dominican Republic, Puerto Plata, Sosúa, El Batey, the garden where the holotype specimen was collected. *Calocybella goethei* basidiomata in field (holotype JBSD130968); spores and basidia (ammoniacal Congo red, holotype JBSD130968); pileipellis (Cotton blue, holotype JBSD130968). Colour codes in the macroscopic descriptions are from Kornerup & Wanscher (1983). Scale bars = 5 µm.

Typus. DOMINICAN REPUBLIC, Puerto Plata, Sosúa, Loc. El Batey, in the garden of a private school (House of Goethe), among the grass of the lawn, 19 Dec. 2020, C. Angelini (holotype JBSD130968, ITS, LSU and *rpb2* sequences GenBank MZ970330, MZ970356 and MZ995427, MycoBank MB 841772).

Notes — Among the lyophylloid genera, *Calocybella*, typified with *C. pudica* (= *C. juncicola* fide Corriol et al. 2017), encompasses species with collybioid habit, a cutis- to trichoderm-like pileipellis, smooth to verrucose spores and clamped hyphae (Vizzini et al. 2015, 2017, Latha et al. 2016). Its sister genus *Gerhardtia* differs mainly by clamp-less hyphae (Vizzini et al. 2015, 2017). *Calocybella goethei* is well differentiated from the other *Calocybella* species described in literature, based on morphological and molecular features. Its yellow lamellae and the co-presence of both smooth and warty mature spores are distinguishing, diagnostic features. The closely related *C. dominicana*, also described from the Dominican Republic, shows a pileus with orange tinges, white lamellae, stipe surface changing to red on bruising and all its spores are verrucose (Vizzini et al. 2017).

Calocybella pudica is presently known from Italy, France, Spain and India (*C. pudica* isolate KUBOT-KRMK, GenBank MW487244; see the phylogenetic tree), and shares with *C. goethei* the presence of dimorphic spores but differs by the entirely orange to brownish orange pileus without a darker umbo, yellow lamellae with orange tinges, yellow rhizoids at the stipe base, a context more or less rapidly changing to red on bruising or when exposed and narrower spores ((3.8–)4.9–6.0(–6.7) × (2.8–)3.0–3.7(–4.3) µm, Q = (1.3–)1.46–1.80(–2.1), Qm = 1.63; Vizzini et al. (2015)). The three new *Calocybella* species recently described from India (Latha et al. 2016) are morphologically quite different from our new species.

Supplementary material**FP1338** Phylogenetic tree.

Alfredo Vizzini, Department of Life Sciences and Systems Biology, University of Torino, Viale P.A. Mattioli 25, I-10125 Torino, Italy; e-mail: alfredo.vizzini@unito.it

Claudio Angelini, Herbario Jardín Botánico Nacional Dr. Rafael Ma. Moscoso, Santo Domingo, Dominican Republic, and

Via Cappuccini, 78/8 – 33170 Pordenone, Italy; e-mail: claudio_angelini@libero.it

Enrico Ercole, Via Murazzano 11, I-10141, Torino (TO), Italy; e-mail: ercole.enrico@gmail.com



Fungal Planet 1339 – 24 December 2021

***Calvatia nordestina* R.L. Oliveira, R.J. Ferreira, P. Marinho, M.P. Martín & Baseia, sp. nov.**

Etymology. In reference to the region where this species was collected, Nordeste, Brazil.

Classification — *Lycoperdaceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata growing gregarious, epigeous, rooting base with incrustations, globose to subglobose and 75–111 mm wide × 20–33 mm high. *Exoperidium* < 0.1 mm thin, fragile, slightly tomentose, evanescent, white (1A1, Kernerup & Wanscher 1978). *Mesoperidium* < 0.1 mm thin, fragile, papyraceous, persistent at the base, smooth with senescence, greyish brown to violet brown (7D3, 11F4). *Endoperidium* < 0.3 mm thin, fragile and brittle at the apex, resistant and persistent at the base, tomentose, greyish brown to reddish brown (9D3, 9D4). *Rhizomorphs* 1–1.5 mm wide × 10–22 mm long, branching, white (1A1) with sand incrustations. *Subgleba* reduced, woolly, compact, dark blond to reddish brown (5D4, 9E4). *Gleba* powdery, not persistent, greyish brown to violet brown (9D3, 11F4), at maturity. *Exoperidium* hyphallic, 2.9–5.1 µm diam, frequent and non-regular septa, double V branching, regular walls ≤ 0.8 µm thin, straight for curves, hyaline, not dextrinoid and acyanophilic. *Mesoperidium* pseudoparenchymatous composed of cells measuring 11.4–39.3 µm wide × 15.4–46.8 µm high, regular walls ≤ 1.1 µm thin, hyaline, not dextrinoid, and acyanophilic. *Apical endoperidium* composed of two layers of hyphae throughout, all brown, dextrinoid and cyanophilic, hyphae 2.3–5.8 µm diam, regular walls ≤ 0.8 µm thickness and curves, frequent and non-regular true septa, double V branching, and mycosclereids globose, subglobose, pyriform, ovoid, ellipsoid, or rectangular, 10.1–23.1 µm high × 4–13.5 µm diam, weakly interconnected, branched, breaking in the septa, regular and walls ≤ 1 µm thickness. Basal *endoperidium* hyphallic, 2.2–4.3 µm diam, rare and non-regular true septa, V-shaped branches, single and double, and in T, regular walls < 0.9 µm thickness, curves, hyaline, dextrinoid, and cyanophilic. *Rhizomorphs* hyphallic, 2.6–5.9 µm diam, regular walls ≤ 0.9 µm thickness and curves, hyaline, not dextrinoid and cyanophilic. Hyphallic *subgleba*, 1.6–4.9 µm diam, rare true septa, branching V, single and double, and T, regular walls ≤ 0.7 µm thickness, curves, pale brown, dextrinoid and acyanophilic. *Paracapillitium* absent. *Capillitium* *Calvatia*-type, 2–3.9 µm diam, pale brown, dextrinoid and cyanophilic; septa frequent and non-regular, V-branching, single and double, and in T, fragmenting in any part of the capillitium or frequent in the septa; regular walls ≤ 0.9 µm thickness, straight, with small, irregular, numerous and conspicuous pits. *Basidiospores* globose, 4.2–6.1 µm ×

4.1–6.1 µm (av. = $5.3 \pm 0.6 \times 5.2 \pm 0.7$; Qm = 1.02; n = 20), equinulated, ornamentation < 0.7 µm length; pedicels present in some basidiospores ≤ 1.4 µm in length.

Habit & Habitat — Basidiomata gregarious, growing on moist soil.

Typus. BRAZIL, Rio Grande do Norte, João Câmara, Serra do Torreão, 17 Feb. 2017, R.L. Oliveira (holotype UFRN-Fungos 3037, ITS sequence GenBank MW830151, MycoBank MB 839179).

Additional material examined. BRAZIL, Rio Grande do Norte, João Câmara, Serra do Torreão, 4 July 2017, R.L. Oliveira (UFRN-Fungos 3038); *ibid.*, 1 May 2020, R.L. Oliveira (UFRN-Fungos 3327); João Câmara, Community of Matão, close to BR 406, 1 May 2020, R.L. Oliveira (UFRN-Fungos 3328, UFRN-Fungos 3329).

Notes — *Calvatia nordestina* is morphologically related to species of the Sect. *Hippoperdon*. Based on morphological and molecular characters, it is close to some other *Calvatia* species, such as *C. cyathiformis*, *C. fragilis*, *C. caatinguensis* and *C. brasiliensis*. *Calvatia cyathiformis* and *C. caatinguensis* have a well-developed subgleba (Zeller & Smith 1964, Crous et al. 2018a), characteristics not found in *C. nordestina*. *Calvatia fragilis* is another morphologically similar species to *C. nordestina*, mainly by an extremely powdery and violacea gleba and reduced subgleba (Morgan 1890); however, *C. fragilis* has larger basidiospores, 6.4–7.2 µm diam (Bates et al. 2009). *Calvatia brasiliensis* has similar macroscopic morphological characteristics to *C. nordestina*, but *C. brasiliensis* differs in the basidiospore ornamentation and capillitium type (Crous et al. 2019a). *Calvatia lilacina* resembles *C. nordestina* by the colour of the basidiomata and basidiospores, but they can be easily distinguished based on other morphological characters. Thus, *Calvatia lilacina* has conical to turbiniform basidiomes with a well-developed, rigid cellular subgleba, easily distinguishable from the fertile portion by the presence of cavities (Berkeley 1845). Such morphological characters are not seen in specimens of *C. nordestina*, which has globose to subglobose basidiomata, reduced, compact and woolly subgleba. Moreover, the ITS nrDNA sequences among *C. lilacina* and *C. nordestina* shown a 1.5 % of different bases to confirm that they are two different species. Clade I represent *C. lilacina* s.str., accommodating sequences from Pakistan, Argentina and Australia (Locality of type). Clade II encompasses specimen sequences from Mexico and the USA, but it is not possible to infer its phylogenetic position in relation to *C. nordestina* or *C. lilacina* since the morphology of these specimens was not analysed.

Colour illustrations. Brazil, Rio Grande do Norte, João Câmara, Serra do Torreão, where the specimens were collected. From top to bottom: basidiospore under SEM; capillitium under SEM; longitudinal section through mature basidioma; mature basidioma *in situ*. Scale bars = 10 mm (basidiomata), 1 µm (SEM photos).

Supplementary material

FP1339 Phylogenetic tree.

Renan de L. Oliveira, Programa de Pós-Graduação em Sistemática e Evolução, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Av. Senador Salgado Filho, 3000, 59072-970 Natal, RN, Brazil; e-mail: brazil_renan77@yahoo.com.br

Renato J. Ferreira, Programa de Pós-Graduação em Biologia de Fungos, Departamento de Micologia, Universidade Federal de Pernambuco, 50670-420 Recife, PE, Brazil; e-mail: renatojuciano@hotmail.com

Paulo Sérgio Marinho Lúcio, Departamento de Biologia Celular e Genética, Universidade Federal do Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil; e-mail: paulomarinho@hotmail.com

María P. Martín, Real Jardín Botánico RJB-CSIC, Plaza de Murillo 2, 28014 Madrid, Spain; e-mail: maripaz@rjb.csic.es

Iuri G. Baseia, Departamento Botânica e Zoologia, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Campus Universitário, 59072-970 Natal, RN, Brazil; e-mail: iuri.baseia@gmail.com

Cladosporium austrolitorale



Fungal Planet 1340 – 24 December 2021

***Cladosporium austrolitorale* C. Gil-Durán & L. Sanhueza, sp. nov.**

Etymology. The epithet *austrolitorale* means 'from the southern seashore' and refers to the coast of Fildes Bay, Antarctica, where the fungus was isolated.

Classification — *Cladosporiaceae*, *Cladosporiales*, *Dothideomycetes*.

Mycelium superficial and immersed, composed of septate, branched, subhyaline to olive brown, smooth hyphae, 2.4–4.1 µm wide. **Conidiophores** dimorphic, macronematous conidiophores arising from lateral or terminal hyphae, erect to slightly flexuous, septate, branched, or unbranched, 20–53.9 µm long, 2.8–3.8 µm wide, pale brown, smooth; micronematous conidiophores arising terminally or laterally from hyphae, 2.3–3.1 µm wide, pale olivaceous brown, with a single apical scar. **Ramoconidia** straight, verrucose, pale brown or pale olivaceous, ellipsoid to subcylindrical, 8.2–22.1 × 3.6–4.6 µm, 0–1-septate. **Secondary ramoconidia** ellipsoid to subcylindrical, verrucose, 8.9–14.2 × 4.9–5.2 µm, 0–1-septate in the middle, with 2–3 distal hila, proliferating sympodially. **Conidia** numerous, catenate, dichotomously branched in all directions, straight, verrucose, with up to five conidia; small terminal conidia subglobose or obovoid, 4.3–5.6 × 3.4–4.6 µm; intercalary conidia obovoid or limoniform, 7.1–11.0 × 3.3–5.3 µm; microcyclic conidiogenesis not observed.

Culture characteristics — (after 2 wk at 20 °C in the dark): On potato dextrose agar (PDA) colonies reach 40–43 mm diam, concentric wavy ring-like pattern with radial grooves, flat colony centre, submerged, profuse sporulation, exudates absent, margin white and undulate; reverse, mycelium immersed in the agar, olivaceous green to olivaceous black. On malt extract agar (MEA), colonies reach 36–39 mm diam, irregular flat growth, elevated centre, opaque dusty, olivaceous green to yellowish green, aerial mycelium absent, exudates absent, white filiform margin; reverse irregular olivaceous black colour. On synthetic nutrient-poor agar (SNA), colonies reach 29–32 mm diam, flat growth, round centre, radiating filaments from the centre, dusty olivaceous green, profuse sporulation mainly in the centre of the colony, exudates absent, filiform margin; reverse olivaceous with white feathery margin. On oatmeal agar (OA) colonies reach 36–40 mm diam, round shape, flat, wrinkled, olivaceous green, abundant aerial mycelium, profuse sporulation, exudates absent, undulate margin.

Cardinal temperature for growth — Optimum 20 °C, maximum 25 °C, minimum 5 °C.

Typus. ANTARCTICA, South Shetland archipelago, King George Island, Fildes Bay, from coastal sea sand, 24 Feb. 2018, L. Sanhueza, LS-1 (holotype CHFC-EA 593 stored in a metabolically inactive state in the Chilean Fungal Collection, culture ex-type CBS 148321, ITS, LSU, *actA* and *tef1* sequences GenBank MN879327, MT003768, MN890007 and MN890010, MycoBank MB 840658).

Colour illustrations. Landscape of the coast of Fildes Bay, Antarctica. *Cladosporium austrolitorale* growing on PDA and SNA; conidiophore, conidia and ramoconidium on SNA after 14 d at 20 °C. Scale bars = 10 µm (conidiophore), 1 µm (conidia and ramoconidium).

Notes — Based on the combined analysis of ITS, *actA* and *tef1* markers, *C. austrolitorale* belongs to the *C. cladosporioides* complex (Bensch et al. 2015) and is phylogenetically related to *C. arenosum*. However, the surface of conidia and ramoconidia of *C. austrolitorale* exhibits verrucose ornamentation, while these structures are smooth in *C. arenosum* (Crous et al. 2020). In addition, colonies of *Cladosporium austrolitorale* have a characteristic concentric wavy ring-like pattern after 2 wk at 20 °C on PDA, and radiating filaments from the colony centre after 2 wk at 20 °C on SNA, characters not found in *C. arenosum* (Crous et al. 2020).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence are *C. perangustum* (ID58, GenBank MN511354.1; Identities 551/551 (100 %), no gaps), *C. arenosum* (CHFC-EA 566, GenBank MN879328.1; Identities 551/551 (100 %), no gaps), *C. globisporum* (DTO 220-D4, GenBank KP701967.1; Identities 551/551 (100 %), no gaps) and *C. asperulatum* (UTHSC DI-13-21, GenBank LN834357.1; Identities 551/551 (100 %), no gaps). The closest hits using the LSU sequence are *C. tenuissimum* (CBS 125995, GenBank NG_069983.1; Identities 608/608 (100 %), no gaps), *C. scabrellum* (CBS 126358, GenBank NG_069960.1; Identities 608/608 (100 %), no gaps), *C. phyllactiniicola* (CBS 126355, GenBank NG_069959.1; Identities 608/608 (100 %), no gaps), and *C. myrtacearum* (CBS 126350, GenBank NG_069958.1; Identities 608/608 (100 %), no gaps). The closest hits using the *actA* sequences were *C. asperulatum* (BP8I3, GenBank KU605799.1; Identities 222/224 (99 %), no gaps), *C. myrtacearum* (CBS 126349, GenBank HM148605.1; Identities 212/227 (93 %), four gaps (1 %)), *C. colombiae* (CBS 274.80B, GenBank FJ936166.1; Identities 215/233 (92 %), eight gaps (3 %)), and *C. lycoperdinum* (CBS 126348, GenBank HM148602.1; Identities 214/232 (92 %), eight gaps (3 %)). The closest hits with *tef1* sequences were *C. arenosum* (CHFC-EA 566, GenBank MN890011.1; Identities 236/243 (97 %), no gaps), *C. asperulatum* (BP3I2, GenBank KU605784.1; Identities 235/242 (97 %), no gaps), *C. angustiterminale* (CPC 15564, GenBank KT600476.1; Identities 215/243 (88 %), five gaps (2 %)), and *C. funiculosum* (UTHSC DI-13-242, GenBank LN834460.1; Identities 194/247 (79 %), seven gaps (2 %)).

Supplementary material**FP1340** Phylogenetic tree.



Fungal Planet 1341 – 24 December 2021

***Coccomyces pycnophyllocladi* P.R. Johnst., sp. nov.**

Etymology. Refers to the similarity to *Coccomyces phyllocladi*, distinguished macroscopically by the presence of asexual morph.

Classification — *Rhytismataceae*, *Rhytismales*, *Leotiomyces*.

Ascomata and pycnidial conidiomata forming within discrete paler areas on fallen leaves of *Phyllocladus alpinus*, these areas often have a diffuse, dark brown margin. *Ascomata* 0.4–0.7 mm, angular, 3–4-sided, wall black, pale zone along future line of opening, lacking lip cells along the radiate opening slits. *Paraphyses* 2.5–3 µm diam, swelling to 4.5–5 µm at the clavate apex, embedded in thick, hyaline gel, extending about 15 µm beyond asci. *Asci* 110–120 × 9–10 µm, cylindrical, tapering gradually to broad subtruncate apex, wall undifferentiated at apex, 8-spored, spores extending almost to base, asci with crozier at base. *Ascospores* 70–80 × 2.5–3 µm unreleased in asci, tapering gradually and slightly towards base, aseptate, hyaline, flattened apical gelatinous cap, small basal gelatinous cap seems to be present but not clearly seen. *Conidiomata* 0.1–0.15 mm, round, dark brown with narrow black margin.

Typus. NEW ZEALAND, Fiordland, Fiordland National Park, Kepler Track, vic. control gates, S45.4648 E167.6740, on dead leaves of *Phyllocladus alpinus* (*Podocarpaceae*), 10 May 2000, P.R. Johnston et al., R1006 (holotype PDD 72098, culture ex-type ICMP 17376, ITS, LSU and mtSSU sequences GenBank MZ702495, MZ702535 and MZ702531, MycoBank MB 841002).

Additional specimen examined. NEW ZEALAND, Buller, Paparoa Range, Croesus Track, on dead leaves *Phyllocladus alpinus*, 3 May 1985, P.R. Johnston (R656), G.J. Samuels & L.M. Kohn (PDD 46953).


Notes — *Coccomyces pycnophyllocladi* differs from the more commonly collected *C. phyllocladi* by the slightly wider asci and ascospores, asci being broadly truncate at the apex, and in having pycnidial conidiomata. Both *C. pycnophyllocladi* and *C. phyllocladi* can be present on the same leaves, the two species distinguishable by the presence or absence of pycnidia within the pale lesions. *Coccomyces clavatus*, also found on *Phyllocladus*, has shorter, clavate ascospores. The flattened apical gelatinous cap of *C. pycnophyllocladi* matches that found in *C. clavatus* (Johnston 1994). Phylogenetically, the

three *Phyllocladus*-inhabiting species known from New Zealand are closely related; *C. phyllocladi* occurs also in Australia (e.g., PDD 68250, GenBank MH578535). The three species have a 96.9–97.5 % match across the ITS region. *Coccomyces phyllocladi* and *C. pycnophyllocladi* have a 96.8 % match across the part of the LSU region amplified with the LROR and LR5 primers (excluding a 218 intron in *C. phyllocladi*, sequences GenBank MZ960460, MZ960459, HM140508).

Both known collections of *C. pycnophyllocladi* are small, and to preserve material no attempt was made to section the apothecia. Based on other observations, the ascomatal structure is likely similar to that described for *C. clavatus* and *C. phyllocladi* (Johnston 1986).

Maximum-likelihood phylogeny based on ITS sequences. Although few nodes are well supported in this phylogeny, it illustrates the phylogenetic distinctness of the new species *Coccomyces pycnophyllocladi*, *Hypoderma subiculatum*, and *Hypoderma aliforme*. Note that generic limits within *Rhytismales* remain morphologically based and most genera as currently accepted are polyphyletic. Phylogeny generated using IQ-TREE v. 1.6.6 (Nguyen et al. 2015; Chernomor et al. 2016), model TIM2e+I+G4 selected using ModelFinder (Kalyaanamoorthy et al. 2017), and bootstrap support values from 1000 ultrafast bootstrap replicates are shown where > 95 % (Hoang et al. 2018). *Marthamyces emarginata* was used as an outgroup.

Colour illustrations. New Zealand, Fiordland, high altitude *Nothofagaceae* forest with *Phyllocladus alpinus*. Ascomata and pycnidia; asci and ascospores; paraphyses; detail of flattened gel cap at ascospore apex (arrowed). Scale bars = 1 mm (ascomata, PDD 72098), 10 µm (all others; asci and paraphyses PDD 72098, gel cap PDD 46953).





Fungal Planet 1342 – 24 December 2021

***Hypoderma aliforme* P.R. Johnst., sp. nov.**

Etymology. From *aliformis*, wing-shaped; reflecting ascospore shape, recalling an aircraft wing in cross section.

Classification — *Rhytismataceae*, *Rhytismatales*, *Leotiomyces*.

Ascomata and pycnidial conidiomata developing on fallen leaves of *Fuscopora solandri* and *F. fusca* within pale areas on host leaf, the pale areas often surrounded by a more or less complete, narrow, black zone line. *Ascomata* 1–1.5 × 0.5–0.7 mm, broad elliptical in outline with broadly rounded ends, wall black, sometimes paler just inside the two ends, single longitudinal opening slit lined with pale grey lip cells. In vertical section ascomata subcuticular, upper wall up to 70 µm thick near edge of opening, gradually thinner towards base of wall, comprising angular cells 5–7 µm diam, with walls thick, dark brown. Exposed face of upper wall along opening slit lined with unbranched cylindrical cells, 25–30 × 2.5–3 µm with walls thin, hyaline. Lower wall of ascomata 5–10 µm thick, comprising 2–3 layers of angular cells with walls thick, brown. *Paraphyses* 1–2 µm diam, circinate at apex, extending 5–10 µm beyond asci. *Asci* 110–140 × 9.5–11(–12.5) µm, clavate-stipitate, tapering to broadly truncate apex, wall undifferentiated at apex, 8-spored, crozier at base. *Ascospores* 18–21 × 3.5–4.5 µm, fusoid-clavate, widest point in upper half, tapering to narrow, rounded to subacute ends, aseptate, hyaline, surrounded by a narrow gelatinous sheath. *Conidiomata* 0.2–0.3 mm diam, more or less round, black-walled. Conidiomata subcuticular, upper wall 5–10 µm thick, of dark tissue with no obvious cellular structure, lower wall 10–15 µm thick of 3–4 layers of angular cells with dark, thick walls, lined with 2–3 layers of cells with hyaline, thin walls on which the conidiogenous cells are formed. *Conidiogenous cells* 8–15 × 1.5–2 µm, solitary, cylindrical, tapering to apex, proliferating sympodially or occasionally percurrently. *Conidia* 3–4 × 1 µm, cylindrical, often slightly sigmoid, ends acute, hyaline, aseptate.

Typus. NEW ZEALAND, Mid Canterbury, vic. Springfield, Kowai Bush, S43.2897 E171.9259, on fallen leaves *Fuscopora solandri* (*Nothofagaceae*), 18 May 2000, P.R. Johnston (R1011) & R.E. Beever (holotype PDD 72102, culture ex-type ICMP 17379, ITS, LSU and mtSSU sequences GenBank MZ702499, MZ702536 and MZ702532, MycoBank MB 841000).

Additional materials examined. NEW ZEALAND, Mid Canterbury, vic. Springfield, Kowai Bush, on fallen leaves *Fuscopora solandri*, 5 May 1995, P.R. Johnston (R938) (PDD 64922, ICMP 17362, ITS, LSU and mtSSU sequences GenBank MZ702500, MZ702538 and MZ702534); North Canterbury, Arthurs Pass, Old Coach Rd, on fallen leaves *Fuscopora solandri*, 5 May 1989, P.R. Johnston (R822), P.K. Buchanan & M. Rajchenberg (PDD 55525). Fiordland, Fiordland National Park, Lake Hauroko, on fallen leaves *Fuscopora solandri*, 8 May 1984, P.R. Johnston (R462) (PDD 46211); Southland, vic. Gore, Piano Flat, Creek Loop Track, on fallen leaves *Fuscopora solandri*, 21 May 2008, P.R. Johnston (R1050.1), R.E. Beever & B.C. Paulus (PDD 94910).




Colour illustrations. New Zealand, Mid Canterbury, Kowai Bush, type locality (photo Jon Sullivan). Infected leaf; detail of ascomata; ascoma in vertical section; ascospores; apex of asci. Scale bars = 1 mm (infected leaf, PDD 72102), 0.5 mm (ascomata, PDD 72102), 100 µm (vertical section, PDD 72102), 10 µm (all others; ascospores and asci, PDD 72102).

Notes — *Hypoderma aliforme* specimens were previously incorrectly identified as *Hypoderma rubi* (Johnston 1990). DNA sequences showed this *Nothofagaceae*-inhabiting fungus to be phylogenetically distinct from *H. rubi*, and this correlates with a consistent difference in ascospore shape. The ascospores of *H. rubi* are more or less naviculate, widest near the centre, taper slightly to rounded apex and more markedly to the narrowly rounded base, whereas *H. aliforme* has fusoid-clavate, radially symmetrical ascospores, tapering to both narrow rounded ends. *Hypoderma rubi* has not been found on *Nothofagaceae* in New Zealand. *Hypoderma sigmoideum* and *Hypoderma obectum* are other *Hypoderma* spp. associated with *Fuscopora*, *H. sigmoideum* especially often being found on the same leaves as *H. aliforme*. *Hypoderma sigmoideum* differs from *H. aliforme* in having red lip cells and longer, sigmoid ascospores; *H. obectum* has ascomata surrounded by subiculum-like hyphae and has longer, bifusiform ascospores.

See the phylogenetic tree provided in Fungal Planet 1341 elsewhere in this paper.



Hypoderma subiculatum



Fungal Planet 1343 – 24 December 2021

***Hypoderma subiculatum* P.R. Johnst., sp. nov.**

Etymology. Refers to the subiculum-like hyphae surrounding the ascomata.

Classification — *Rhytismataceae*, *Rhytismatales*, *Leotiomyces*.

Ascomata 1.5–4 × 0.4–0.6 mm, elliptical, ends acute, wall black, pale creamy lip cells along the single, longitudinal opening slit, groups of ascomata often surrounded by woolly, white subiculum, subiculum hyphae encrusted with clumps of white crystals. In vertical section ascomata subcuticular, upper wall up to 165 µm thick near edge of opening, narrowing towards base of wall where it is about 30 µm thick, comprising angular cells 6–8 µm diam, with walls irregularly encrusted with dark brown material. Exposed face of upper wall along opening slit lined with unbranched cylindrical cells 9–20 × 2.5–4 µm with walls thin, hyaline. Lower wall of ascomata 20–35 µm thick, comprising several layers of somewhat tangled hyphae with walls irregularly encrusted with dark brown material. Subhymenium 20–25 µm thick, comprising hyaline *textura intricata*. **Paraphyses** 1 µm diam, slightly and irregularly swollen and circinate near apex, occasionally branched. **Asci** (100–)115–125 × 14–16 µm clavate, long stipitate, tapering gradually to rounded apex, wall undifferentiated at apex, 8-spored, crozier at base. **Ascospores** 22–27(–30) × 3–4 µm, cylindrical, straight, tapering slightly towards base, ends rounded, aseptate, hyaline, surrounded by a 1 µm thick gelatinous sheath. **Conidiomata** 0.5–1 × 0.25 mm, cylindrical with rounded ends, wall pale grey to grey. **Conidiomata** with poorly developed upper wall, lower wall comprising 3–4 layers of cells 4–5 µm diam, angular, walls thick and dark, lined with 1–2 layers of thin walled, pale brown cells on which the conidiogenous cells are formed. **Conidiogenous cells** 10–15 × 1.5–2.5 µm, cylindrical, taper towards apex, proliferating percurrently and sympodially. **Conidia** 4–4.8 × 1.5 µm, cylindrical, straight, apex rounded, base truncate, aseptate, wall hyaline.

Typus. NEW ZEALAND, Westland, Haast, Ship Creek, beach loop track, S43.7599 E169.1440, on dead leaves *Phormium tenax* (*Asphodelaceae*), 18 May 2018, P.R. Johnston (D2528) & M. Padamsee (holotype PDD 117664; ITS sequence GenBank MZ702490, MycoBank MB 841001).

Additional materials examined. NEW ZEALAND, Stewart Island, Doughboy Bay, near cave, S47.0332 E167.7072, on dead leaves *Phormium* sp., 29 Apr. 2002, P.R. Johnston (R1023), R. Leschen & S.R. Whitton (PDD 78367, culture ICMP 17397, ITS, LSU and mtSSU sequences GenBank MZ702491, MZ702537 and MZ702533); Dunedin, Swampy Spur Rd, S45.8045 E170.4827, on dead leaves *Phormium* sp., 16 May 2008, P.R. Johnston (R1042), R.E. Beever & B.C. Paulus (PDD 95046).

Colour illustrations. New Zealand, Westland, Ship Creek, beach near mouth of creek, *Phormium* along foreshore. Ascomata; ascoma in vertical section; detail of ascomatal opening in vertical section; asci; ascospores; conidiogenous cells. Scale bars = 1 mm (ascomata, PDD 117664), 100 µm (vertical section, PDD 78367), 10 µm (all others; vertical section of ascomatal opening, PDD 78367, all other images, PDD 117664).

Notes — To date *Hypoderma subiculatum* has been found only in the south of New Zealand. The distinctive white subiculum is sometimes mostly worn off and in this case a specimen may be confused with two other *Hypoderma* spp. also found on *Phormium*, *H. cordylines* and *H. cookianum*. These two species both have macroscopically similar ascomata, although they never have subiculum hyphae. *Hypoderma cookianum* lacks lip cells along the opening slit and has large, bifusiform ascospores; *H. cordylines* has reddish coloured lip cells and shorter, broader ascospores (14–21 × 4.5–6 µm, Johnston 1990). All three species are phylogenetically distinct.

See the phylogenetic tree provided in Fungal Planet 1341 elsewhere in this paper.



Fungal Planet 1344 – 24 December 2021

***Crepidotus innuopurpureus* McMull.-Fish., T. Lebel, Senn-Irlet, sp. nov.**

Etymology. Named for the brightly coloured pigments (magenta to violet) on pileus and lamellae, particularly margins of the lamellae, that are lost with exposure to the elements, so often only 'hints of purple' remain, *innuo* (L. for hint) and *purpura* (L. for purple).

Classification — *Crepidotaceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata small, kidney-shaped, younger specimen hoof-shaped. **Pileus** 3–10 mm (radius), convex to near plane, outer surface matt, clay pink, mid-pink, grey red, to lilac, becoming ochraceous to brown with age or exposure, finely tomentose, and some basidiomata radially wrinkled, outer rim paler to concolourous, margin inrolled in younger specimens, to undulating in some older specimens. **Lamellae** close to subdistant, conspicuously magenta, clay pink, mid-pink to mid-red to brown-red (8B5-8C5, Kornerup & Wanscher 1978) with strongest colour concentrated at margins. Lamellulae present, in two tiers. **Attachment** astipitate, sessile with lateral, eccentric to dorsally attached basal pad of white to pale tomentum. **Spores** 5.5–7.1(–8) × 5–6.3 µm (av. 6.8–5.8 µm), Q = 1–1.4 (av. = 1.14, n = 73), globose to subglobose, brown, distinctly but finely warty, ornamentation 0.5–0.8 µm high. **Basidia** four-spored. **Cheilocystidia** variable in shape, from lageniform to cylindrical subcapitate or clavate to broadly utriform, sometimes branched, often coloured and often with external crystals. **Pileipellis** a thin cutis with diverticulate hyphae, containing dark red to violet pigment, pigment is concentrated in scattered patches. **Pigment deposits** – inside elements coloured, some with crystals present on outside of cheilocystidia, and patchy in concentration, densest at lamellae edges and on the pileipellis, pigment colour dissipates in weak (3 %) potassium hydroxide. **Clamp connections** present at base of basidia and cheilocystidia, and on pileipellis, trama and basal pad hyphae.

Habit, Habitat & Distribution — Small, gregarious, fan-shaped mushrooms found on bark of living rough barked rainforest trees and stags or logs, sometimes without bark; often amongst bryophytes. Currently known from rainforest, with subtropical rainforest at Mary Cairncross Scenic Reserve near Maleny, Blackall Range, Queensland and in tropical rainforest below sandstone cliff at in the Jameson Arch, near Mt Agnes, West Kimberley, Western Australia. Expected to have a tropical and subtropical distribution across northern Australia.

Colour illustrations. Remnant subtropical rainforest, Mary Cairncross Scenic Reserve, Maleny, Queensland, Australia, holotype site. Basidiomata (photo credit M.D. Barrett); SEM mature spore (photo credit B. Senn-Irlet); dried specimen with arrows highlighting pigment; microscopic elements: basidiospores, pigmented hyphae in pileipellis, basidia, cheilocystidia variable shapes, some with crystals (drawing credit B. Senn-Irlet & K. Syme). Scale bars = 5 µm, except for dried specimen 5 mm.

Typus. AUSTRALIA, Queensland, Blackall Range, Mary Cairncross Scenic Reserve, subtropical rainforest, on dead wood of leafy trees, 27 Feb. 1996, T.R. Lohmeyer (holotype MEL 2503290; ITS and LSU sequences MZ870345 and MZ870347, MycoBank MB 840921).

Additional material examined. AUSTRALIA, Western Australia, West Kimberley Region, Jameson Arch, tropical rainforest, 28 Jan. 2007, M.D. Barrett, MDB F82/07 (ITS and LSU sequences MZ870344 and MZ870346).

Notes — Other pink, peach, lilac and purple coloured *Crepidotus* occur across Australia but *C. innuopurpureus* is distinctive microscopically with the strong pigments found within pileal elements and cheilocystidia, and coloured crystals observed on the outside tip of some cheilocystidia. These patches of pigment may survive drying and are most obvious on lamellae edges and on the surface of the cap but fade with age and likely exposure to light and other environmental elements. The bright magenta pigment patches are often still obvious in dried specimens; the pigment dissolves in seconds if examined under KOH. The overall colour of the basidiomes is a pale pink brown to lilac brown, that becomes a pale dull brown colour with any exposure to light.

Crepidotus innuopurpureus is part of a small but strongly supported clade, including *C. parietalis*, which is currently known from Australia, India and New Zealand. *Crepidotus parietalis* has small yellowish basidiomata, that similarly has crystals on the outside tips of the cheilocystidia, warty spores, and loses pigment in KOH (Horak 1978, 2018). The Indian collection (GenBank MK567976) is labelled as '*C. roseus*' (Manoj et al. 2018). However, none of the descriptions of *Crepidotus roseus* mention crystals or pigmentation in the pileal elements or cheilocystidia (Singer 1947, Guzmán-Dávalos et al. 2017). Further collections and analysis of Indian collections are needed to determine relationships with our new species. Examination and analysis of more tropical and subtropical Australasian collections will likely provide further taxa in this intriguing group.

Supplementary material**FP1344** Phylogenetic tree.

Cuphophyllus bondii



Fungal Planet 1345 – 24 December 2021

***Cuphophyllus bondii* Lebeuf & I. Saar, sp. nov.**

Etymology. Name honours Sir Robert Bond, last premier of the British Colony of Newfoundland and first Prime Minister of the Dominion of Newfoundland. He solved the 'French shore' problem, allowing permanent settlement in peace, and he bequeathed his property to the people of Newfoundland and Labrador, which became Sir Robert Bond Park, type location of *Cuphophyllus bondii*.

Classification — *Hygrophoraceae*, *Agaricales*, *Agaricomycetes*.

Pileus 12–35 mm diam, at first convex, becoming applanate or slightly depressed at the disk with a low umbo, the margin incurved at first, becoming straight, reflexed, wavy, then completely upturned with age; surface smooth, lubricous, hygrophanous, orange-grey, greyish orange, pinkish white, or between orange-white and pale orange (6B2, 6B3, 7A2, 5½A2–3 in Kernerup & Wanscher 1978). **Lamellae** decurrent, arcuate at first, distant, increasingly intervenose with age, 2–4 mm wide, with 1–2 tiers of lamellulae, whitish to very pale orange, yellowing in exsiccata. **Stipe** 36–64 × 3.5–10 mm, equal, straight or curved, smooth, dry, with age fibrillose-striate and developing small scales in the upper third, stuffed then hollow, whitish or concolourous with the lamellae in the upper half, with increasing yellow tones towards the base. **Context** thin, thick at disk; odour and taste not recorded. **Spore print** white. **Basidiospores** (134 spores, four basidiomata, two collections) 6–8 × 4.5–6 (–6.5) µm, av. 6.9 × 5.4 µm, Q = 1.17–1.51, Q_{av} = 1.29, broadly ellipsoid to ellipsoid, subamygdaliform, inamyloid, smooth. **Basidia** 40–60 × 6.5–9 µm, 4-spored, more rarely 2- or 3-spored, clavate; sterigmata 3–8 µm long. **Subhymenium** ramose, not gelatinised, made of short 2–4 µm wide cells. **Hymenial cystidia** absent. **Lamellar trama** irregular-interwoven, made up of cylindrical and inflated hyphae 25–85 × 4–12 µm. **Pileipellis** an ixocutis 50–100 µm wide, made of gelatinised, spirally-incrusted or smooth hyphae 1–5 µm diam. **Context** made of tightly interwoven inflated, smooth hyphae 15–80 × 4–14 µm diam. **Stipitipellis** a cutis made of 3–5 µm wide smooth hyphae. **Clamp connections** present in all tissues, including occasional medallion clamps in pileipellis.

Habitat & Distribution — Solitary or gregarious in grassland, in the fall. Currently only known from the type location, on the Avalon Peninsula.

Typus. CANADA, Newfoundland and Labrador, Whitbourne, Sir Robert Bond Park, N47°24'42.0" W53°32'15.8", 56 m a.s.l., in grassland with some *Alnus* at a distance, 4 Oct. 2020, D. Spencer (holotype DAOM 984902, isotype TUF117862, ITS sequence GenBank MZ681950, MycoBank MB 839736).


Additional materials examined. CANADA, Newfoundland and Labrador, Whitbourne, Sir Robert Bond Park, N47°24'44.2" W53°32'15.7", 55 m a.s.l., in grass under *Quercus* and *Acer*, 25 Sept. 2018, R. Lebeuf, Foray NLA18A-034 (DAOM 981262, TUF117704, ITS sequence GenBank MZ681949).

Notes — *Cuphophyllus bondii* is characterised by the pale orange to pinkish tones of the pileus, its pale lamellae, yellow stipe base, mostly broadly ellipsoid to ellipsoid spores measuring 6–8 × 4.5–6.5 µm and growth in grass in the fall. A preliminary description of the paratype was published by Voitek et al. (2020a). A few other North American species of *Cuphophyllus*

Colour illustrations. Collection site of the paratype of *Cuphophyllus bondii*, in grass (photo credit Roger Smith). Lower inserts show holotype in situ. Note the presence of *Pilosella* sp., often observed in the vicinity of hygrocyboid fungi; basidia in Congo Red; spores in Congo Red (above) and KOH 3% (below). Scale bars = 10 mm (basidiomata), 10 µm (basidia and spores).


display brownish or orangish pileal colour: *C. lamarum*, recently described from Newfoundland and Labrador (Voitek et al. 2020b), also shows yellow tones at the base of the stipe, but it grows in sphagnum bogs in the summer (July); *C. hygrocyboides*, likely circumboreal, described from the French Alps and recently reported (Voitek et al. 2020b) from the alpine zone in British Columbia, Canada, has a darker bright orange-brown pileus and a pale orange-brown stipe lacking yellow tones at the base. *C. colemannianus* has a darker reddish brown, orange-brown to dark brown pileus and a white stipe without yellow tint.

Based on a blastn search of UNITE database, the closest hits of ITS sequence from the isotype are *Cuphophyllus flavipes* (GenBank MK547066; Identities = 679/705 (96 %), 18 gaps (3 %)), *C. flavipesoides* (GenBank MK573931; Identities = 681/708 (96 %), 19 gaps (3 %)), and *C. pseudopallidus* (GenBank HQ185706; Identities = 434/454 (96 %), 13 gaps (3 %)). Bayesian inference analysis (BI) was performed with MrBayes v. 3.2.6 (Ronquist et al. 2012) with 5 million generations, applying default values of other prior settings. The first 500 K generations without a stable likelihood score were discarded. Maximum likelihood (ML) analysis was performed with RAXML-HPG BlackBox v. 8.2.9 (Stamatakis 2014), at the CIPRES Science Gateway (Miller et al 2010; <http://www.phylo.org/>). The best tree of the BI analysis is presented.



Novel species is highlighted in a light green box. The scale bar indicates the expected number of substitutions per site, and bootstrap support values of the ML analyses ($\geq 70\%$) and Bayesian Posterior Probabilities (≥ 0.95) are above and below branches, respectively. The tree was rooted to the sequences of *C. cinerellus*, *C. esteriae* and *C. lamarum*. The alignment and tree were deposited in TreeBASE (study 28644, <http://purl.org/phylo/treebase/phyloids/study/TB2:S28644>).

Curvularia stenotaphri



Fungal Planet 1346 – 24 December 2021

***Curvularia stenotaphri* N.T. Tran, Geering, Y.P. Tan & R.G. Shivas, sp. nov.**

Etymology. Name refers to the plant genus, *Stenotaphrum*, with which the fungus is associated.

Classification — *Pleosporaceae*, *Pleosporales*, *Dothideo-mycetes*.

Hyphae pale brown, smooth, branched and septate, 3–5 µm wide. **Conidiophores** erect, straight to flexuous, geniculate towards apex, brown, smooth, septate, up to 500 µm long, lateral or terminal, unbranched or sparingly branched. **Conidiogenous cells** intercalary and terminal, brown, smooth, polytretic with darkened scars. **Conidia** cylindrical to ellipsoidal, the third (or rarely second) cell from base often swollen and straight on one side and convex on the opposite side, rounded at the apex, 21–31 × 11–14 µm, 3-distoseptate, brown to dark brown, swollen cell larger and darker than others; **hila** conspicuous, slightly protuberant, thickened, darkened, 2–3 µm wide.

Culture characteristics — Colonies on potato dextrose agar (PDA) approx. 4.5 cm diam after 7 d at 25 °C, surface with little aerial mycelium, olivaceous black; reverse leaden black. Stromata abundant on maize leaf agar after 7 d at 25 °C, immersed and superficial, branched or unbranched.


Typus. AUSTRALIA, Queensland, Wivenhoe Pocket, on roots of *Stenotaphrum secundatum* (*Poaceae*), 19 Feb. 2020, N.T. Tran, A. Teo & A.D.W. Geering (holotype BRIP 71303, preserved as metabolically inactive culture, culture ex-type BRIP 71303, ITS, *gapdh* and *tef1* sequences GenBank MZ681952, MZ695824 and MZ695819, MycoBank MB 840702).

Additional materials examined. AUSTRALIA, New South Wales, Cornwallis, on leaves and roots of *S. secundatum*, 2 Mar. 2020, N.T. Tran & A.D.W. Geering, BRIP 71307 (ITS, *gapdh* and *tef1* sequences GenBank MZ681953, MZ695825 and MZ695820); Louth Park, *ibid.*, 4 Mar. 2020, N.T. Tran & A.D.W. Geering, BRIP 71316 (ITS, *gapdh* and *tef1* sequences GenBank MZ681956, MZ695828 and MZ695823); Maitland, *ibid.*, 4 Mar. 2020, N.T. Tran & A.D.W. Geering, BRIP 71314 (ITS, *gapdh* and *tef1* sequences GenBank MZ681955, MZ695827 and MZ695822); Wilberforce, *ibid.*, 3 Mar. 2020, N.T. Tran & A.D.W. Geering, BRIP 71309 (ITS, *gapdh* and *tef1* sequences GenBank MZ681954, MZ695826 and MZ695821).

Colour illustrations. Turf farm in South-East Queensland, Australia, where the type specimen of *C. stenotaphri* was collected. Colony of *Curvularia stenotaphri* on PDA; stroma; conidiophores on maize leaf on modified Sach's agar; conidiophores; conidia. Scale bars = 1 cm (colony), 1 mm (stroma), 100 µm (conidiophores on maize leaf), 10 µm (all others).

Notes — *Curvularia stenotaphri* was isolated from rotting roots and leaf spots of *Stenotaphrum secundatum* collected from localities in New South Wales and Queensland. *Curvularia stenotaphri* is distinguished from other species by comparison of sequences of ITS, *gapdh*, and *tef1* gene regions. *Curvularia stenotaphri* resembles many of the species with four-celled conidia and an asymmetrically swollen darker third cell, e.g., *C. lunata*, *C. hominis* and *C. muehlenbeckiae* (Madrid et al. 2014). Therefore, morphology alone does not reliably separate these and similar species.

Curvularia stenotaphri ITS megablast search showed similarities to *C. muehlenbeckiae* (GenBank MN893921.1; Identities = 530/532 (99 %), one gap (0 %)), *C. lunata* (GenBank MT444990.1; Identities = 530/532 (99 %), one gap (0 %)), and *C. lycopersici* (GenBank MT590310.1; Identities = 530/532 (99 %), one gap (0 %)). The *gapdh* megablast search showed similarities to *C. hominis* (GenBank LT715809.1; Identities = 527/536 (98 %), no gaps), *C. lunata* (GenBank AF081411.1; Identities = 496/506 (98 %), no gaps), and *C. plantarum* (GenBank MN053038.1; Identities = 525/534 (98 %), no gaps). The *tef1* megablast search showed similarities to *C. platzii* (ex-type strain BRIP 27703b, GenBank MH433669.1; Identities = 863/869 (99 %), no gaps), *C. polytrata* (ex-type strain CGMCC 3.19338, GenBank MN263984.1; Identities = 920/926 (99 %), no gaps), and *C. plantarum* (GenBank MN053007.1; Identities = 863/869 (99 %), no gaps).



Phylogenetic tree of selected *Curvularia* species based on a maximum likelihood analysis of a combined multilocus alignment (*gapdh*, ITS, and *tef1*). Analyses were performed on the Geneious v. 11.1.2 platform (Biomatters Ltd.) using RAXML v. 8.2.11 (Stamatakis 2014) and MrBayes v. 3.2.6 (Ronquist & Huelsenbeck 2003), both based on the GTR substitution model with gamma-distribution rate variation. Branch lengths are proportional to substitutions per site. RAXML bootstrap (bs) values > 70 % and Bayesian posterior probabilities (pp) > 0.8 are given on the branches (bs/pp). *Alternaria alternata* strain CBS 916.96 and *Bipolaris maydis* strain CBS 136.29 were used as outgroup. Novel taxon is highlighted in colour. Ex-type strains are marked with an asterisk (*).

Nga T. Tran & Andrew D.W. Geering, Centre for Horticultural Science, Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Dutton Park 4102, Queensland, Australia;
e-mail: n.tran3@uq.edu.au & a.geering@uq.edu.au

Yu Pei Tan, Plant Pathology Herbarium, Department of Agriculture and Fisheries, Dutton Park 4102, Queensland, Australia;
e-mail: yupeit.tan@daf.qld.gov.au

Roger G. Shivas, Centre for Crop Health, University of Southern Queensland, Toowoomba 4350, Queensland, Australia;
e-mail: roger.shivas@usq.edu.au



Fungal Planet 1347 – 24 December 2021

***Cytospora hippophaicola* Spetik, Eichmeier, Gramaje, Stuskova & Berraf-Tebbal, sp. nov.**

Etymology. Named after the host genus *Hippophae* from which the fungus was collected.

Classification — *Cytosporaceae*, *Diaporthales*, *Sordariomycetes*.

Conidiomata pycnidial, brown to black, solitary or aggregated with white to yellow conidial masses, surface covered by white mycelium. **Conidiophores** hyaline, straight or slightly curved, unbranched. **Conidiogenous cells** cylindrical to clavate, straight. **Conidia** hyaline, allantoid, aseptate, smooth-walled, $(3.5\text{--}4.0\text{--}5.0\text{--}5.5) \times (1.0\text{--}1.2\text{--}1.4\text{--}1.5) \mu\text{m}$, av. $\pm$ SD $4.3 \pm 0.2 \times 1.3 \pm 0.1 \mu\text{m}$ ($n = 30$).

Culture characteristics — Colony on potato dextrose agar (PDA) > 90 mm diam after 7 d at 22 °C, > 90 mm at 25 °C, no growth at 35 °C, medium dense, flat, fluffy mycelium, white, reverse white. On malt extract agar (MEA) > 90 mm diam after 7 d at 25 °C, dense, flat, fluffy mycelium with a uniform texture, white, reverse white. Colonies on oatmeal agar (OA) > 90 mm diam after 7 d at 25 °C, dense, flat, fluffy, mycelium forming slightly erumpent concentric circles, white, reverse white. Sexual morph not observed.

Typus. CZECH REPUBLIC, Lednice, isolated from the wood of asymptomatic *Hippophae rhamnoides* (*Elaeagnaceae*), Nov. 2020, M. Spetik (holotype BRNU 680004, ex-type culture CBS 147584 = MEND-F-0547; ITS, LSU, *tub2*, *tef1-α*, *act* and *rpb2* sequences GenBank MZ702814, MZ702873, MZ712165, MZ712155, MZ712150 and MZ712160, MycoBank MB 841161).

Additional material examined. CZECH REPUBLIC, Mutenice, isolated from the wood of symptomatic *Vaccinium corymbosum*, Feb. 2021, J. Pecenka (culture MEND-F-0554, ITS, LSU, *tub2*, *tef1-α*, *act* and *rpb2* sequences GenBank MZ702815, MZ702872, MZ712166, MZ712156, MZ712151 and MZ712161); Mutenice, isolated from the wood of symptomatic *V. corymbosum*, Feb. 2021, J. Pecenka (culture MEND-F-0555, ITS, LSU, *tub2*, *tef1-α*, *act* and *rpb2* sequences GenBank MZ702816, MZ702873, MZ712167, MZ712157, MZ712152 and MZ712162); Mutenice, isolated from the wood of symptomatic *V. corymbosum*, Feb. 2021, J. Pecenka (culture MEND-F-0558, ITS, LSU, *tub2*, *tef1-α*, *act* and *rpb2* sequences GenBank MZ702817, MZ702874, MZ712168, MZ712158, MZ712153 and MZ712163); Mutenice, isolated from the wood of symptomatic *V. corymbosum*, Feb. 2021, J. Pecenka (culture MEND-F-0559, ITS, LSU, *tub2*, *tef1-α*, *act* and *rpb2* sequences GenBank MZ702818, MZ702875, MZ712169, MZ712159, MZ712154 and MZ712164).

Notes — *Cytospora hippophaicola* was isolated for the first time from the wood of asymptomatic *Hippophae rhamnoides*. Subsequently, the same species has been isolated from the wooden tissues of symptomatic highbush blueberry (*Vaccinium corymbosum*). The symptoms included necrotic lesions under the bark, necrotic twigs and dieback of whole plants. Based on the phylogenetic data *C. hippophaicola* is closely related to *C. galegicola* (strain MFLUCC 18-1199). Both species can be distinguished based on ITS and *rpb2* sequence data. Morpho-

logically, the shape of conidia is similar to those of *C. galegicola*. However, *C. hippophaicola* produces smaller conidia, $4.3 \pm 0.2 \times 1.3 \pm 0.1$ vs $7.6 \times 1.3 \mu\text{m}$ in *C. galegicola*. Also, the colonies of *C. hippophaicola* on PDA grow faster > 90 mm diam at 22 °C after 7 d vs those of *C. galegicola* 5–5.5 cm diam after 15 d, at room temperature (Shang et al. 2020).

Based on a megablast search of NCBI's nucleotide database, the closest hits using the ITS sequence had the highest similarity to *Cytospora* sp. (strain H2/3b, GenBank MG020290.1; Identities = 545/546 (99 %), one gap (0.2 %)), *C. ambiens* (as *Valsa ambiens*; strain CBS 423.52, GenBank MH857109.1; Identities = 543/545 (100 %), no gaps) and *Cytospora* sp. (strain SL69_85_W4, GenBank MN105524.1; Identities = 543/546 (99 %), one gap (0.2 %)). The closest hits using the LSU sequence had the highest similarity to *C. ambiens* (as *Valsa ambiens*; strain AR3516, GenBank AF362564.1; Identities = 954/956 (99 %), one gap (0.1 %)), *V. cenisia* (strain AR3522, GenBank AF408385.1; Identities = 961/967 (99 %), one gap (0.1 %)) and *C. galegicola* (as *Cytospora* sp., strain MFLUCC 18-1199, GenBank MK571756.1; Identities = 803/803 (100 %), no gaps); closest hits using the *rpb2* sequence are *C. ambiens* (as *V. ambiens*; strain AFTOL-ID 2131, GenBank DQ862025.1; Identities = 913/913 (100 %), no gaps), *C. ambiens* (as *V. ambiens*; strain AR3516, GenBank EU219347.1; Identities = 891/891 (89 %), no gaps) and *C. galegicola* (as *Cytospora* sp., strain MFLUCC 18-1199, GenBank MN685820.1; Identities = 768/795 (97 %), 14/795 (1 %) gaps). The closest hits using the *tef1-α* sequence had the highest similarity to *Cytospora* sp. (strain ColPat634, GenBank MK490945.1; Identities = 287/300 (96 %), one gap (0.3 %)), *Cytospora* sp. (strain ColPat633, GenBank MK490944.1; Identities = 287/300 (97 %), one gap (0.3 %)) and *C. longispora* (strain 10F-57, GenBank MG971615.1; Identities = 288/311 (93 %), 8 gaps (4 %)). The closest hits using the *act* sequence had the highest similarity to *C. ribis* (strain CFCC 50026, GenBank KP310843.1; Identities = 249/254 (98 %), one gap (0.4 %)), *Cytospora* sp. (strain XJAU2776_1, GenBank MN044591.1; Identities = 251/260 (97 %), two gaps (0.8 %)) and *C. galegicola* (as *Cytospora* sp., strain MFLUCC 18-1199, GenBank MN685810.1; Identities = 215/215 (100 %), no gaps); closest hits using the *tub2* sequence are *C. ribis* (strain CFCC 50026, GenBank KP310826.1; Identities = 682/705 (95 %), six gaps (0.9 %)), *C. ribis* (strain CFCC 50039, GenBank KR045683.1; Identities = 603/669 (90 %), 31 gaps (4 %)) and *C. carbonacea* (strain CFCC 50058, GenBank KP310823.1; Identities = 623/709 (88 %), 28 gaps (4 %)).

Colour illustrations. *Hippophae rhamnoides* shrub in Lednice, Czech Republic. Conidia; section through conidiomata; conidiomata on WA; culture on PDA, MEA and OA. Scale bars = 1 mm (conidiomata on WA), 100 μm (section), 5 μm (all others).

Supplementary material**FP1347** Phylogenetic tree.

Milan Spetik, Ales Eichmeier, Katerina Stuskova & Akila Berraf-Tebbal, Mendeleum – Institute of Genetics, Faculty of Horticulture, Mendel University in Brno, Valticka 334, Lednice, 69144, Czech Republic;

e-mail: milan.spetik@mendelu.cz, ales.eichmeier@mendelu.cz, xstusko1@mendelu.cz & qqberraf@mendelu.cz

David Gramaje, Instituto de Ciencias de la Vid y del Vino (ICVV), Consejo Superior de Investigaciones Científicas (CSIC) – Universidad de La Rioja – Gobierno de La Rioja, Ctra. LO-20, Salida 13, 26007, Logroño, Spain; e-mail: david.gramaje@icvv.es



Fungal Planet 1348 – 24 December 2021

Didymella tabebuicola W.A.S. Vieira, I.G. Duarte, A.G.G. Amaral, A.F. Lima & M.P.S. Câmara, *sp. nov.*

Etymology. Name refers to the host *Tabebuia aurea*, from which it was isolated.

Classification — *Didymellaceae*, *Pleosporales*, *Dothideo-mycetes*.


On potato dextrose agar (PDA). *Conidiomata* pycnidial, separate, dark brown, globose, unilocular, with central ostiole, 55.4–100.5 µm high, 44.4–87.8 µm diam. *Chlamydospores* unicellular, yellowish brown, intercalary, in chains, globose to subglobose, thick-walled, 4.5–6.6 µm diam. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* enteroblastic, phialidic, ampulliform, determinate, hyaline, smooth-walled, (5.1–)5.3–7.7(–9.5) × (2.7–)3–4.2(–4.5) µm, av. ± S.D. 6.5 ± 1.6 × 3.6 ± 0.6 µm. *Conidia* solitary, hyaline, aseptate, smooth, guttulate, subcylindrical, straight, apex slightly obtuse, base truncate, (3.9–)4.2–4.9(–5.8) × (1.6–)1.7–2(–2.5) µm, mean ± S.D. 4.6 ± 0.4 × 1.9 ± 0.2 µm. *Sexual morph* not observed.

Culture characteristics — Colonies on PDA, 27–36 mm diam after 7 d, margin regular, covered by floccose aerial mycelia, olivaceous, with white edges; reverse greyish sepia (Rayner 1970).

Typus. BRAZIL, Pernambuco, Recife, from leaf spots on *Tabebuia aurea* (*Bignoniaceae*), 3 Feb. 2020, W.A.S. Vieira (holotype VIC 47504, culture ex-type COAD 3340 = LM1189; ITS, LSU, *rpb2* and *tub2* sequences GenBank MZ703618, MZ703623, MZ712360 and MZ712364; MycoBank MB 841209).

Additional material examined. BRAZIL, Pernambuco, Recife, from leaf spots on *T. aurea*, 3 Feb. 2020, W.A.S. Vieira, LM1187, LM1188, LM1190, ITS (MZ703616, MZ703617, MZ703619), LSU (MZ703621, MZ703622, MZ703624), *rpb2* (MZ712358, MZ712359, MZ712361) and *tub2* (MZ712362, MZ712363, MZ712365) sequences GenBank.

Notes — Similar *T. aurea* leaf spots were reported by Viana et al. (2012) and associated with a *Phoma* sp. based on phenotypic features. *Didymella tabebuicola* is closely related with *D. coffeae-arabicae*, *D. keratinophila*, *D. mitis* and *D. sancta*, species previously identified as *Phoma* from section *Peyronellaea* (Aveskamp et al. 2009, Chen et al. 2017, Hou et al. 2020). Based on BLASTn searches, the ITS sequences had highest similarity to *D. keratinophila* (strain UTHSC DI16-200, GenBank NR_158275.1, identity = 494/498 (99 %), 1 gap). LSU sequences had highest similarity to *D. heteroderae* (strain CBS 109.92, GenBank GU238002.1, Identities = 852/852(100 %), no gaps). *rpb2* sequences had highest similarity to *D. mitis* (strain CBS 443.72, GenBank MT018137.1, Identities = 575/595(97 %), no gaps). *tub2* sequences had highest similarity to *D. keratinophila* (strain UTHSC DI16-200, GenBank NR_158275.1, Identities = 331/341(97 %), no gaps). However, *D. tabebuicola* is recovered as monophyletic in all independent gene trees.



Colour illustrations. *Tabebuia aurea* growing on campus of Universidade Federal Rural de Pernambuco, where the species was collected. *Tabebuia aurea* symptomatic leaf; aerial and reverse view from colonies on PDA; pycnidium; chlamydospores; conidiogenous cells; conidia. Scale bars = 1 cm (leaf and colonies), 10 µm (pycnidium), 5 µm (all others).

Maximum Likelihood tree inferred with RAxML (Stamatakis 2014) from combined ITS, LSU, *rpb2* and *tub2* sequences of 51 *Didymella* isolates. Ex-types are highlighted in **bold**. Bootstrap support values from ML ≥ 70 % are shown above the branches. The tree is rooted at the midpoint.

Willie A.S. Vieira, Ingrid G. Duarte, Ana G.G. Amaral, Athaise F. Lima, Marcos P.S. Câmara,

Universidade Federal Rural de Pernambuco, Rua Dom Manuel de Medeiros s/n, 52171-900, Recife, Pernambuco, Brazil;

e-mail: andersonvieira12@gmail.com, ingridgduarte@gmail.com, gabriele.160713@gmail.com, athaiseff17@gmail.com & marcos.camara@ufrpe.br

Entoloma cinnamomeum



Fungal Planet 1349 – 24 December 2021

***Entoloma cinnamomeum* O.V. Morozova, Vila, Finy, D. Ageev & Dima, sp. nov.**

Etymology. From Latin *cinnamomeus*, referring to the cinnamon brown colour of the young and fresh basidiomata.

Classification — *Entolomataceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata medium-sized, collybioid. **Pileus** 15–25 mm diam, hemispherical, then expanding to convex and plano-convex, sometimes slightly depressed, with deflexed then straight margin, faintly hygrophanous, translucently striate up to half of the radius in well-developed specimens, densely radially fibrillose-squamulose in the centre and around it, with squamules more distant and finely towards the margin, moderately dark brown, orange-brown (5D4–6, 5E4–7, 6D4–6, 6E4–7; Kernerup & Wanscher 1978), sometimes with red brown tones (cinnamon) especially in young specimens, with darker contrasting spot in the centre and darker radial stripes, paler in dry conditions (5C3–4, 6C3–4). **Lamellae** moderately distant, adnate-emarginate, whitish, pale or beige, becoming greyish pink, with irregular, concolourous or whitish edge. **Stipe** 30–60 × 1.5–2 mm, cylindrical or slightly broadened towards the base, almost glabrous, polished, sometimes minutely longitudinally striate, with pruinose apex especially in young specimens, concolourous with the pileus in the upper part, contrasting with whitish lamellae, greyish brown to greyish at the base or completely greyish (5C2–3, 5D3–4), sometimes yellowish in the base, with white tomentum in the lower part. **Context** white, brownish under the surface of the pileus. **Smell** indistinct, **taste** not reported. **Basidiospores** (9–)10–11.5(–12.5) × (6–)7–8(–8.5) µm, Q = (1.3–)1.4–1.5(–1.8), heterodiametrical, with 5–7 angles in side-view. **Basidia** 28–34 × 9.5–10.5 µm, 4-spored, narrowly clavate to clavate, clampless. **Lamella edge** fertile or heterogeneous. **Cheilocystidia** absent, but some cystidia like cylindrical cells can be observed. **Hymenophoral trama** regular, made up of 4–15 µm wide, cylindrical hyphae. **Pileipellis** a cutis with transition to a trichoderm, a trichoderm in the centre of cylindrical hyphae, 5–12 µm wide with ascending more or less inflated to fusiform or ellipsoid, sometimes constricted terminal elements, 30–110 × 6–30 µm, with yellow-brown in KOH intracellular pigment. **Stipitipellis** a cutis of cylindrical parallel hyphae with some clusters of cylindrical hairs in the apex of the stipe. **Clamp connections** absent.


Habit, Habitat & Distribution — In small groups on soil on calcareous grasslands. Known from Russia (European part and West Siberia), Hungary and Spain (Catalonia).

Colour illustrations. Russia, Tver' region, Zubtsov District, calcareous grassland in the right bank of the Derzha River (type locality). Spores; pileipellis; stipitipellis and basidiomata in situ (from holotype); basidiomata in situ (FP-2014-09-20). Scale bars = 1 cm (basidiomata), 10 µm (spores and microstructures).

Typus. RUSSIA, Tver' Region, Zubtsov District, 2 km to SE from Mozgovo Village, on soil in the calcareous grassland in the right bank of the Derzha River, N56.213861° E34.8025°, 11 Sept. 2015, O.V. Morozova (holotype LE 311801; ITS and LSU sequences GenBank OK161243 and OK161274, MycoBank MB 841838). Previously reported as *Entoloma longistriatum* (Morozova et al. 2016).

Additional material examined. HUNGARY, Tolna County, Bikács, on dry, sandy, calcareous grassland, 20 Sept. 2014, P. Finy (FP-2014-09-20, ITS sequence GenBank OK161245). — RUSSIA, Altay Republic, Chemalinsky District, vicinities of Elanda Village, bank of Katun' River, N51.229145° E86.085518°, on soil at the riverbank, 22 Aug. 2019, D. Ageev (LE 312535, ITS sequence GenBank OK161246). — SPAIN, Catalonia, el Far, Sant Martí Sacalm (Selva, Girona), 910 m a.s.l., among mosses and grasses, in an open area of a *Quercus humilis* woods, on basic soil, 6 June 2014, J. Vila & X. Llimona (JVG 1140606-10, ITS sequence GenBank OK161244).

Notes — *Entoloma cinnamomeum* nests within the *sarcitulum* clade. Morphologically it corresponds to the main characteristics of this clade – predominance of the brown colours in the whole basidiomata, deeply translucently striate pileus, and polished stipe. The presence of the greyish tinge near the stipe base makes it similar to *E. longistriatum* in the broad sense, including *E. majusculum* (Noordeloos 1992). *Entoloma cinnamomeum* can be distinguished from representatives of this group by the absence of cheilocystidia and darker colour, sometimes with red brown tones.



Phylogenetic tree derived from Maximum Likelihood analysis based on nrITS1-5.8S-ITS2 sequence data. Analysis was performed in PhyML v. 3.0 (Guindon et al. 2010) using the non-parametric Shimodaira-Hasegawa version of the approximate likelihood-ratio test (SH-aLRT) and the GTR+I+Γ model of evolution. ML bootstrap support (BS) values are shown at the nodes (BS > 70 %).

Olga V. Morozova, Komarov Botanical Institute of the Russian Academy of Sciences, 197376, 2 Prof. Popov Str., Saint Petersburg, Russia; e-mail: OMorozova@binran.ru

Jordi Vila, Passatge del Torn, 4, 17800 Olot, Spain; e-mail: micocistus@hotmail.com

Péter Finy, Zsombolyai u. 56, 8000 Székesfehérvár, Hungary; e-mail: peter@finy.hu

Dmitry V. Ageev, LLC 'Signatec', 630090, Inzhenernaya Str. 22, Novosibirsk, Russia; e-mail: dim@diamondsteel.ru

Bálint Dima, Department of Plant Anatomy, Institute of Biology, Eötvös Loránd University, Pázmány Péter sétány 1/C, H-1117, Budapest, Hungary; e-mail: cortinarius1@gmail.com



Fungal Planet 1350 – 24 December 2021

***Entoloma kovalenkoi* O.V. Morozova, E.S. Popov & A.V. Alexandrova, sp. nov.**

Etymology. Named in honour of Alexander Kovalenko, an exceptional Russian mycologist, who headed the Laboratory of the Systematics and Geography of Fungi of the Komarov Botanical Institute RAS for many years, who brought together all professional and amateur Russian mycologists, and was the initiator and inspirer of the study of the mycobiota of Vietnam by Russian scientists as well as of many other significant projects.

Classification — *Entolomataceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata medium-sized, inocyboid. **Pileus** 10–25 mm diam, firstly campanulate, then conical, broadly conical without distinct papilla, with deflexed then straight, slightly crenulate or ragged margin, not hygrophanous, not translucently striate, entirely radially fibrillose to hairy, light- to deep orange (5A5–8, 6A7–8; Kernerup & Wanscher 1978). **Lamellae** moderately distant, adnate-emarginate, almost free, ventricose, pale orange, salmon (5A4–5), with minutely serrate whitish edge. **Stipe** 40–90 × 2–4 mm, cylindrical or slightly broadened towards the base, hollow, longitudinally fibrillose, pale yellow, yellowish orange, paler than pileus (4A5–7; 5A4–5), white tomentose at base. **Context** pale orange, salmon, exuding sparse yellow latex. **Smell** indistinct, **taste** not reported. **Basidiospores** (8–)9–10(–11) × (7–)8–9(–10) µm, Q = (1–)1.1(–1.2), isodiametrical, cuboid, rarely with five angles in side-view. **Basidia** 45–59 × 13–19.5 µm, 4-spored, clavate, constricted in the middle part, clamped. **Cheilocystidia** 46–95 × 11–18 µm, fusoid, lageniform, clavate or subcapitate with slightly attenuate apex, sometimes septate, thin-walled, not pigmented, forming sterile lamellae edge. **Lamellae trama** regular, consists of cylindrical hyaline hyphae 4–10 µm broad and abundant bright yellow oleiferous hyphae. **Pileipellis** a transition between cutis and trichoderm of cylindrical hyphae up to 10 µm broad with ascending cylindrical to fusoid terminal elements up to 15 µm broad. **Pileitrama** consists of cylindrical hyphae 4–6 µm broad with yellowish intracellular pigment, with abundant bright yellow oleiferous hyphae. **Clamp connections** present in all tissue.


Habitat & Distribution — In small groups on rotten wood in middle mountain evergreen mixed forest. Known from Vietnam.

Typus. VIETNAM, Gia Lai Province, K'Bang District, Son Lang Commune, Kon Chu Rang Nature Reserve, N14.49328° E108.54419°, 970 m a.s.l., on rotten wood in middle mountain evergreen mixed forest with a predominance of *Podocarpaceae* (*Dacrydium elatum*, *Dacrycarpus imbricatus*), *Magnoliaceae*, *Burseraceae* (*Canarium*), *Myrtaceae* (*Syzygium*), 27 May 2016, E.S. Popov & O.V. Morozova (holotype LE 312529, ITS, LSU, mtSSU and *tef1α* sequences GenBank OK257210, OK257207, OK256168 and OK256169, MycoBank MB 841775).

Additional specimens examined. VIETNAM, Gia Lai Province, K'Bang District, Son Lang Commune, Kon Chu Rang Nature Reserve, N14.49328° E108.54419°, 970 m a.s.l., on rotten dead wood in middle mountain evergreen mixed forest with a predominance of *Podocarpaceae* (*Dacrydium elatum*, *Dacrycarpus imbricatus*), *Magnoliaceae*, *Burseraceae* (*Canarium*), *Myrtaceae* (*Syzygium*), 1 June 2016, O.V. Morozova & A.V. Alexandrova (LE 312530, ITS and LSU sequences GenBank OK257211 and OK257208).


Colour illustrations. Vietnam, Gia Lai Province, Kon Chu Rang Nature Reserve, middle mountain evergreen mixed forest, type locality. Spores; cheilocystidia; pileipellis; basidiomata (from holotype); longitudinal section of the basidioma (LE 312530). Scale bars = 1 cm (basidiomata), 50 µm (pileipellis), 10 µm (spores and cheilocystidia).

Notes — *Entoloma kovalenkoi* is characterised by bright orange basidiomata with entirely radially fibrillose not hygrophanous pileus, longitudinally striate stipe, cuboid spores and fusoid, lageniform, clavate or capitate cheilocystidia with attenuate apex. Among the species with cuboid spores, similar cheilocystidia have been observed in *E. latericolor* (Horak 1976). However, *E. latericolor* has a cinnamon to brick red pileus and is known from New Zealand. *Entoloma quadratum* (= *E. salmoneum*), also coloured in orange tones, possesses a glabrous hygrophanous surface of the pileus becoming rimose-fibrillose with age and another type of cheilocystidia – cylindrical to narrowly clavate cells in dense clusters up to 120 µm long of 'serrulatum' type (Horak 1976, Noordeloos & Morozova 2010, Morozova et al. 2012). The new species resembles the blue-coloured species of *Entoloma virescens* complex by the not hygrophanous radially fibrillose pileus and abundant yellow milky juice, leading to a change in the colour of the context when damaged. Phylogenetically the new species is rather distant from all known *Entoloma* species with cuboid spores from subgenus *Cubospora* (Karstedt et al. 2019).



Phylogenetic tree derived from Bayesian analysis, based on mtSSU sequence data. Analysis was performed under the GTR model of evolution, for 3 M generations, using MrBayes v. 3.2.1 (Ronquist et al. 2012). Posterior probability (PP > 0.94) values from the Bayesian analysis are added at the nodes. The scale bar represents the expected number of nucleotide changes per site.

Entoloma verae



Fungal Planet 1351 – 24 December 2021

Entoloma verae O.V. Morozova, Noordel., Reschke, F. Salzmann & Dima, *sp. nov.*

Etymology. Named in honour of the Russian mycologist Vera Malysheva, as one of the collectors of the type specimen of the new species and one of the first investigators of fungi of Zhiguli Mts, the type locality.

Classification — *Entolomataceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata medium-sized, collybioid. **Pileus** 15–40 mm diam, hemispherical with slightly depressed centre, then expanding to convex and plano-convex with slightly umbilicate centre, with deflexed then straight margin, slightly hygrophanous, translucently striate usually up to 1/2–2/3 of radius, olive, olive yellow, greyish yellow, yellow-green, with darker striae and centre (2B5–8, 2C4–8, 2D4–6; Kornerup & Wanscher 1978), covered with brownish scales, especially in the centre, becoming smooth towards the margin. **Lamellae** moderately distant, adnate-emarginate, with 3–5 lamellulae, adnate, decurrent with short tooth, slightly emarginate to arcuate, whitish, becoming pinkish, with entire or irregular, concolourous edge. **Stipe** 20–70 × 1–3 mm, cylindrical or slightly broadened towards the base, smooth, polished, yellowish green (29A6–8), staining bright greenish blue when bruised, white tomentose at the base. **Context** greenish, discolouring to bright greenish blue when bruised. **Smell** indistinct or not reported, **taste** not reported. **Basidiospores** 10–15.5 × 7–9 µm, av. 12.5 × 8 µm, Q = 1.3–2.0, Q_{av} = 1.6, heterodiametrical with 6–8 angles in side-view. **Basidia** 33.5–39 × 9–11.5 µm, 2–4-spored, clavate, clampless. **Lamella edge** fertile or heterogeneous, composed of basidiola-like clavate cells intermixed with rare basidia. **Cheilocystidia** not differentiated. **Hymenophoral trama** regular, made up of 5–10 µm wide, cylindrical hyphae. **Pileipellis** a trichoderm to hymenoderm in the central part with elongated to spherical terminal elements, 40–150 × 20–35 µm, a cutis with transition to a trichoderm towards the margin composed of ascending cylindrical to fusiform hyphae, 10–20 µm broad. **Caulocystidia** absent. **Clamp connections** absent.

Habit, Habitat & Distribution — Solitary or in small groups on soil on calcareous grasslands. Known from Russia (European part and Caucasus), Germany, the Netherlands and China (GenBank JQ281488).

Colour illustrations. Russia, Samara region, the bank of the Volga River, Zhigulevsky Nature Reserve (type locality, photo credit V. Malysheva). Basidioma *in situ* from L-0607931 (photo credit F. & R. Salzmann); spores; cheilocystidia; pileipellis in central part and near the margin of pileus (all from holotype). Scale bars = 1 cm (basidiomata), 10 µm (spores and microstructures).

Typus. RUSSIA, Samara Region, Stavropolsky District, Zhiguli Nature Reserve, vicinities of Bakhilova Polyana Village, way to Gudronny Village, N53.414444° E49.73425°, 16 Sept. 2003, *E. Malysheva & V. Malysheva* (holotype LE 227613; ITS and LSU sequences GenBank OK161251 and OK161277, MycoBank MB 841839).

Additional materials examined. GERMANY, Heimberg, near Schloss Böckelheim, oligotrophic meadow, 27 Oct. 2017, *K. Reschke* (KaiR990, ITS sequence GenBank MZ611683). – THE NETHERLANDS, prov. Limburg, Nijswiller-Noord, N50.81533° E5.95455°, 21 Aug. 2019, leg. *F. & R. Salzmann* (L-0607931, ITS sequence GenBank OK161252) – RUSSIA, Karachaevo-Cherkesia Republic, Teberda Nature Reserve, Jamagat gorge, N43.45374° E41.82818°, a.s.l. 1880 m, 13 Aug. 2009, leg. *O. Morozova* (LE 312551, ITS sequence GenBank OL744072)..

Notes — During the phylogenetic study of the subgenus *Cyanula*, two genotypes fitting the current morphological concept of *Entoloma incanum* were revealed with 14 substitution and indel differences between them. For the one that is widespread in Europe (including the country of the type locality in Sweden), Siberia and the Far East of Russia, i.e., *Entoloma incanum* s.str., we have chosen a neotype corresponding to the protologue (Fries 1821) and the current understanding of the species (Noordeloos 1992):

Agaricus incanus Fr., Syst. mycol. (Lundae) 1: 209 (1821) MycoBank: MB 372149. Neotype (designated here, MBT 10003730): Sweden, Östersund, Torvalla, Ängsmon västra, Tysjöarna NR, 2 Sept. 2016, *O. Morozova* (LE 312503, ITS and LSU sequences GenBank OK161247 and OK161275).

The second species is described here as new to science. *Entoloma verae* differs morphologically from typical *E. incanum* by the generally larger spores measuring 12.5 × 8 µm, on average, vs 10.5 × 7.5 µm on average in the neotype. But the size of spores can vary significantly, sometimes related to the presence of the 2-spored basidia.

See the phylogenetic tree provided in Fungal Planet 1349 elsewhere in this paper.

Olga V. Morozova, Komarov Botanical Institute of the Russian Academy of Sciences, 197376, 2 Prof. Popov Str., Saint Petersburg, Russia; e-mail: OMorozova@binran.ru

Machiel E. Noordeloos, Naturalis Biodiversity Center, section Botany, P.O. Box 9517, 2300 RA Leiden, The Netherlands; e-mail: m.noordeloos@mac.com

Kai Reschke, Mycology Research Group, Faculty of Biological Sciences, Goethe University Frankfurt am Main, Max-von-Laue Straße 13, 60439 Frankfurt am Main, Germany; e-mail: Reschke@em.uni-frankfurt.de

Finy Salzmann, Kloosterweg 5 6301 WK, Valkenburg a/d Geul, The Netherlands; e-mail: reisalzmann@gmail.com

Bálint Dima, Department of Plant Anatomy, Institute of Biology, Eötvös Loránd University, Pázmány Péter sétány 1/C, H-1117, Budapest, Hungary; e-mail: cortinarius1@gmail.com



Fungal Planet 1352 – 24 December 2021

***Entomortierella hereditatis* J. Trovão, J. Pawłowska & A. Portugal, sp. nov.**

Etymology. Latin for the word 'heritage', in honour of the UNESCO World Heritage Site of 'University of Coimbra: Alta and Sofia' where the isolate was collected.

Classification — *Mortierellaceae*, *Mortierellales*, *Mortierellomycetes*.

Hyphae hyaline, 2–3 µm wide, forming abundant solitary gemmae both on aerial and substrate hyphae. Weakly branched aerial hyphae reach up to 900 µm in length, containing multiple gemmae. **Gemmae** intercalary and terminal on 2–3 µm wide stalks, filled with oil drops, hyaline, globose, (8.3–)11.7(–19.4) (SD = 2.31) µm diam, with smooth cell wall, (0.83–)1.2(–1.4) (SD = 0.32) µm thick. No sexual or asexual reproduction structures detected in 2 % potato dextrose agar (PDA), half and quarter strength PDA, 2 % water agar (WA), malt extract agar, potato carrot agar, Sabouraud dextrose agar, Miura's medium and 0.3 % Gerber baby food agar. The species is a non-halotolerant, facultative alkaliphile able to grow under pHs from 6 to 11.

Culture characteristics — *Colonies* on PDA, after 7 d at 25 °C in the dark: colonies radiate, fast-growing, with daily increments of 5–7 mm, reaching up to 45 mm diam, with typical slightly zonate growth and lobate pattern on the edge; no garlic odour detected. Colonies dull white on top and yellowish in reverse. On WA, the characteristic colony shape does not develop.

Typus. PORTUGAL, Coimbra, isolated from a biofilm covering a deteriorated limestone wall in the Old Cathedral of Coimbra, 22 Nov. 2016, *I. Tiago* (holotype MUM-H 18.47, culture ex-type MUM 18.47 = DSM 106915, ITS and LSU sequences GenBank MG938350 and MG932787, MycoBank MB 825064).

Additional material examined. PORTUGAL, Coimbra, isolated from a biofilm covering a deteriorated limestone wall in the Old Cathedral of Coimbra, 22 Nov. 2016, *I. Tiago*, MUM 20.28, ITS sequence GenBank MG938351; *ibid.*, MUM 20.29, ITS sequence GenBank MG938352.

Colour illustrations. The sampled biofilm in the Old Cathedral of Coimbra (photo credit J. Trovão). Seven-day-old colony on PDA; abundant gemmae with oil droplets; chains of gemmae. Scale bars = 100 µm (gemmae with oil droplets), 50 µm (chains of gemmae).

Notes — The phylogenetic analyses shows that this isolate belongs to the genus *Entomortierella* (Vandepol et al. 2020), representing 98 % LSU sequence similarity to *Entomortierella bejakovae* (CBS 274.71; CBS 275.71; CBS 109658; CBS 102878; CBS 123.72; CBS 806.68; CBS 109595; CBS 267.71) and 85 % ITS sequence similarity to '*Mortierella*' *calciphila* (WA18944). As ITS sequence similarity of strains MUM-H 18.47 and WA18944 is much lower than 97 %, proposed by Nagy et al. (2011), we propose to delimit it as a separate species. Both *E. hereditatis* and '*Mortierella*' *calciphila* are linked to limestone habitats. Nonetheless, '*Mortierella*' *calciphila* was isolated from soil with the presence of limestone and *E. hereditatis* from a biofilm covering a biodeteriorated dolomitic limestone monument. Additionally, in similar conditions '*Mortierella*' *calciphila* produced asexual reproduction structures while *E. hereditatis* is morphologically simple, without any sexual and asexual reproductive forms. In *Entomortierella*, the lack of reproduction structures was also observed in '*Mortierella*' *formicae*, a species isolated from ants (Hyde et al. 2017). Moreover, *E. hereditatis* forms single solitary gemmae, while '*Mortierella*' *calciphila* and '*Mortierella*' *formicae* produce gemmae in clusters. The gemmae of *E. hereditatis* are also smaller than '*Mortierella*' *calciphila*, usually reaching 11–25(–18) µm; *Entomortierella bejakovae*, usually reaching 20–45(–60) µm; and '*Mortierella*' *formicae*, usually reaching 8.7–23.43(–18.66) µm (Liu et al. 2016, Hyde et al. 2017). Sometimes, superficial hyphae segments are converted into elongated chlamydospores with oil drops and thick wall as similarly observed in cultures of '*Mortierella*' *formicae*. The characteristic *E. hereditatis* single gemmae can be also observed in some other *Mortierellaceae* species, e.g., '*Mortierella*' *paraensis* (Khalabuda 1973, Gams 1976, Domsch et al. 1993). However, the phylogenetic analysis clearly supports the separation of this lineage from other representatives of this genus.

Supplementary material

FP1352-1 Phylogenetic tree.

FP1352-2 Phylogenetic tree.

João Trovão & Igor Tiago, Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, 3004-531 Coimbra, Portugal; e-mail: jtrovaosb@gmail.com & itiago@uc.pt.

António Portugal, Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, 3004-531 Coimbra, Portugal; and Fitolab - Laboratory for Phytopathology, Instituto Pedro Nunes, 3030-199 Coimbra, Portugal; e-mail: aportuga@bot.uc.pt

Julia Pawłowska, Institute of Evolutionary Biology, Faculty of Biology, Biological and Chemical Research Centre, University of Warsaw, ul. Żwirki i Wigury 101, 02-089 Warsaw, Poland; e-mail: julia.z.pawlowska@uw.edu.pl

Fusarium chuoi



Fungal Planet 1353 – 24 December 2021

***Fusarium chuoi* R. Hill, Gaya, D.T. Vu, Sand.-Den. & Crous, sp. nov.**

Etymology. From *chuối*, Vietnamese vernacular name for *Musa* spp., from which the ex-type strain was isolated.

Classification — *Nectriaceae*, *Hypocreales*, *Sordariomycetes*.

On SNA and CLA, sporulation abundant from aerial conidiophores and sporodochia. *Aerial conidiophores* erect or prostrate, copiously branching laterally and sympodially, giving rise to macro-, and rarely, microconidia; *aerial conidiogenous cells* mono- and polyphialidic, subulate to subcylindrical, smooth- and thin-walled, proliferating sympodially, 6.5–40.5 × 2.5–4 µm, with apical flared collarette and periclinal thickening; aerial conidia of two types: *microconidia* often produced on prostrate conidiophores, rarely on aerial mycelium, aggregating in false heads, ellipsoidal, subcylindrical to slightly falcate, 0–1-septate, 8–15 × 2–29.5 µm; *macroconidia* fusiform to falcate, straight to apically dorsiventrally curved, apex curved to pointed, base obtuse to papillate, 1–3-septate, smooth- and thin-walled; 1-septate conidia: (14–)18–27.5(–29.5) × (2.5–)3–4 µm (av. 22.8 × 3.2 µm); 2-septate conidia: 26–28.5 × 3–4 µm (av. 27.4 × 3.6 µm); 3-septate conidia: (28–)31.5–43(–50.5) × 3–4 µm (av. 37.3 × 3.5 µm). *Sporodochia* saffron, luteous to ochreous coloured (Rayner 1970), formed abundantly on the agar surface and carnation leaves under nuv. *Conidiophores in sporodochia*, densely and irregularly branched, bearing apical whorls of 2–4 monophialides; sporodochial monophialides subcylindrical, 10–26 × 2.5–4.5 µm, smooth- and thin-walled, with a distinct apical collarette. *Sporodochial conidia* (macroconidia) falcate, almost straight to gently curved, tapering at both ends, apex curved to blunt, base poorly- to well-developed foot-shaped, 1–6-septate, hyaline, smooth- and thin-walled; 1-septate conidia: (14.5–)15–20.5(–24) × 3–4.5 µm (av. 17.9 × 3.9 µm); 2-septate conidia: 21.5–32 × 3–4.5 µm (av. 26.4 × 3.5 µm); 3-septate conidia: (33–)43–61(–71.5) × (3–)4–5 µm (av. 51.8 × 4.2 µm); 4-septate conidia: (50.5–)55–69(–74.5) × 3.5–5 µm (av. 62.3 × 4.2 µm); 5-septate conidia: 54 × 4.5 µm (rare); 6-septate conidia: (49.5–)56.5–71(–73) × (3.5–)4–4.5(–5) µm (av. 63.8 × 4.3 µm). *Chlamydospores* not observed.

Culture characteristics — Colonies on potato dextrose agar (PDA) and oatmeal agar (OA) growing in the dark at 24 °C covering and entire 9 cm Petri dish in 7 d. Colony surface peach to vinaceous, flat, velvety to felty with abundant floccose aerial mycelium forming concentric rings; colony margins undulate. Reverse flesh to salmon with diffuse coral to brick pigment throughout the medium.

Typus. VIETNAM, Hà Tĩnh Province, Hương Sơn District, Sơn Kim commune, N18°25'37.38" E105°12'53.95", inside seed of *Musa itinerans* (*Musaceae*), 9 Nov. 2014, D.M. Thu, L.T. Phong & T.T. Duong, isol. R. Hill (holotype CBS H-24901, culture ex-type CBS 148464; ITS, LSU, *cmdA*, *rpb1*, *rpb2*, *tef1* and *tub2* sequences GenBank OK586454, OK586452, OK626304, OK626306, OK626302, OK626308 and OK626310, MycoBank MB 841865).

Colour illustrations. Flowers, fruits, leaves and seeds of *Musa itinerans* (background photo by D.T. Vu); from top to bottom and left to right: colony on PDA after 14 d at 24 °C in darkness (left = obverse, right = reverse), sporodochia formed on CLA, aerial conidiophore, aerial conidiogenous cells, aerial conidia, sporodochial conidia. Scale bars: black = 20 µm, white = 10 µm.

Additional material examined. VIETNAM, Nghệ An Province, Con Cuông District, Châu Khê commune, N19°1'48.73" E104°43'31.97", inside seed of *M. itinerans*, 18 Nov. 2014, L.T. Phong, V.V. Tung & T.T. Duong, isol. R. Hill (culture CBS 148465; ITS, LSU, *cmdA*, *rpb1*, *rpb2*, *tef1* and *tub2* sequences GenBank OK586455, OK586453, OK626305, OK626307, OK626303, OK626309 and OK626311).

Notes — *Fusarium chuoi* resides in the Asian clade of the *Fusarium fujikuroi* species complex (FFSC: O'Donnell et al. 1998, Yilmaz et al. 2021, Crous et al. 2021b). Based on nucleotide searches using the *Fusarium* Pairwise ID engine on the Fusarioid-ID database (www.fusarium.org, Crous et al. 2021) the closest hit using the ITS sequence was *Fusarium siculi* (strain CBS 142422; identities = 449/450 (99 %), no gaps). The closest hit using the LSU sequence was *F. siculi* (strain CBS 142422; identities = 804/805 (99 %), no gaps). Closest hit using the *cmdA* sequence was *Fusarium fractiflexum* (strain NRRL 28852; identities = 426/434 (98 %), no gaps). Closest hit using the *rpb1* sequence was *F. fujikuroi* (strain NRRL 13566; identities = 687/702 (98 %), no gaps). Closest hit using the *rpb2* sequence was *Fusarium globosum* (strain CBS 428.97; identities = 856/867 (98 %), no gaps). Closest hit using the *tef1* sequence was *F. fractiflexum* (strain NRRL 28852; identities = 619/643 (96 %), 2 gaps (0.3 %)). The phylogenetic results, however, showed that *F. chuoi* is not directly related to any of the previously described species of FFSC (see Suppl. material FP1353), clustering as the second basal-most species of that clade after *F. sacchari*.

Asian *Fusarium* spp. in the FFSC are characterised by mono- and polyphialides producing oval to ellipsoid, rarely pyriform to globose (i.e., *F. annulatum*, *F. fujikuroi* and *F. globosum*) microconidia organized in chains or false heads; 3–5-septate sporodochial conidia and lacking chlamydospores. The elaborate, profusely branched aerial conidiophores of *F. chuoi* are comparable to those of *F. concentricum*, *F. lumajangense*, *F. mangiferae* and *F. sacchari*, all the latter species producing oval, ellipsoidal to allantoid microconidia on false heads. Aerial conidiophores of *F. chuoi*, however, mostly produce macroconidia, while microconidia grouped on false heads are restricted to short, mostly unbranched and prostrate conidiophores formed on the surface on the culture media.

Several Asian species of the FFSC have been reported from *Musa* spp. i.e., *F. annulatum*, *F. concentricum*, *F. fujikuroi*, *F. lumajangense* and *F. sacchari* (Leslie & Summerell 2006, Maryani et al. 2019, Farr & Rossman 2021). The two strains representing *F. chuoi* were isolated as endophytes from asymptomatic seeds of wild banana (*Musa itinerans*), which had been collected predispersal and stored in the Millennium Seed Bank for ~2.5 years at -20 °C prior to isolation.

Supplementary material


FP1353 Phylogenetic tree.

Rowena Hill & Ester Gaya, Comparative Fungal Biology, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3DS, UK; e-mail: R.Hill@kew.org & EGaya@kew.org

Dang Toan Vu, Research Planning and International Cooperation Department, Plant Resources Center, An Khanh, Hoai Duc, Hanoi 152900, Vietnam; e-mail: toanvd.prc@mard.gov.vn

Marcelo Sandoval-Denis & Pedro W. Crous, Westerdijk Fungal Biodiversity Institute, Phytopathology, P.O. Box 85167, 3508 AD, Utrecht, The Netherlands; e-mail: m.sandoval@wi.knaw.nl & p.crous@wi.knaw.nl

Fusarium subflagellisporum



Fungal Planet 1354 – 24 December 2021

Fusarium subflagellisporum T.F. Nóbrega & R.W. Barreto, *sp. nov.*

Etymology. Name refers to the short whip-like projections on apical cells of the sporodochial conidia.

Classification — *Nectriaceae*, *Hypocreales*, *Sordariomycetes*.

Mycelium branched, septate, hyaline, smooth, 2–5 µm diam. **Aerial conidiophores** branched or simple and reduced to conidiogenous cells. **Sporodochia** not observed on carnation leaf agar (CLA), occasionally formed on the surface of synthetic nutrient poor agar (SNA) after 14 d in the dark and potato dextrose agar (PDA) after 7 d under a 12 h daily light regime, both at 25 °C. **Conidiogenous cells** formed on aerial mycelium or in sporodochia, monophialidic, cylindrical to subcylindrical, conical or doliiform, 8–12 × 2–4 µm. **Aerial macroconidia** formed in SNA, abundant, formed directly on phialides, 1–5-septate, septa distinct, fusiform, mostly straight to slightly curved, 18–39 × 2–6 µm, apical cell curved or conical, basal cell foot-shaped or rounded at base, hyaline, thin-walled, smooth. **Sporodochial macroconidia** (on SNA), very abundant orange masses, 4–5-septate, elongate, bearing short whip-like projections, curved dorsiventrally, tapering towards both ends, 37–74 × 3–5 µm, apical cell, generally elongate basal cell either typical foot-shaped or, sometimes subconical, hyaline, thin-walled, smooth. **Microconidia** very rare, formed on phialides of aerial mycelium in SNA, 0–1-septate, ovoid to slightly fusoid, 4–11.5 × 1–4 µm, hyaline, thin-walled, smooth. **Chlamydospores** abundant, terminal or intercalary, either single or in chains, globose, 8–15 µm, aseptate, firstly subhyaline, becoming brown at maturity, smooth or rugose.

Culture characteristics — Fast-growing (up to 85 mm diam in 7 d on PDA under a 12 h daily light regime at 22 °C); raised, margin entire, aerial mycelium dense, felty, vinaceous centrally with alternating zonate rings white and rose towards the edge, reverse vinaceous; abundant sporulation.


Typus. BRAZIL, Pernambuco, Petrolina, Special Fruit Farm (S9°19'17.4" W40°40'29.9"), isolated from hypertrophied floral and vegetative branches of mango trees, 30 Aug. 2019, T.F. Nóbrega (holotype VIC 47377, ex-type culture COAD 2989, *tef1-α* and *rpb2* sequences GenBank MT774486 and MZ970426, MycoBank MB 841204).

Notes — *Fusarium subflagellisporum* is a new member from the *Brachygibbosum* clade (*Fusarium brachygibbosum*). It belongs to the *Fusarium sambucinum* species complex as indicated by *tef1-α* and *rpb2* sequence analyses. Isolates MRC 2486 (from peanuts, USA), FRC R-7642 (from sorghum debris, Puerto Rico), FRC R-8881 (from Nigeria), FRC R-9025 (from millet debris on soil, Zimbabwe), and NRRL 66930 (from corn-cultivated soil, USA) which have been reported in another

study (O'Donnell et al. 2018, Laraba et al. 2021) all grouped in the same clade with *F. subflagellisporum* and are regarded here as belonging to the same species. However, no descriptions of morphology were provided for such isolates and no species names were proposed. Instead, only temporary codes were given to those isolates, listed as *Fusarium* sp. nov. 27 in Laraba et al. (2021) and *Fusarium* sp. nov. 10 in O'Donnell et al. (2018). Information on those isolates clearly indicates that *F. subflagellisporum* is widely distributed and occurs on a broad range of substrates. Laraba et al. (2021) also found that all isolates of *F. subflagellisporum*, listed as *Fusarium* sp. nov. 27, evaluated in their study produced the mycotoxin neosolaniol. It is not known whether the Brazilian isolate of *F. subflagellisporum* from mango also produces neosolaniol.

Fusarium subflagellisporum is phylogenetically close to *F. brachygibbosum* and *F. transvaalense*. Nevertheless, those two species do not produce sporodochial macroconidia with short whip-like projections on apical cells such as in *F. subflagellisporum*.

Fusarium subflagellisporum was isolated from a mango sample (stem and inflorescence) bearing typical symptoms of mango malformation disease. Isolation from those samples yielded *Fusarium* colonies belonging to two morphotypes. One morphotype was identified as *F. sterilihyphosum* – a well-known etiological agent of mango malformation that belongs to the *Fusarium fujikuroi* species complex (Lima et al. 2009). The other was described here as *F. subflagellisporum*. Inoculation on healthy young mango plants with *F. sterilihyphosum* led to development of mango malformation symptoms whereas inoculations with isolate COAD 2989 did not result in any disease symptoms.



Bayesian Inference tree constructed from the combined datasets of *tef1-α* and *rpb2* sequences from species belonging to the *Fusarium sambucinum* species complex, including one isolate of *Fusarium subflagellisporum* sp. nov. obtained in this study (indicated in **bold**). Bayesian posterior probabilities (≥ 0.90) and bootstrap support values ($\geq 70\%$) are given at the nodes. The tree is rooted to *Fusarium fujikuroi* NRRL 66452.



Fungal Planet 1355 – 24 December 2021

***Genea zamorana* Cabero, P. Alvarado & B. Martín, sp. nov.**

Etymology. The epithet refers to the province where the holotype was found (Zamora, Spain).

Classification — *Pyronemataceae*, *Pezizales*, *Ascomycota*.

Ascocarps subglobose, broadly lobed, somewhat gibbous; dark brown in colour, darkened with age; surface covered with high warts (0.9–1.1 mm), more or less distant between them, irregular or polygonal (4–6 sides), with a flat top, often cracked, producing an overall rough matt look to the ascocarp; ascocarps measure 1.3–1.6 cm diam; apical orifice conspicuous, irregularly cylindrical; small mycelium tuft tightly attached to the base. **Odour** not remarkable. **Inner chamber** not divided; with sinuose foldings of the wall containing whitish hymenium layers; chamber walls about 2 mm wide, covered by a brownish epithecium formed by minute scale-like warts. **Peridium** formed by two layers: 1) an external pseudoparenchymatic layer about 350 µm thick, composed of subglobose to angular hyaline elements, 25–30 × 35–40 µm mostly arranged perpendicularly to the peridium surface, which is dark brown; the innermost elements (in contact with the next layer) display a conspicuous reddish colour not observed in other species; and 2) an inner prosenchymatic layer about 50 µm thick that fuses with the hymenium. **Hymenium** arranged as a palisade, composed of asci and cylindrical or filiform septate paraphyses, formed of elements, 35.5–46 × 3.5–4.5 µm, fusing in a epithecium made of subglobose to angular cells. **Asci** cylindrical, indehiscent, inamyloid, stalked, 215–230 × 20–22 µm, when turgid, collapsing soon; stalk central, straight, lacking a crozier at the base, measuring about 30 × 12 µm (length × width). **Ascospores** uniseriate, 6–8 per ascus, facing different directions, subglobose to ovoid, measuring (21.5–)23–25.5(–26) × 18.5–20(–20.5) µm (ornamentation excluded); ornamented with tronco-conical or almost cubical warts measuring 2.3–2.8 × 2–3 µm (width × height), developing small secondary warts on them that produce an irregularly crown-like look.


Habitat & Distribution — Ascomata in groups of 2–5 in summer (July–September). Found in central Spain, in a slightly sandy siliceous soil with a pH of 5.5, surrounded by a *Quercus pyrenaica* forest with *Pteridium aquilinum* and *Erica* sp. receiving 650–750 mm rain/year. Altitude about 995 m a.s.l.

Typus. SPAIN, Castilla y León, Zamora, Sanabria, Remesal, 995 m a.s.l., in a slightly sandy siliceous soil with a pH of 5.5, surrounded by a *Quercus pyrenaica* forest with *Pteridium aquilinum* and *Erica* sp. receiving 650–750 mm rain/year, 27 Sept. 2020, J. Cabero (holotype AH 49255, ITS, LSU and *tef1* sequences GenBank OK138914, OK138913 and OK143509, isotype JC2020927KY (J. Cabero personal fungarium), MycoBank MB 841174).

Additional materials studied. SPAIN, Castilla y León, Zamora, Sanabria, llanes, 995 m a.s.l., in a slightly sandy siliceous soil with a pH of 5.5, surrounded by a *Quercus pyrenaica* forest with *Pteridium aquilinum* and *Erica* sp. receiving 650–750 mm rain/year, 18 July 2017, J. Cabero (JC20170718); Sanabria, Quintana de Sanabria, 995 m a.s.l., in a slightly sandy siliceous soil with a pH of 5.5, surrounded by a *Quercus pyrenaica* forest with *Pteridium aquilinum* and *Erica* sp. receiving 650–750 mm rain/year, 18 July 2017, J. Cabero (JC20140920).

Notes — This species is phylogenetically nested in the clade of *G. fragrans*, closer related to *G. pseudobalsleyi* (ITS 90 % similar; LSU 97 % similar; *tef1* 98 % similar). This clade includes also other species such as *G. anthracina*, *G. amici*, *G. balsleyi*, *G. compressa* and *G. pseudoverrucosa* (Alvarado et al. 2016, 2018, 2020). It differs from these taxa because of the remarkable warts in its external surface, which are much smaller in the other species. In addition, *G. zamorana* presents reddish tones in the innermost rows of the pseudoparenchymatic peridium layer, a feature not observed in any other species of genus *Genea*.

Phylogenetic tree. A 50 % majority rule 28S rDNA- *tef1* consensus phylogram of the species related to *Genea fragrans* (with *Genea arenaria* as outgroup) obtained using MrBayes from 1 200 sampled trees. Nodes were annotated if they were supported by ≥ 0.95 Bayesian posterior probability (left) or ≥ 70 % maximum likelihood bootstrap proportions (right). Nonsignificant support values are exceptionally represented inside parentheses. Samples newly sequenced in this study are in **bold**.



Colour illustrations. Spain, Castilla y León, Zamora, forest of *Quercus pyrenaica*. Ascomata; peridium; paraphyses; spores under LM (from the holotype). Scale bars = 20 µm.

Julio Cabero, C/ El Sol 6, 49800 Toro, Zamora, Spain; e-mail: fotovideocabero@hotmail.com
 Pablo Alvarado, ALVALAB, C/ Dr. Fernando Bongera, Severo Ochoa bldg. S1.04, 33006 Oviedo, Spain; e-mail: pablo.alvarado@gmail.com
 Berta Martín, Servicio Territorial de Agricultura, Ganadería y Desarrollo Rural de Zamora, C/ Prado Tuerto 17, 49019 Zamora, Spain;
 e-mail: berta.martin@zamoradipu.es

Gymnopus nigrescens



Fungal Planet 1356 – 24 December 2021

Gymnopus nigrescens* Bañares, G. Moreno, P. Alvarado & Antonín, sp. nov.Etymology.* After the blackish colour of dried lamellae.Classification — *Omphalotaceae*, *Agaricales*, *Agaricomycetes*.

Basidiocarps annual, with a central stipe. *Cap* 2–3.5(–4) cm, convex, expanding to convex-subapplanate, sometimes with depressed or subumbilicate centre, light reddish brown, darker at centre and paler towards margin, hygrophanous, translucently striate, rough and sometimes sulcate towards margin, greasy to subviscid when moist. *Gills* rather distant, *L* = 24–30, *l* = 15–25, adnexed to almost free, beige-cream with flesh colour tinge, dark brown to black when dried. *Stem* 40–60 × 2–3 mm, dark brown to blackish towards base, concolourous to pileus in upper part, cylindrical, sometimes compressed, with longitudinal groove, glabrous, scarcely compressed-tomentose at base, also strigose and subradicant on substrate. *Spores* (4.4–)4.7–7.2(–7.5) × 2.2–4.5(–5) µm, av. 5.9 × 3.4 µm, *Qav* = 1.8, *n* = 50, ellipsoid to lacrimoid. *Basidia* 4-spored, 23–26 × 7–8 µm. *Basidioles* very abundant, fusoid to claviform. *Cheilocystidia* 25–56(–72) × 3.5–8.5(–14) µm, claviform, cylindrical, deformed with wide projections, rarely lobed at apex and sometimes capitate. *Pileipellis* with lobate terminal cells or poorly developed Dryophila-structure hyphal elements (Antonin & Noordeloos 2010), sometimes with broad lobes, terminal elements up to 21 µm wide, that weakly turn green in KOH. *Stipitipellis* a cutis of cylindrical elements up to 7 µm wide, that strongly turn green in KOH. *Caulocystidia* 2.5–5(–8) µm wide, abundant, cylindrical, flexuose, obtuse, septate, thick-walled, sometimes with short to elongated (up to 12 µm) obtuse projections. *Clamps* present.

Habitat & Distribution — Growing gregarious on bryophytes (*Scleropodium touretii*) in a dense forest of *Erica arborea*. Presently known only from the Canary Islands (Spain).

Typus. SPAIN, Canary Islands, Tenerife, Chanajiga (T.M. Los Realejos), N28°20'37.42" W16°34'50.00", 1170 m a.s.l., 22 Nov. 2019, Á. Bañares & O. Bermúdez (holotype TFC Mic. 25409, isotype in AH 49188; ITS, LSU and *rpb2* sequences GenBank MZ542560, MZ542560 and MZ546414, MycoBank MB 840636).

Additional materials examined. *Marasmiellus* sp. nov. ad int.: FRANCE, Corsica, Valdo Moltifao, roots of *Alnus glutinosa*, 7 Nov. 2019, G. Moreno, F. Esteve-Raventós, A. Altés, F. Pancorbo & P.A. Moreau (AH 51247, ITS and LSU sequences GenBank MZ542557 and MZ542557). *Gymnopus pubipes*: SPAIN, Guadalajara, Albalate de Zorita, under *Quercus ilex* ssp. *ballota*, 15 Oct. 1999, F. Esteve, M. Villarreal, L. Montoya & V. Bandala (paratypus AH 26931, ITS and LSU sequences GenBank MZ542558 and MZ542558). *Gymnopus fuscopurpureus*: CZECH REPUBLIC, Moravian Karst, Adamov, Josefůvské údolí valley, 13 Sept. 2018, H. Ševčíková (BRNM 809119, ITS, LSU and *rpb2* sequences GenBank MZ542559, MZ542559 and MZ546413).

Colour illustrations. Spain, Canary Islands, Tenerife, Chanajiga, on bryophytes in a dense forest of *Erica arborea*, where the holotype was collected. Basidiocarps in situ; basidiocarps in herbarium; pileipellis with Dryophila-structure and clamps; cheilocystidia; caulocystidia; spores (from the holotype). Scale bars = 1 cm (basidiomata), 0.5 cm (basidiomata in herbarium), 10 µm (pileipellis, cheilocystidia and spores under LM), 10 µm (caulocystidia, left 20 µm), 10 µm (spores).

Notes — *Gymnopus nigrescens* is characterised by its light reddish brown pileus, flesh coloured lamellae that turn blackish when dried, poorly-developed Dryophila-structure in the pileipellis, and walls of all hyphae turning green in alkali.

Based on these characters, *G. nigrescens* belongs to sect. *Levipedes*, subsect. *Alkalivirentes* (Antonin & Noordeloos 2010). Other related species with fleshy coloured lamellae are *Gymnopus fuscopurpureus*, which differs by its darker pileus, lamellae not blackening, densely tomentose stipe and differently shaped caulocystidia, and *G. obscuroides* with a long stipe (to 9 cm) and a light brown reduced pileus of 0.9–1.8 cm wide. The sequences of *G. fuscopurpureus* obtained in the present work from specimen BRNM 809119 were significantly similar to those of *G. nigrescens*, but displayed a considerable distance in most markers (ITS: 93 % similar mainly due to a single 30 bp insertion, LSU: 99.5 % similar, *rpb2*: 95 % similar). In turn, the ITS sequences of *G. obscuroides* available in public databases (GenBank KX958398 and KX958399) were identical to others identified as *G. alkalivirens* (GenBank GU234141 and GU234022), which nests inside the genus *Marasmiellus*. Other similar genetic sequences in public databases were obtained from specimens identified as *Collybia brunneola* (GenBank MH874036, 99.32 % similar to LSU of *G. nigrescens*) and *Gymnopus spongiosus* (GenBank KY026706, also 99.32 % similar to LSU of *G. nigrescens*). *Collybia brunneola* is considered an independent species related to *G. dryophilus* (Vilgalys & Miller 1983), so the exact identity of the sample identified with this name needs to be confirmed. *Gymnopus spongiosus* is an American species with white to cream-white or creamy-buff lamellae, a stipe surface subglabrous at apex, and tomentose or strigose towards the base, slightly larger basidiospores (6.2–8.4 × 3.5–4.2 µm), inconspicuous, often collapsed, 24.5–50 µm long, cylindric to strangulated or irregularly lobed and knobbed cheilocystidia, a pileipellis composed of branched hyphae, often bifurcate, not diverticulate or coralloid, coarsely encrusted with a yellow brown pigment, caulocystidia cylindric to contorted, branched, septate (Halling 1996). According to Mata et al. (2006) it is similar to *G. erythropus*.


Supplementary material**FP1356** Phylogenetic tree.

Angel Bañares, Departamento de Botánica, Ecología y Fisiología Vegetal, Universidad de La Laguna, Apdo. 456, E-38200 La Laguna, Tenerife, Islas Canarias; e-mail: angelb@idecnet.com

Gabriel Moreno, Departamento de Ciencias de la Vida (Unidad Docente de Botánica), Facultad de Ciencias, Universidad de Alcalá, E-28805 Alcalá de Henares, Madrid, Spain; e-mail: gabriel.moreno@uah.es

Pablo Alvarado, ALVALAB, Dr. Fernando Bongera st. Severo Ochoa bldg. S1.04, 33006 Oviedo, Spain; e-mail: info@alvalab.es
Vladimír Antonín, Department of Botany, Moravian Museum, Zelný trh 6, 659 37 Brno, Czech Republic; e-mail: vantonin@mzm.cz

Hesperomyces parexochomi



Fungal Planet 1357 – 24 December 2021

Hesperomyces parexochomi* Mironova & Haelew., sp. nov.Etymology.* Referring to the host genus, *Parexochomus*.**Classification** — *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*.

Associated with ladybird hosts in the genus *Parexochomus* (*Coccinellidae*, *Coleoptera*). *Thallus* (251–)299–348–396(–441) μm long from foot to perithecial apex; coloured yellowish. *Cell I* irregularly quadrilateral, broadening distally, $2.3\times$ longer than broad, with very oblique septum I–II and short septum I–III. *Cell II* $2.3\times$ longer than broad, (34–)38–43–48(–53) $\times$ (15–)17–19–21(–24) μm , subtrapezoidal in section. *Cell III* always smaller than cell II, (8.5–)9.3–12.0–14.4(–18.9) $\times$ 10.3–14.3–18.9 μm , with rounded dorsal cell wall. *Primary appendage* consisting of 4–6 superposed cells, (49–)53–59–66(–73) μm long; in the same axis as cell I and III, separated from the latter by the constricted primary septum; its basal cell as long as broad, cup-shaped, larger than each of the remaining cells; second to terminal cells each carrying a single antheridium externally, the terminal cell also carrying a second upwardly directed antheridium. *Antheridia* flask-shaped, with slightly curved efferent necks; the original ascospore apex still present as a short, pointed process between the fourth-to-last and second-to-last antheridium. *Cell VI* $2\times$ longer than broad, (34–)42–58–74(–108) $\times$ (21–)25–29–32(–36) μm , with subparallel margins to broadening distally, septum VI–VII oblique. *Perithecium* (165–)215–253–292(–314) $\times$ (47–)56–63–70(–79) μm , asymmetrical, with the anterior margin convex and the posterior one almost straight; broadest at the second and third tiers, then gradually tapering towards the apex; remainder of the trichogyne apex visible at basis of fifth tier cell; apex a complex with two short lower lobes, two upper (terminal) lobes, and two prominent lips surrounding the ostiole; lower lobes fingertip-shaped; terminal lobes unicellular, elongated, 34–40–44 μm in length, curved upwards and outwardly; ostiole with two lips, one lip triangular, the other slightly shorter, and blunt or rounded. *Ascospores* (59–)63–66–70(–74) μm long, with conspicuous slime sheath only surrounding the larger cell.

Typus. SPAIN, Canary Islands, Las Palmas, Fuerteventura Island, desert near Costa Calma, N28°09'56.2" W14°13'27.4", on *Parexochomus quadriplagiatus* (*Coccinellidae*, *Coleoptera*), 27 Mar. 2017, J. Romanowski, slide D. Haelew. 1465c (holotype at GENT, one subadult and five adult thalli from sternites; isotype DH1465b, three adult thalli from sternites, SSU, ITS and LSU sequences GenBank MZ994889, MZ994868 and MZ994879, MycoBank MB 840989).

Colour illustrations. Xerophilous plants, habitat for *Parexochomus nigripennis*. *Parexochomus nigripennis* with thalli of *Hesperomyces parexochomi* on elytral tips; *P. quadriplagiatus* with thalli of *H. parexochomi* on left elytron; decorative plants at Caleta de Fuste, habitat for *P. quadriplagiatus*; primary appendage, with arrowhead pointing at original spore apex; adult thallus. Scale bars = 200 μm (hosts), 50 μm (thallus), 10 μm (appendage).

Additional material examined. SPAIN, Canary Islands, Las Palmas, Fuerteventura Island, Caleta de Fuste, N28°23'11.4" W13°51'50.9", on *Parexochomus nigripennis* (*Coccinellidae*, *Coleoptera*), 13 Feb. 2017, J. Romanowski, slides D. Haelew. 1377a (GENT, one juvenile and seven adult thalli from left metatrochanter) and 1377b (GENT, nine thalli from sternites); *ibid.*, desert near Costa Calma, N28°09'56.2" W14°13'27.4", on *P. quadriplagiatus*, 27 Mar. 2017, J. Romanowski, slide D. Haelew. 1466a (GENT, one subadult thallus from right epipleuron and one adult thallus from sternite); *ibid.*, Lajares, N28°40'44.1" W13°56'13.5", on *P. nigripennis*, 20 Oct. 2018, J. Romanowski, slide D. Haelew. 1690e (GENT, one adult thallus from right elytron); *ibid.*, slide D. Haelew. 1691e (GENT, two adult thalli from elytral tips); *ibid.*, slides D. Haelew. 1693c (GENT, one juvenile and 11 adult thalli from left elytral tip) and 1693d (GENT, 14 adult thalli from tip of right elytron); *ibid.*, slide D. Haelew. 1694b (GENT, two adult thalli from left antenna).

Additional materials sequenced. See Supplementary material page.

Notes — This species is a part of the near-cryptic *Hesperomyces virescens* species complex (Haelewaters et al. 2018). *Hesperomyces parexochomi* is phylogenetically distinct from its closest relatives. Morphologically, it is very similar to other taxa within *H. virescens* s.lat., except for the position of the original spore apex. It is situated between the fourth-to-last and second-to-last antheridium in *H. parexochomi* whereas it occurs on the last or penultimate antheridium in other members of the complex. Thalli removed from *Parexochomus nigripennis* and *P. quadriplagiatus* are morphologically similar and form a single phylogenetic clade. This is in line with the hypothesis that the host specificity within the *H. virescens* complex exists at the generic level (Haelewaters et al. 2018, Haelewaters & De Kesel 2020).

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence had highest similarity to *H. virescens* ex *Harmonia axyridis* (isolate D. Haelew. 1268d, GenBank MG757830.1; Identities = 789/853 (92 %), 13 gaps (1 %)), *H. virescens* ex *Harmonia axyridis* (isolate D. Haelew. 1268b, GenBank MG757829.1; Identities = 789/853 (92 %), 13 gaps (1 %)), and *H. virescens* ex *Harmonia axyridis* (isolate D. Haelew. 943b, GenBank MZ994889.1; Identities = 789/853 (92 %), 13 gaps (1 %)). Closest hits using the **LSU** sequence are *H. virescens* ex *Adalia bipunctata* (isolate D. Haelew. 1232a, GenBank MG745351.1; Identities = 884/899 (98 %), one gap (0 %)), *H. virescens* ex *Adalia bipunctata* (isolate D. Haelew. 1199h, GenBank MG745347.1; Identities = 884/899 (98 %), one gap (0 %)), *H. virescens* ex *Adalia bipunctata* (isolate D. Haelew. 1231a, GenBank MG745350.1; Identities = 884/900 (98 %), two gaps (0 %)) and *H. virescens* ex *Psyllobora vigintimaculata* (isolate D. Haelew. 1251b, GenBank MG745356.1; Identities = 884/901 (98 %), no gaps).

Supplementary material**FP1357-1** Phylogenetic tree.**FP1357-2** Table. Species of *Hesperomyces* included in this study.

Danny Haelewaters, Research Group Mycology, Department of Biology, Ghent University, K.L. Ledeganckstraat 35, 9000 Ghent, Belgium / Faculty of Science, University of South Bohemia, Branišovská 31, 370 05 České Budějovice, Czech Republic; e-mail: danny.haelewaters@gmail.com

Polina Mironova & Pieter Asselman, Research Group Mycology, Department of Biology, Ghent University, K.L. Ledeganckstraat 35, 9000 Ghent, Belgium; e-mail: polina.mironova@ugent.be & pieter.asselman@ugent.be

Jerzy Romanowski & Piotr Ceryngier, Institute of Biological Sciences, Cardinal Stefan Wyszyński University, Wóycickiego 1/3, 01-938 Warsaw, Poland; e-mail: j.romanowski@uksw.edu.pl & p.ceryngier@uksw.edu.pl



Fungal Planet 1358 – 24 December 2021

Heterophoma rehmanniae A. Shibata, Tonegawa, Sei. Kato, Toy. Sato & Hirooka, *sp. nov.*

Etymology. Named after the host, *Rehmannia glutinosa*, from which our fungus was isolated.


Classification — *Didymellaceae*, *Pleosporales*, *Dothideomycetes*.

Conidiomata on oatmeal agar (OA) pycnidial, solitary or aggregated, sub-globose, glabrous, superficial and immersed into the agar, 71–441 × 55–253 µm in size. **Ostiole** single, slightly papillate or non-papillate. **Mycelium** hyaline, thin-walled, smooth, becoming dark-brown and hypertrophic with age. **Conidiogenous cells** phialidic, hyaline, smooth, globose to lageniform, 5.2–9.3 × 3.7–7.7 µm in size. **Conidia** ellipsoidal to cylindrical, hyaline, thin-walled, smooth, aseptate, (2.9–)3.4–4.2(–4.6) × (1.2–)1.4–1.7(–1.8) µm with small polar guttles. **Chlamydospores** absent.

Culture characteristics — **Colonies** on malt extract agar (MEA) reaching 51–61 mm at 20 °C and 61–67 mm diam at 25 °C after 7 d, margin regular, floccose, (pale) olivaceous grey to dull green; reverse dark olivaceous green near centre. **Colonies** on OA reaching 46–50 mm at 20 °C and 58–60 mm at 25 °C after 7 d, margin regular, woolly, pale olivaceous grey, sometimes black pycnidia formed; reverse concolourous. NaOH test negative.

Typus. JAPAN, Tokyo, Kodaira city, on leaves of *Rehmannia glutinosa* f. *hueichingensis* (*Scrophulariaceae*), Aug. 2015, A. Shibata (holotype TNS-F-93697, culture ex-type HM15-014C = MAFF247505; ITS, LSU, *rpb2* and *tub2* sequences GenBank LC656204, LC656208, LC656201 and LC656212, MycoBank MB 841842).

Phylogenetic tree of the combined LSU, ITS, *rpb2* and *tub2* MAFFT-aligned datasets obtained using Maximum Likelihood. A heuristic search was performed in RAXML-NG (Kozlov et al. 2019) with support at the nodes calculated using bootstrap analyses with 100 replicates. The ML bootstrap support values ≥ 70 % are indicated at the nodes. *Stagonosporopsis hortensis* (CBS 104.42 and CBS 572.85) was used as outgroup. T: ex-type strain; R: representative strain.



Colour illustrations. Flowers of *Rehmannia glutinosa*. Upper row: colony on OA after 7 d at 20 °C. Middle row: colony on MEA after 7 d at 20 °C; conidioma on leaf spot. Bottom row: conidiomata; conidiogenous cell; conidia on OA. Scale bars = 200 µm (conidiomata bottom row), 50 µm (conidioma middle row), 5 µm (conidiogenous cell and conidia).

Additional materials examined. JAPAN, Tokyo, Kodaira city, on leaves of *R. glutinosa* var. *purpurea*, Aug. 2015, A. Shibata (HM15-015 = 15-M0092 = TNS-F-93698, culture HM15-015C = MAFF247506; ITS, LSU, *rpb2* and *tub2* sequences GenBank LC656205, LC656209, LC656202 and LC656213); Toyama, Toyama city, on leaves of *R. glutinosa* f. *hueichingensis*, Sept. 2015, T. Sato (HM15-362 = TKP2 = TNS-F-93699, culture HM15-362C = MAFF 245313; ITS, LSU, *rpb2* and *tub2* sequences GenBank LC656206, LC656210, LC656203 and LC656214); Toyama, Toyama city, on leaves of *R. glutinosa* var. *purpurea*, Sept. 2015, T. Sato (HM15-364 = TAP1 = TNS-F-93700, cultures HM15-364C = MAFF 245316; ITS, LSU and *tub2* sequences GenBank LC656207, LC656211 and LC656215).

Notes — In our Maximum Likelihood tree using LSU, ITS, *rpb2* and *tub2*, our four isolates represented an independent clade in the genus *Heterophoma*. The isolates were different from seven existing species of *Heterophoma*, i.e., *H. nobilis*, *H. poolensis*, *H. adonis*, *H. sylvatica*, *H. verbascicola*, *H. verbasci-densiflori* and *H. novae-verbascicola*, based on shape and size of conidia, and NaOH reaction (see Table in Supplementary material; De Gruyter & Noordeloos 1992, De Gruyter et al. 1993, Chen et al. 2015, 2017, Hou et al. 2020a).

Supplementary material

FP1358 Table. Morphological characteristics of *Heterophoma rehmanniae* and published descriptions of *Heterophoma* spp.



Fungal Planet 1359 – 24 December 2021

***Hyphodermella pallidostraminea* Bukharova & Volobuev, sp. nov.**

Etymology. Name refers to the pale straw-yellow colour of hymenial surface.

Classification — *Phanerochaetaceae*, *Polyporales*, *Agaricomycetes*.

Basidiomata annual, resupinate, adnate, effuse, ceraceous to crustaceous when dry, at first orbicular, then confluent, up to 15 mm diam. Margin white, determinate, narrow, fibrillose, with age thinning out. **Hymenial surface** yellowish, tiger yellow (090 90 50), fresh yellow (095 90 40) or pea green (095 80 40) (all colour codes are given following the RAL Design colour chart), smooth to slightly tuberculate, cracked. Subiculum thin, indistinct, white. **Hyphal system** monomitic; generative hyphae simple-septate. Subicular hyphae 3.5–5 µm diam, hyaline, thick-walled or with slightly thickened walls, encrusted with numerous prismatic crystals, loosely interwoven, more or less parallel arranged, occasionally branched. Subhymenial hyphae 2.5–3.5 µm diam, hyaline, thin-walled, smooth, compactly interwoven, frequently ramified. **Cystidia** absent, but cylindrical to fusoid, cystidioid hyphal ends occasionally present, thin-walled, not encrusted, projecting above the hymenium up to 35 µm. **Basidia** 17–25 × 5.2–6 µm, clavate, with a simple septum at the base, 4-sterigmate. **Basidiospores** (5.0–)5.4–6.6(–6.7) × (2.9–)3.0–3.5(–3.6) µm, n = 30, L = 5.96, W = 3.23, Q = 1.82–1.87, smooth, hyaline, thin-walled, ellipsoid, inamyloid, non-dextrinoid, with tiny oil drops.


Habitat & Distribution — On dry, dead *Actinidia* sp. branch, in a herbaceous mixed coniferous-broadleaf forest. Hitherto only known from the type locality in the south of the Russian Far East.

Typus. RUSSIA, Jewish Autonomous Oblast, Obluchensky District, Bastak State Nature Reserve, 15 km north-northeastward from the Kirga settlement, at the foot of Mount Skalistaya, N49°00'47.3" E132°53'53.6", on dry dead branch of *Actinidia* sp. (*Actinidiaceae*) in mixed coniferous-broadleaf forest, 24 Aug. 2009, N. Bukharova (*Vasilyeva*) (holotype LE 286968, isotype VLA M-22719, ITS and LSU sequences GenBank OK138912 and OK138911, MycoBank MB 841840).

Notes — *Hyphodermella pallidostraminea* is a minute-sized corticioid fungus characterised macroscopically by pale-yellowish smooth to tuberculate hymenophore and white pruinose margin of basidiocarps. The main distinguishing microscopic feature among other representatives of the genus *Hyphodermella* is the spore size (up to 3.5 µm in width) that is overlap-

ping only with the recently described *H. zixishanensis*, but the latter species differs by the reddish to brown hymenial surface and the absence of cystidioid hyphal ends projecting above the hymenium (Wang et al. 2021). The previous attempt to find a taxonomic interpretation for the studied specimen of *H. pallidostraminea* without a molecular analysis led to the fact that this material was erroneously identified as *Sistotremastrum niveocreum* (Bukharova & Zmitrovich 2014), also being a deciduous-dwelling fungus having a monomitic hyphal system, similar spore sizes and shape, but differing by clamps on hyphae.

Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits using the ITS sequence had highest similarity to *H. rosae* (strain FP-150552, GenBank KP134978; Identities = 625/657 (95 %), 13 gaps (1 %)), *H. corrugata* (voucher K(M) 237661, GenBank MK496919; Identities = 633/666 (95 %), 13 gaps (1 %)) and *H. rosae* (voucher CFMR DLL2011-177, GenBank KJ140672; Identities = 613/646 (95 %), 13 gaps (2 %)). Closest hits using the LSU sequence are *Geliporus exilisporus* (voucher TNM GC 1702-15, GenBank LC379153; Identities = 847/859 (99 %), no gaps), *H. rosae* (strain GC 1604-113, GenBank MZ637147; Identities = 849/862 (98 %), no gaps) and *H. corrugata* (voucher MA-Fungi 5527, GenBank JN939597; Identities = 849/862 (98 %), no gaps).



Phylogenetic tree derived from Maximum Likelihood analysis based on nrITS1-5.8S-ITS2 data. The analysis was performed in the IQ-TREE Web Server (Trifinopoulos et al. 2016) with 1000 ultrafast bootstrap replicates. Maximum Likelihood bootstrap support values shown above branches (BS > 80 %). The new species described in this study is in **bold** face. The scale bar represents the expected number of nucleotide changes per site.

Colour illustrations. Coniferous-broadleaf mixed forest in the Bastak State Nature Reserve (Jewish Autonomous Oblast, Russia). Basidiocarp; basidia and spores (all from holotype). Scale bars = 2 mm (basidiocarp), 10 µm (cystidioid hyphal ends), 5 µm (spores and basidia).

Sergey V. Volobuev, Komarov Botanical Institute of the Russian Academy of Sciences, 2, Professora Popova street, 197376 Saint Petersburg, Russia; e-mail: sergvolobuev@mail.ru

Nadezhda V. Bukharova, Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far Eastern Branch of the Russian Academy of Sciences, Pr-t 100-let Vladivostoka 159, 690022 Vladivostok, Russia; e-mail: nadya808080@mail.ru



Fungal Planet 1360 – 24 December 2021

***Inocybe corsica* Esteve-Rav., Pancorbo & G. Moreno, sp. nov.**

Etymology. Name refers to the island of Corsica (Corse), where the samples were collected.

Classification — *Inocybaceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata agaricoid and stipitate. **Pileus** 10–25 mm, at first conical-campanulate, then convex to plano-convex, broadly umbonate to subumbonate, slightly hygrophanous; margin straight, regular to hardly wavy with age, fissurate at times, velipellis not present; colour yellow ochraceous (Mu 10YR 6/8, 7/6) when young or moistened, from uniform to light ochraceous in the centre (Mu 10YR 8/2) in some basidiomata; surface radially fibrillose, smooth, not rimose, slightly striated towards the margin in adult basidiomata, with greasy aspect, not sticky when moist. **Lamellae** moderately crowded ($L = 35\text{--}45$; $I = 1\text{--}2$), free to adnate, ventricose, initially whitish, becoming pale grey to beige, then pale brown, edge concolourous, entire to finely crenulate. **Stipe** 22–40 × 3–4 mm, straight to curved towards base, bulbous to bulbous-napiform, not marginately bulbous; colour whitish, becoming beige to ochraceous with age (Mu 10YR 8/2; 7/3); surface densely pruinose along the entire surface. **Cortina** absent. **Context** fibrose, whitish, concolourous to the stipe surface. **Smell** herbaceous, **taste** not recorded. **Colour of exsiccatum** ochre-brownish (pileus and stipe), not blackening. **Spores** (8.0–)8.7–9.7–10.9(–12.2) × (5.4–)6.7–7.7–8.7(–9.1) µm, Qm: (1.0–)1.09–1.25–1.42(–1.7) ($n = 193 / N = 2$), mostly subsisdiametric to sometimes subheterodiametric, distinctly nodulose (though exceptionally sub-entolomatoid) under the optical microscope, provided with 7–10 distinct knobs variable in height (1.3–2.3 µm high), yellowish, apicula distinct. **Basidia** (24–)27–34.8–41(–49) × (10.9–)11.3–13.2–15.8(–17.1) µm; Qm: (1.81–)2.0–2.6–3.2(–3.4), 4-spored, rarely 2-spored, clavate, sterigmata 2.5–7 µm long. **Lamella edge** practically sterile, composed by numerous protruding hyaline to brownish cheilocystidia mixed with abundant mostly hyaline to brownish clavate paracystidia. **Pleurocystidia** abundant, (38–)45–58.1–72(–80) × (11.3–)12.7–16.5–21.2(–22.4) µm, Qm: (1.78–)2.5–3.6–4.8(–5.6), ($n = 110 / N = 2$), broadly lageniform to rarely fusiform, hyaline or with brownish content, base often attenuate but not pedicellate, rather crystalliferous at the apex, walls (1.0–)1.2–1.6–2.2(–2.4) µm thick, reaching ~3 µm at the apex, hyaline to hardly yellowish in 10% NH_4OH . **Cheilocystidia** numerous, (43–)47–58.2–73(–79) × (11.0–)13.3–16.5–20.3(–22.7) µm, Qm: (2.21–)2.7–3.5–4.8(–7.1), similar in size and shape to pleurocystidia. **Stipitipellis** a cutis of parallel hyphae (3.5–)4.3–6.0–7.9(–10.3) µm, bearing numerous caulocystidia along the entire length of the stipe, (45–)49–63.2–83(–90) × (8.3–)8.8–12.2–15.8(–16.8) µm, Qm: (3.8–)4.0–5.2–7.6(–8.4) ($n = 73 / N = 2$), more slender but similar in shape to hymenial cystidia, mixed with numerous

clavate to subcylindrical hyaline paracystidia. **Pileipellis** a cutis formed by parallel cylindrical cells, 4.5–13 µm wide, somewhat constricted at septa, showing minute pale yellow ochraceous pigment, minutely encrusting and also diffuse intracellular. **Lamellar trama** of parallel hyphae, 6–16 µm broad, showing diffuse yellowish intracellular pigment. **Clamp connections** abundant in all tissues.

Habitat & Distribution — Gregarious in acidic soil; in humid and herbaceous areas, next to willows (*Salix atrocinerea*), in somewhat muddy and nitrified soils; the collecting area is located in a Mediterranean environment of holm oak (*Quercus ilex* subsp. *ilex*) and cork oak (*Quercus suber*) mixed forest. No sequence similar to the ones generated from our material are available in GenBank.

Typus. FRANCE, Corsica, Corti, Saint-Jean, Maison San Giovanni, N42°17'38.04" W9°10'04.61", 427 m a.s.l., along a ditch on mossy and wet ground, near *Salix atrocinerea*, with some *Juncaceae* and *Rosa* sp. scrub, close to *Quercus suber*-*Q. ilex* subsp. *ilex* forest, in acidic sandy soil, 8 Nov. 2019, F. Pancorbo, A. Altés, G. Moreno & F. Esteve-Raventós (holotype AH 51900, ITS and LSU sequences GenBank MZ308644 and MZ308646, MycoBank MB 839085).

Additional material examined. Same location, date, habitat and collectors, AH 55201 (paratypus), ITS and LSU sequences GenBank MZ308645 and MZ308647).

Notes — Colour codes are taken from Munsell (1994), terminology follows Vellinga (1988) and Kuyper (1986). *Inocybe corsica* is probably a species of Mediterranean distribution linked to *Salicaceae* (found under *Salix atrocinerea*), of humid and waterlogged areas in acidic soils; macroscopically it resembles several species of the *Xanthomelas* clade (Esteve-Raventós et al. 2015, Vauras & Larsson 2016, Bandini et al. 2019, 2020) because of its medium-small size, and is characterised by the presence of a non-marginate napiform bulb and flesh that does not blacken markedly with age or desiccation. Our analyses of the ITS region (Supplementary Fig. 1) show that *I. corsica* nests in the well supported *I. sect. Marginatae* clade (ML-BS 98 %, BPP 1), within which it forms a clade together with a sequence of *I. diabolica* (ML-BS 95 %, BPP 0.98). The two sequences of *I. corsica* obtained are identical and form a fully-supported clade (ML-BS 100 %, BPP 1). From the group of species linked to the *Salicaceae*, both *I. salicis* (= *I. straminipes*) and *I. lacunarum* (Vauras & Larsson 2016), show different spores, provided with numerous (17–22) prominent knobs (Kühner 1955); another species bound to *Salix* and *Populus* and discovered in Japan is *I. populea*, very close to *I. salicis* in morphological features but showing an indistinctly marginate to claviform stipe base and smaller cystidia (Kobayashi & Courtecuisse 2000).

(text continues on Supplementary material page FP1360)

Colour illustrations. France, Corsica, Corti, Saint-Jean, Maison San Giovanni, *Salix atrocinerea* with *Quercus suber* in muddy and mossy soil, near a dirt road, locality where the samples were collected. Basidiomata; pleurocystidia; cheilocystidia; caulocystidia at stipe base; spores under optical microscope; spores under SEM (bottom right). Scale bars = 50 µm (cystidia), 10 µm (spores), 2 µm (spores under SEM).

Supplementary material

FP1360 Phylogenetic tree.

Fernando Esteve-Raventós, Gabriel Moreno & Alberto Altés, Universidad de Alcalá, Facultad de Ciencias, Departamento de Ciencias de la Vida (Botánica), 28805 Alcalá de Henares, Madrid, Spain; e-mail: fernando.esteve@uah.es, gabriel.moreno@uah.es & alberto.altes@uah.es

Fermín Pancorbo, Sociedad Micológica de Madrid, Real Jardín Botánico. C/ Claudio Moyano 1, 28014 Madrid, Spain; e-mail: fermin@socmicolmadrid.org

Ibai Olariaga, Rey Juan Carlos University, Dep. Biology and Geology, Physics and Inorganic Chemistry, C/ Tulipán s/n, 28933 Móstoles, Madrid, Spain; e-mail: ibai.olariaga@urjc.es

Inocybe nivea



Fungal Planet 1361 – 24 December 2021

Inocybe nivea* E. Larss., sp. nov.Etymology.* Refers to the white colour of the mushroom.Classification — *Inocybaceae*, *Agaricales*, *Agaricomycetes*.

Pileus 5–20 mm diam, as young campanulate, conical to convex mostly with an obtuse to broad umbo, sometimes more prominent, margin incurved, later becoming convex to plano-convex with broad or a more prominent umbo, initially pure white, sericeous smooth, silky, with age becoming pale buff to isabelline, when affected by harsh weather cracking and becoming scaly, velipellis indistinct. *Cortina* white visible and rather abundant in young basidiomata. *Lamellae* moderately crowded, interspersed with lamellulae, L = 25–40, 2–3 mm broad, ventricose, sinuate to emarginate, first pale beige with a greyish tone, later yellowish brown, edge fimbriate white. *Stipe* 15–35 × 1–4 mm, equal to slightly bulbous but without bulb, solid, whitish, at apex pruinose descending to about 1/3, downwards smooth, in lower part fibrillose. *Context* whitish to pale buff. *Smell* spermatic, *taste* indistinct. *Basidiospores* (8.5–)9.6–9.7–9.8(–11.1) × (4.6–)5.7–5.8–5.9(–6.5) µm, n = 80, Q = (1.5–)1.62–1.65–1.66(–1.9), smooth ellipsoid, amygdaliform to subamygdaliform with obtuse apex and a small distinct apiculus, yellowish brown. *Basidia* 30–35 × 9–11 µm, n = 20, subclavate to clavate, 4-spored, hyaline, sterigmata 4.5–5.8 µm. *Pleurocystidia* 45–65 × 11–18 µm, n = 50, fusiform, lageniform to utriform with a slender pedicel, thick walled, 1–3 µm thick, hyaline, crystalliferous at apex. *Cheilocystidia* similar to pleurocystidia but shorter, 38–52 × 13–18 µm, n = 20, lageniform to broadly utriform, densely arranged, thick walled, 2.5–4.5 µm thick. *Paracystidia* hyaline, pyriform to clavate 15–28 × 10–13 µm, n = 10. *Caulocystidia* at apex similar to pleurocystidia, abundant, with crystals, less so further down 45–65 × 11–18 µm, n = 10, fusiform to more cylindrical, caulocystidioid hairs thin-walled, sometimes septate, 30–60 × 5–12 µm, cauloparacystidia few. *Pileipellis* an interwoven cutis of smooth hyaline hyphae, 6–12 µm wide. *Clamp connections* present.

Ecology & Distribution — Associated with *Salix polaris* and *Salix* spp. on nutrient rich and somewhat calcareous soils. Basidiomata so far only known from an arctic habitat on Svalbard where it grew in a rather moist area among mosses and herbs associated with *S. polaris*, and from a subalpine *Betula pubescens* forest in Norway, where it grew in a moist area associated with *Salix* spp. Blast search of NCBI's GenBank nucleotide database and the UNITE database gave no further information on distribution and occurrence as no matches of additional ITS sequence data were available.


Colour illustrations. *Inocybe nivea* habitat in the arctic zone growing associated with *Salix polaris*, on Svalbard, Spitsbergen, Revneset. *In situ* basidiomata of the holotype (GB-0207613); hymenial pleurocystidia; caulocystidia from stipe apex; basidiospores. Scale bars = 20 µm (pleuro- and caulo-cystidia), 10 µm (spores).

Typus. SVALBARD AND JAN MAYEN, Spitsbergen, Revneset, arctic habitat with mosses and herbs, associated with *Salix polaris*, 12 Aug. 2015, E. Larsson, EL76-15 (holotype GB-0207613, isotype O, ITS-LSU sequence GenBank OK090776, MycoBank MB 841189).

Additional materials examined. SVALBARD AND JAN MAYEN, Spitsbergen, Revneset, arctic habitat with mosses and herbs, associated with *Salix polaris*, 12 Aug. 2015, E. Larsson, EL81-15, GB-0207614. — NORWAY, Nordland, Saltdal, Jukerdalsura NR, subalpine area on gravelly ground with *Betula pubescens*, *Alnus incana*, and *Salix* spp., 10 Aug. 2012, E. Larsson, EL43-12, GB-0207612 (ITS-LSU sequence GenBank OK090775).

Notes — *Inocybe nivea* belongs in the *I. geophylla* group, a group that is identified to host a high phylogenetic diversity and the name *I. geophylla* has been widely applied to many taxa in North America and Europe (Ryberg et al. 2008, Matheny & Swenie 2018), and in addition *I. geophylla* is reported associated with a large number of frondose and coniferous trees (Kuyper 1986). In macro- and micro-morphology they all are rather similar, but the geographic distribution and ecological preferences may explain part of the observed phylogenetic diversity. Further work on the group is needed.

Inocybe nivea is a rather small species within the group, characterised by a conical to plano-convex white pileus, about 15–20 mm diam in mature basidiomata, and seems to be associated with *Salix* spp. The pileus is silky and smooth but can in the arctic zone often be affected by cold and windy weather conditions that makes the cutis break up and form large scales. This was not observed in the subalpine basidiomata. In *I. geophylla* the pileus is usually much larger and they have different host preferences and ecology.



Phylogram obtained using PAUP v. 4.0a (Swofford 2003) based on ITS and LSU data showing the position of *I. nivea* in the *I. geophylla* group. Heuristic searches with 1000 random-addition sequence replicates and tree bisection-reconnection (TBR) branch swapping were performed. Relative robustness of clades was assessed by the bootstrap method using 1000 heuristic search replicates with 100 random taxon addition sequence replicates and TBR branch swapping. Bootstrap support values are indicated on branches. *Inocybe nivea* is marked in **bold** and the holotype is indicated.



Fungal Planet 1362 – 24 December 2021

***Laccaria diospyricola* Salna Nanu & T.K.A. Kumar, sp. nov.**

Etymology. Basidiomata of the species were always found among roots of *Diospyros paniculata*.

Classification — *Hydnangiaceae*, *Agaricales*, *Agaricomycetes*.

Pileus 5–15 mm diam, slightly depressed when young, becoming concave to infundibuliform at maturity; surface flesh coloured (Methuen 6B5–6B6, 7A2–7A4), cream (6A2–6A3) towards the centre, with occasional scattered violaceous spots, hygrophanous, minutely pubescent throughout, and covered with small superficial patches of whitish mycelium; margin uplifted at maturity. **Lamellae** adnate to slightly decurrent, concolourous with the pileus (6B5–6B6), thick, distant, up to 1.5 mm wide, with lamellulae in 2–3 tiers; edge serrate, concolourous with the sides. **Stipe** 10–30 × 1–3 mm, central, terete, almost equal, slightly tapering towards the base, flesh coloured (6B5–6B6) without violaceous spots, pubescent; base with mycelial pad. **Basidiospores** (two observations two collections, five sporocarps, 20 basidiospores) 7–9 × 6–8 µm (excluding the spines), Q = 1.0–1.28, Qm = 1.15, subglobose to globose, echinulate, thin-walled, hyaline, inamyloid; echinulae up to 1 µm long. **Basidia** 23–66 × 9–15 µm, clavate, thin-walled, 4-spored. Sterigmata up to 11 µm long. **Cheilocystidia** 40–90 × 4–10 µm, flexuose, cylindrical to narrowly clavate, hyaline, thin-walled, inamyloid. Lamellar trama subregular; hyphae 3–6 µm wide, hyaline, inamyloid, slightly thick-walled (up to 1 µm thick). **Pileipellis** an agglutinated cutis, occasionally disrupted by trichodermal patches of ascending or erect hyphae; hyphae 2–7 µm wide, thin-walled. **Stipitipellis** a cutis frequently disrupted by clustered or scattered caulocystidia; hyphae 2–6 µm wide, inamyloid, slightly thick-walled (up to 1 µm thick) with pale yellow content. **Caulocystidia** 38–185 × 8–16 µm, flexuose, inamyloid, hyaline, slightly thick-walled (up to 1 µm thick). **Stipe context** composed of 3–10 µm wide hyphae which are hyaline, inamyloid, slightly thick-walled (up to 1 µm thick), with yellowish brown plasmatic pigment. **Clamp connections** observed on all hyphae.

Habitat & Distribution — Scattered or in small groups, on sandy soil, in the root bed of *Diospyros paniculata*. Basidiocarps collected from sacred groves of two different regions of Kerala State, India.

Typus. INDIA, Kerala State, Kozhikode District, Thurayilkavu, on the ground under *Diospyros paniculata* (*Ebenaceae*), 10 Aug. 2018, S. Nanu (holotype CAL1771, nrITS sequence GenBank MK776767, MycoBank MB 833411).

Additional material examined. INDIA, Kerala State, Kannur District, Poon-gottukavu, 12 Sept. 2018, S. Nanu (CAL1770).

Colour illustrations. India, Kerala State, Kozhikode district, Thurayilkavu, a sacred grove with different plants including *Diospyros paniculata*. SEM photograph of basidiospores; cheilocystidia; caulocystidia; basidiomata. Scale bars = 10 mm (basidiomes), 10 µm (all others).

Notes — Small basidiomata, infunduliformis to applanate pileus, absence of two-spored basidia, cutis type of pileipellis with trichodermal patches and presence of caulocystidia are the diagnostic features of *L. diospyricola*. This species is closely related to *L. violaceotincta* and *L. vinaceoavellanea*. *Laccaria diospyricola* is found near or on the root bed of *D. paniculata*, whereas, *L. violaceotincta* was reported from a freshwater swamp forest dominated with *Myristica* species (Latha et al. 2019). *Laccaria diospyricola* differs from *L. violaceotincta* by having small basidiomata, absence of 2-spored basidia, and a cutis type of pileipellis disrupted by trichodermal patches. Hongo (1971) originally described *L. vinaceoavellanea* from Japan, having brownish to vinaceous basidiomata, greyish red to white pileus, distant and decurrent gills, tough, solid and striate stipe, globose and inamyloid spores with ornamentation, and presence of clamp connections in all hyphae. *Laccaria diospyricola* basidiomata show some resemblance with *L. vinaceoavellanea* in having globose to subglobose basidiospores with spines, and in having clamp connections in all hyphae. Different colour of basidiomata, absence of tough, solid and equal stipe, and molecular data are the characters that distinguish *L. diospyricola* from *L. vinaceoavellanea*. *Laccaria yunnanensis* also shows similarities with *L. diospyricola*. However, the presence of adnate to slightly decurrent lamellae, densely pubescent and terete stipe, larger cheilocystidia, and caulocystidia separates *L. diospyricola* from *L. yunnanensis*.

A BLAST search in GenBank nucleotide database using the newly generated ITS sequence of *L. diospyricola*, resulted in a close hit with *Laccaria violaceotincta* (GenBank MK141034), showing a sequence similarity of 94.26 %. The second closest hit (93.74 %) in the BLAST search was with *L. vinaceoavellanea* (GenBank LC098741). Phylogenetic analyses using maximum parsimony (MP) and maximum likelihood (ML) methods generated trees that were congruent with regards to the topology and the relative positions of the taxa. The bootstrap support values from the MP analysis were similar to ML support values. In both the MP and ML trees, *Laccaria diospyricola* settled in a separate well-supported clade along with other tropical species, *L. violaceotincta*, *L. yunnanensis*, and *L. vinaceoavellanea*. *Laccaria diospyricola* resolved in a branch distinct from all other species, and the clade received a bootstrap support value of 99 % in the ML analysis and MP analysis.

Supplementary material**FP1362** Phylogenetic tree.


Lasiosphaeria deviata



Lasiosphaeria deviata A.N. Mill. & Læssøe, *sp. nov.*

Colour illustrations. Cattle-grazed old meadow partly overgrown with *Crataegus*. Ascomata; ascomal neck; longitudinal section through ascomal wall; asci; ascospores and phialides. Photo credits: Nikolaj Gervig Correll (habitat), Andrew N. Miller & Thomas Læssøe. Scale bars = 500 µm (ascomata), 10 µm (all others).

Notes — *Lasiosphaeria devata* possesses the typical tomentose ascomata with a 3-layered ascomal wall known for the genus but lacks yellow centrum pigments (Miller & Huhndorf 2004a, b). It can be distinguished from all other species of *Lasiosphaeria* by its aberrant arnium-like ascospores, which are dark brown, ellipsoid and possess hyaline, lash-like appendages at each end of the ascospore. *Arnium* has been shown to be highly polyphyletic (Kruys et al. 2015). Other tomentose species of *Arnium* include *A. olerum*, *A. sulphureum* and *A. tomentosum* (Lundquist 1972). *Arnium olerum* possesses a whitish tomentum, whereas *A. tomentosum* possesses a brown tomentum. *Arnium sulphureum* possesses a yellow tomentum, but only around the ascomal neck, is semi-immersed, has larger ascospores and grows on horse dung (Muroi et al. 1987). Neither the holotype at NHL nor the isotype at TNS of *A. sulphureum* can be found, so this name should be considered a *nomen dubium*. *Lasiosphaeria devata* occurs as a sister taxon to *Cercophora gossypina*, which has a white tomentum and typical cercophora-like ascospores.



Maximum likelihood tree generated from a PhyML analysis of a combined ITS-LSU-*Btub* sequence alignment in Seaview v. 4.5.4 (Gouy et al. 2010). Missing loci are shown as dashes. *Lasiosphaeria deviata* is in **bold**. Bootstrap branch support $\geq 75\%$ are shown above or below nodes and thickened branches indicate Bayesian posterior probabilities $\geq 95\%$. GenBank accession numbers for the combined ITS-LSU-*Btub* alignment are given after taxon names.

Lepiota sayanensis



Fungal Planet 1364 – 24 December 2021

***Lepiota sayanensis* E.F. Malysheva & Malysheva, sp. nov.**

Etymology. The specific epithet refers to the geographic name of type locality, the Sayan Mountains in southern Siberia.

Classification — *Verrucosporaceae*, *Agaricales*, *Agaricomycetes*.

Basidiocarps small. **Pileus** 10–15 mm diam, convex, plano-convex with low prominent umbo, beige (RAL 1001) (the RAL colour standard – K7 colour fan, edition 2019, <https://www.ral-farben.de/en/ral-k7-colour-fan>, was used for basidiocarps description) or sand yellow (RAL 1002), around centre beige red (RAL 3012) or orange-brown (RAL 8023), smooth and glabrous, with contrasting cream (RAL 9001) or whitish, felted to fringed margin. **Lamellae** crowded, free, ventricose, whitish at first and soon cream coloured (RAL 9001), with concolourous even edge. **Stipe** 30–40 × 1–3 mm, cylindrical or slightly wider at base, cream (RAL 9001) or sand yellow (RAL 1002), ochre brown (RAL 8001) or beige brown (RAL 8024) at utmost base, smooth, hollow, with white tomentum at base. **Annulus** whitish and distinct, funnel-shaped, persistent, with flaky rim. **Context** thin, cream or white. **Odour** indistinct. **Basidiospores** 4.1–5.2 × (2.0–)2.7–3.4 µm, (n = 60, Q = 1.3–1.7, Qav = 1.6), ellipsoid to oblong, slightly thick-walled, smooth, without germ pore, non-dextrinoid. **Basidia** 16–20 × 4–6 µm, 2- and 4-spored, with clamp connection. Lamella edge fertile. **Cheilocystidia** absent. **Pleurocystidia** absent. **Pileipellis** hymeniderm, made up of cells, 24–45 × 10–22 µm, varying in shape: narrowly to broadly clavate, broadly utriform, or sphaeropedunculate, with short to long pedicel, thick-walled, with pale yellowish intracellular pigment. **Clamp connections** present, numerous.

Habitat & Distribution — Growing solitary on litter and soil. So far known only from type locality.


Typus. RUSSIA, Krasnoyarsk Territory, Sayano-Shushensky Biosphere Reserve, vicinity of Kerema field station, the mouth of the Bolshaya Kerema River, N52°07'07.6" E92°13'35.8", mixed forest (*Betula pendula* with *Pinus sylvestris* and *Larix sibirica*), on litter, 23 Aug. 2015, E. Malysheva (holotype LE 312933, ITS and LSU sequences GenBank MZ997391 and MZ997392, MycoBank MB 841097).

Notes — *Lepiota sayanensis* is characterised by its small basidiocarps, pale coloured pileus with white, fringed margin, persistent and funnel-shaped annulus, ellipsoid to oblong, non-dextrinoid basidiospores, fertile lamellae edge without cheilo- and pleurocystidia, and hymeniderm pileipellis with tightly packed elements.

This new species is very similar and morphologically almost indistinguishable from *Lepiota neophana* (including var. *europaea* and f. *papillata*). The only differences of our species from the latter are a smooth, non-wrinkled pileus, persistent and funnel-shaped annulus on the stem, which is flimsy and evanescent in *L. neophana*, and the presence of 2-spored basidia. According to E.C. Vellinga (Vellinga 2010), *L. neophana* most

likely represents a species complex and may combine several species that differ phylogenetically and in their geographic distribution, and our species is very close to this complex (see phylogenetic tree below). The ITS sequences from *L. neophana* show a clear genetic distance from *L. sayanensis* that reaches 4 %.

Morphologically close species, *Lepiota psalion*, with similar small and pale-coloured basidiocarps, differs in shorter basidiospores (3.0–4.5 × 2.5–3.5 µm; Vellinga 2001, Vizzini et al. 2019) and the presence of cheilocystidia. The new species differ similarly from *L. coloratipes* (*Lepiota rufipes* f. *phaeophylla*), which has cheilocystidia, shorter basidiospores (3.0–3.6(–4) × 2.0–2.8 µm) and evanescent annulus on stipe (Vizzini et al. 2014). Another morphologically close species, *Lepiota recondita*, can be distinguished by an incomplete, often disappearing with age, annulus, and the presence of cheilocystidia (Vizzini et al. 2019).



Maximum likelihood phylogeny from a nrITS alignment performed on RAXML server v. 1.0.0 (<https://raxml-ng.vital-it.ch/#/>) (Kozlov et al. 2019) with 100 rapid bootstrap (BS) replicates, under the GTR+G model, showing relationships of some *Lepiota* species with a hymeniform pileipellis, with *Chamaemyces fracidus* as outgroup. A Bayesian analysis was performed with MrBayes v. 3.2.5 (Ronquist et al. 2012) with 7 M generations under the same model. Support values (BS/PP) are given above the branches. All tips are labelled with taxon name and GenBank accession number. The newly generated sequence is in **bold**.

Colour illustrations. Russia, bank of Yenisei River in Sayano-Shushenskiy Reserve. Mature basidiocarp; pileipellis elements; basidiospores (all from holotype). Scale bar = 0.5 cm (basidiocarp), 10 µm (all others).



Fungal Planet 1365 – 24 December 2021

***Micropsalliota albofelina* D. D. Ivanova & O.V. Morozova, sp. nov.**

Etymology. From Latin '*albus*' (white) and '*felis*' (cat), referring to the surface of the basidiomata, covered with white hairs that resemble white feline fur.

Classification — *Agaricaceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata delicate and tiny, pristine white when fresh, discolouring to brown, brown in exsiccata, reddish brown (6C5, 6D5–6; Kernerup & Wanscher 1978) when bruised or cut. **Pileus** 4–5 mm diam, convex, surface hirsute due to presence of thin hairs, pure white. **Context** less than 1 mm. **Lamellae** free, with 1–2 series of lamellulae, 0.5–0.7 mm broad, white as the whole basidiomata. **Annulus** absent. **Stipe** 17–29 × 1.0–1.5 mm, cylindrical, similarly thick across the whole length, concolourous with the pileus, surface hirsute. **Odour** faint, **taste** not reported. **Basidiospores** 5.5–7.5 × 3.5–4 µm, Q = 1.4–2, Qav = 1.8 (n = 21) ellipsoid, with apical thickening. **Basidia** 12.5–17.5 × 6.5–8 µm, 4-spored, clavate, hyaline, smooth (in the light microscope). **Cheilocystidia** 33–34.5 × 10–13 µm, irregularly lecythiform, capitate, with a long, narrow and fragile neck (3–4 µm diam) and thickened base (10–13 µm diam), capitulum 4–6 µm diam, pigment absent. **Pleurocystidia** none. **Pileipellis** a cutis with transition to a trichoderm, composed of cylindrical hyphae 8–15 µm wide, a bit constricted in septa, slightly incrustated. **Stipitipellis** of cylindrical hairs 5–15 µm wide, some with capitate apices.

Habit, Habitat & Distribution — On naked soil in tropical evergreen mixed forests. Known from Vietnam.


Typus. VIETNAM, Binh Phuoc Province, Bu Gia Map District, Bu Gia Map National Park N12.19155° E107.20422°, 520 m a.s.l., on naked soil of bank of stream in tropical evergreen mixed forests with the participation of palms, 20 Aug. 2011, O. V. Morozova (holotype LE 312536, ITS and LSU sequences GenBank OK257212 and OK257209, MycoBank MB 841774).

Notes — *Micropsalliota albofelina* is characterised by the following features: delicate, tiny pristine white discolouring to brown basidiocarps, convex pileus, cylindrical stipe without annulus, and cheilocystidia with long neck and well-distinguished capitulum. The most distinctive feature of this species is the presence of thin white hairs which cover the entire basidioma.

There are some species of *Micropsalliota* with tiny white basidiocarps: *M. alba*, *M. albella*, *M. albocericea*, *M. dentatmarginata*, *M. pudica*, and *M. pusillissima*. However, none of them is marked by so peculiarly a fuzzy surface and most of them (with the exception of *M. pudica*) possess the stipe with an annulus. Besides, *Micropsalliota alba* (described from India) differs by having wider conical pilei (4–8 mm diam), although has similar cheilocystidia with a capitulum and long neck to that of *M. albofelina* (Heinemann & Little Flower 1983). *Micropsalliota albella* (from Thailand) is distinguished by non-capitate cheilocystidia (He et al. 2020). *Micropsalliota albocericea* (known from India and Thailand) is characterised by distant ventricose lamellae,

Colour illustrations. Vietnam, Binh Phuoc Province, Bu Gia Map District, Bu Gia Map National Park, evergreen mixed forest, type locality. Spores; basidium; stipitipellis; cheilocystidia; pileipellis; basidiomata *in situ*; discolouring basidiomata (all from holotype). Scale bars = 1 cm (basidiomata), 10 µm (all others).

cymbiform spores and ventricose-capitate cheilocystidia (Zhao et al. 2010). *Micropsalliota dentatmarginata* (from China) has convex pilei and cheilocystidia with thickened base, long neck and capitate apex, but differs by wider pilei (12–15 mm diam) with appendiculate dentate margin (Li et al. 2021). *Micropsalliota pusillissima* (from Thailand) differs by smaller size of basidiomata (pileus 1–3 mm, stipe 5–12 × 0.2 mm with annulus), and ventricose-capitate cheilocystidia (Zhao et al. 2010). *Micropsalliota pudica* (from India) lacks an annulus and possesses capitate cheilocystidia, but is distinguished from *M. albofelina* by a convex-campanulate silky smooth pileus with appendiculate margin and yellow-orange discolouring of the basidiomata (Heinemann & Leelavathy 1991).



Phylogenetic tree derived from a Bayesian analysis, based on nrITS1-5.8S-ITS2 data. Analysis was performed under a GTR model of evolution, for 3 M generations, using MrBayes v. 3.2.1 (Ronquist et al. 2012). Posterior probability (PP > 0.95) values from the Bayesian analysis are shown at the nodes. The scale bar represents the expected number of nucleotide changes per site. The novel species is shown in **bold**.

Daria D. Ivanova, The Herzen State Pedagogical University of Russia, 191186, 48 Moyka Embankment, Saint Petersburg, Russia; e-mail: krakatau1349@yandex.ru

Olga V. Morozova, Komarov Botanical Institute of the Russian Academy of Sciences, 197376, 2 Prof. Popov Str., Saint Petersburg, Russia; e-mail: OMorozova@binran.ru



Fungal Planet 1366 – 24 December 2021

***Mollisia inferiseptata* Matočec, I. Kušan, Pošta, Tkalčec & Mešić, sp. nov.**

Etymology. Named after the septate spores occupying lower positions of *pars sporifera* while still inside living mature asci.

Classification — *Mollisiaceae*, *Helotiales*, *Leotiomyces*.

Ascomata apothecial, plate shaped when young, becoming richly lobed when fully mature, superficial, sessile, irregular from the top view, *0.5–1.8 mm diam, solitary or few apothecia merged, thin-fleshed in relation to size, notably leathery in consistence. Hymenium ashy grey, undulate; margin sharp, brownish grey, very finely mealy, entire, lobed, fully stretched out in maturity; excipular surface ochre-brown from apothecial base to the margin, mealy. Basal hyphae not visible. *Hymenium* *85–97 µm thick. *Asci* cylindrical with conical-subtruncate to subpapillate apex, *69.8–80.5 × 6.6–7.2 µm with *pars sporifera* *25.8–36.4 µm, †58.3–68.7 × 5–6 µm, 8-spored, base pleurorhynchous, arising from repetitive croziers, apical apparatus refractive and readily visible already in water, in Lugol's solution (IKI) apical ring medium to strongly amyloid (2–3bb) of *Calycina*-type. *Ascospores* 4–6(–8) in lower positions of *pars sporifera* inside *mature asci 2-celled and *10.4–16.4 × 2.5–3.2 µm, *Q = 3.6–5.3, while the remaining (0–)2–4 upper ones are 1-celled and *8.2–11.7 × 2.6–3 µm, *Q = 2.8–4.1, (Σn = 40), subscutiform or ellipsoid-bacilliform when septate, piscoid when 1-celled, ± heteropolar, slightly bent, hyaline, smooth, binucleate in 2-celled spores and uninucleate in 1-celled spores, freshly ejected with temporary fine sheath, biseriate inside *asci, containing very low amount of minute lipid bodies (LBs) and containing two to several small to medium sized slightly refractive globose bodies per cell, clearly visible in dark field technique; in IKI and brilliant cresyl blue (CRB) sporoplasm unstained. *Paraphyses* subcylindrical, rounded at apex, apical cell *(18.2–)22–43.3(–60.6) × (3.0–)3.3–4.1 µm, †2.5–3.1 µm wide, straight, simple, *containing single or several cylindrical vacuolar bodies (VBs), wall thin and hyaline; in KOH without yellow reaction; in IKI VBs not stained, in CRB turquoise blue to deep blue, immediately collapsing after adding KOH. *Subhymenium* *11–15 µm thick, hyaline, richly beset with highly repetitive croziers, composed of hyaline densely packed epidermoid and shortly cylindrical cells *3–5.3 µm wide. *Medullary excipulum* *24–36 µm thick at the middle flank, composed of rather compact hyaline *textura porrecta-intricata*, cells *2.1–3.4 µm wide, notably gelatinous, intercellular spaces lilac in CRB, devoid of crystals and KOH-soluble cytoplasmic bodies. *Ectal excipulum* *31–43 µm thick at the middle flank, composed of *textura globulosa-angularis*, cells *9.7–16.8 × 6.7–12.6 µm, may contain both VBs and LBs, in the inner part walls brownish, in the outer part thickened, hazel brown, *0.6–0.8 µm thick. Outermost cells with very thick blackish-brown plaque on the surface exposed to the outer environment, firmly cementing together all surface cells; excipulum devoid of intra- or intercellular crystals. *Marginal tissue* *26–29 µm thick, hairless, composed of brownish clavate cells, running at the low angle towards the surface, terminal cells

Colour illustrations. Croatia, Konavle area, forest of *Laurus nobilis* near the Ljuta river spring (type locality). Apothecia; mature living asci, crozier and paraphyses in 'H₂O'; paraphyses in 'CRB'; ascus tips in IKI; freshly ejected mature ascospores in 'H₂O' (dark and bright field, all from the same asci); subicular hyphae in 'H₂O'; middle excipular flank in 'H₂O'; marginal texture in 'H₂O'; 30-d-old colonies on wort agar (two pictures). Scale bars = 10 mm (colonies), 5 mm (single colony), 1 mm (apothecia), 10 µm (microscopic elements).

cylindrical-clavate, thick-walled, cemented together with dark brown matter on upper portions, *8.2–13.3 × 4.5–5.8 µm, may contain VBs. *Subicular hyphae* always present but of limited extent, not reaching upper flank, hyphae sparsely branched, anastomosed, and septate, *3.1–5.5 µm diam, thick-walled; walls brown and rough, *(0.8–)1.1–1.7 µm thick.

Colonies after 30 d in the dark at 20 °C on wort agar (2 % sucrose) 19.1–20 mm diam, flat discoid, slightly elevated in centre, with sparse very loosely woven whitish grey covering; margin densely radially diffuse, milky cream, forming lobed outline; underlying layer more compact, brown at the centre and near the margin, greyish-black in middle portion with slight concentric pattern; reverse margin as above, middle portion dull grey, creamy ochre at the centre, without diffuse pigmentation around colonies. Exudate drops and crystalline matter absent, but colonies produce large quantities of irregular, dark rusty brown to dark purplish brown to blackish matter. Conidiogenesis absent even within 9 mo. Aerial, covering loosely woven hyphae hyaline to subhyaline; underlying hyphae ± compact, porrectoid, pale ochre, but becoming substromatal, with thickened walls in old cultures; all hyphae smooth-walled, septate, sparsely branched, nearly without refractive content, *2.3–4 µm wide, wall *0.4–0.6 µm thick, often producing loops without any ectochroic pigment. Conidiophores and conidia absent. Asterisk (*) denotes living and cross (†) dead state. Ascus amyloidity is termed after Baral (1987), spore shape after Kušan et al. (2014), and wort agar is prepared after Fassatová (1986).

Distribution & Habitat — Known so far only from the type locality in the Konavle area, Croatia. Type collection was found on rotten trunk of *Laurus nobilis* on the forest floor, exposed to a continual water spraying under a dense canopy in the evergreen thermo-Mediterranean forest.

Typus. CROATIA, Dubrovnik-Neretva County, Konavle area, 20 m S from the Ljuta river spring, N42°32'22.5" E18°22'49.1", 70 m a.s.l., on rotten decorticated lying *Laurus nobilis* trunk, in a forest of *Laurus nobilis*, with *Ligustrum vulgare*, *Smilax aspera*, *Asparagus* sp. and *Hedera helix*, 20 Dec. 2020, N. Matočec, (holotype CNF 2/11123, ex-type culture CBS 147848, ITS and LSU sequences GenBank OK135173 and OK135174, MycoBank MB 841175).


Notes — Based on our phylogenetic analysis resulting in the concatenated ITS-LSU phylogram (see Supplementary material FP1366), our new species *Mollisia inferiseptata* belongs to the *Mollisia* s.str. species group. This clade is characterised by lignicolous apothecial species, or caulicolous as in *Neomollisia gelatinosa* (Ekanayaka et al. 2019), having a grey coloured hymenium, or greyish blue in *M. novobrunsvicensis* and *M. rava* (Tanney & Seifert 2020), or some shade of olivaceous grey in *M. cinereo-olivascens* (Saccardo 1889, Le Gal & Mangenot 1961), and ectal excipulum brown or greyish brown at least to the marginal rim. All species of this clade have an euamyloid apical ring.

(Notes continues on Supplementary material page FP1366)

Supplementary material

FP1366 Phylogenetic tree.

Murispora kazachstanica



Fungal Planet 1367 – 24 December 2021

Murispora kazachstanica Akhmetova, Kovács & D.G. Knapp, *sp. nov.**Etymology.* Referring to the origin country of the species, Kazakhstan.*Classification* — *Amniculicolaceae*, *Pleosporales*, *Dothideomycetes*.

Murispora kazachstanica differs from its closest phylogenetic neighbour, *Murispora fagicola* (MFLUCC 13-0600, ITS: NR_155379.1; LSU: NG_060797.1; SSU: NG_065639.1; *tef1*: KT709188.1), by unique fixed alleles in the ITS, LSU, SSU and *tef1* loci based on alignments of the separate loci deposited in TreeBASE as study 28727: ITS positions: 1 (A), 2 (A), 7 (T), 25 (A), 27 (C), 47 (C), 53 (T), 56 (G), 63 (T), 64 (T), 65 (A), 66 (T), 67 (C), 69 (A), 70 (T), 77 (A), 87 (A), 88 (A), 92 (T), 103 (C), 114 (A), 284 (C), 318 (T), 320 (A), 321 (G), 324 (T), 325 (C), 349 (T), 362 (C), 363 (C), 365 (T), 366 (T), 390 (A), 399 (A), 401 (T), 402 (T), 403 (T), 404 (T), 405 (T), 411 (del.), 415 (G), 417 (T), 419 (A); LSU positions: 44 (A), 135 (C), 147 (T), 150 (T), 152 (A), 234 (C), 365 (T), 376 (T), 432 (T), 433 (C), 477 (T), 478 (C), 647 (T), 835 (T); SSU positions: 51 (A), 978 (G); *tef1* positions: 39 (C), 63 (T), 96 (C), 165 (C), 192 (C), 199 (C), 200 (A), 201 (C), 222 (C), 225 (C), 252 (T), 312 (C), 513 (T), 549 (C), 615 (T), 678 (T), 687 (T), 702 (T), 729 (C), 774 (T), 789 (A), 795 (T), 798 (C), 801 (A).

Culture characteristics — Cultures covered the Petri dish on potato dextrose agar (PDA), malt extract agar (MEA) and oatmeal agar (OA) media within 3 wk at 25 °C in the dark. Diameter of the colonies on synthetic nutrient-poor agar (SNA) and Modified Melin-Norkrans (MMN) media are less than 20 mm. The margin is usually undulate on PDA, MEA, and MMN, and entire on PDA, MEA. Cultures sterile, hyphae are pigmented and usually form chlamydospores. Isolates stain the agar from orange-brown to deep red on PDA.

Typus. KAZAKHSTAN, Shortandy, from healthy roots of *Triticum aestivum*, 10 Sept. 2018, G. Akhmetova (holotype BP111491), barcode HNHM-MYC-009908, culture ex-type KG103 = CBS 148424, ITS, LSU, SSU and *tef1* sequences GenBank MZ996880, MZ997323, MZ996882 and OK019410, MycoBank MB 841081.


Additional material examined. KAZAKHSTAN, Shortandy, from roots of *Hordeum vulgare*, 10 Sept. 2018, G. Akhmetova paratype BP111490, barcode HNHM-MYC-009907, culture KG93 = CBS 148424, ITS, LSU, SSU and *tef1* sequences GenBank MZ996879, MZ997322, MZ996881 and OK019409.

Notes — Based on a megablast search of NCBI's GenBank nucleotide database, the closest hits of *Murispora kazachstanica* (KG103 = CBS 148425) using the ITS sequence are *Murispora aquatica* (GenBank MN325085.1; Identities = 423/450 (94 %), six gaps (1 %)), *Murispora fagicola* (GenBank NR_155379.1; Identities = 349/376 (93 %), five gaps (1 %)), and *Murispora medicaginicola* (GenBank NR_155380.1; Identities = 401/436 (92 %), four gaps (0 %)). The closest hits using the LSU sequence are *Murispora galii* (GenBank KT709175.1; Identities = 887/898 (99 %), one gap (0 %)), *Murispora cardui* (GenBank NG_059607.1; Identities = 885/898 (99 %), no gaps), and *Murispora hawksworthii* (GenBank KT709180.1; Identities = 885/898 (99 %), one gap (0 %)). The closest hits

Colour illustrations. The barley (*Hordeum vulgare*) field in Shortandy, Northern Kazakhstan, where *Murispora kazachstanica* strain KG93 was isolated in 2018 from healthy roots of the plant. Colony of KG103 (upper) and KG93 (middle) on PDA media; pigmented septate hyphae and chlamydospores of strain KG93 (lower) on PDA media. Scale bar = 20 µm.


using the SSU sequence are *Murispora hawksworthii* (GenBank NG_063592.1; Identities = 1051/1052 (99 %), one gap (0 %)), *Murispora cicognanii* (GenBank NG_063591.1; Identities = 1050/1052 (99 %), one gap (0 %)), and *Murispora medicaginicola* (GenBank NG_063590.1; Identities = 1048/1052 (99 %), three gaps (0 %)). The closest hits using the *tef1* sequence are *Murispora medicaginicola* (GenBank KT709191.1; Identities = 761/779 (98 %), no gaps), *Murispora aquatica* (GenBank MN337969.1; Identities = 708/725 (98 %), no gaps), and *Murispora cicognanii* (GenBank MK109804.1; Identities = 783/802 (98 %), no gaps).

No sporulation of both *Murispora kazachstanica* isolates was observed in any of the media PDA, MEA, SNA, OM, MMN and WA supplemented with various autoclaved plant tissues (methods described in Knapp et al. 2015).



Maximum Likelihood (RAxML) tree of concatenated ITS, LSU, SSU, and *tef1* sequence alignments of *Murispora* species and representative taxa of related families sensu Wanasinghe et al. (2015) and Bao et al. (2019). All the GenBank accession numbers of the sequences used for the analysis can be found in Bao et al. (2019). *Murispora kazachstanica* isolates are highlighted in **bold**. The RAxML analysis was performed by raxmlGUI v. 1.3 (Silvestro & Michalak 2012), bootstrap support values ($\geq 70\%$) are shown above branches and before slashes. The Bayesian analysis was performed with MrBayes v. 3.2.1 (Ronquist et al. 2012) and Bayesian posterior probabilities (≥ 0.90) are shown below branches and after slashes. *Quadricrura septentrionalis* (CBS 125428), *Tetraploa sasicola* (KT 563) and *Triplosphaeria maxima* (KT 870) served as multiple outgroups. The scale bar indicates the expected number of changes per site.

Neodevriesia kalakoutskii



Fungal Planet 1368 – 24 December 2021

Neodevriesia kalakoutsii Kochkina, Kachalkin, Ivanushkina, Trubitsyn & Ozerskaya, sp. nov.

Etymology. Named in honour of Lev Kalakoutsii, a famous Russian microbiologist, for his contributions to the study of microbial diversity and development of the All-Russian Collection of Microorganisms.

Classification — *Neodevriesiaceae*, *Mycosphaerellales*, *Dothideomycetes*.

Mycelium consisting of branched, septate, pale brown, smooth, 2–3 µm wide hyphae. **Conidiophores** rarely micronematous, predominantly macronematous, solitary, erect, arising from superficial mycelium, subcylindrical, predominantly unbranched, medium brown, smooth, straight, thick-walled, 10–50 × 2–3 µm, 0–2-septate. **Conidiogenous** cells terminal, medium to dark brown, subcylindrical, smooth, 5–15 × 2–3 µm; rarely proliferating sympodially; conidiogenous loci, thickened and darkened, 1–1.5 µm wide. **Ramoconidia** medium brown, smooth, subcylindrical, (9–)11–15(–17) × 2.5–3.5 µm, 0–1-septate, predominantly with two conidiogenous loci, scars darkened and thickened, 1 µm diam. **Conidia** occurring in long branched chains, pale to medium brown, smooth, subcylindrical to narrowly fusoid-ellipsoidal or obclavate, 0(–1)-septate, hila darkened, 1–1.5 µm wide, sometimes situated on denticles up to 2.5 µm long, intercalary conidia with flattened ends, (6–)7–11(–15) × 2.5–3.5 µm; apical conidia with rounded apex, (4–)5–7(–9) × 2–2.5 µm. Fragmenting conidial segments from aerial hyphae abundant.

Culture characteristics — On potato dextrose agar (PDA), 20 °C after 1 mo colonies grey (3E1; Kornerup & Wanscher 1978), dense, aerial mycelium abundant, felty to woolly, erumpent in the centre, flat and entire at the margin, reverse olive grey (3F2); colonies reaching 15–16 mm diam (at 20 °C), 12–14 mm diam (at 15 °C), 3–4 mm diam (at 5 °C), no growth (at 25 °C).


Typus. NORWAY, West Spitsbergen, near Barentsburg on the southern bank of Isfjorden, isolated from permafrost, 30 Oct. 2020, V.E. Trubitsyn, WS-F1 (holotype VKM H-0002 preserved in a metabolically inactive state, culture ex-type VKM F-4872 = CBS 148320, SSU, ITS-D1/D3 domains of LSU nrDNA, *tef1*, *rpb1* and *rpb2* sequences GenBank MZ317563, MZ025963, OD937129, OU070195 and OU070196, MycoBank MB 840057).

Notes — Based on a blastn search of NCBI's GenBank nucleotide database, the closest hits to *Neodevriesia kalakoutsii* using the ITS sequence are uncultured fungus clones RS-Apr-ITS18 and RS-Oct-ITS03 from saline efflorescence on rock shelter (GenBank KY096661 and KY096647; Identities = 516/517 (99.81 %), one gap), *N. metrosideri* (strain CBS 145084, GenBank NR_161141; Identities = 497/518 (95.95 %), eight gaps) and *N. lagerstroemiae* (strain CPC 14403, GenBank GU214634; Identities = 494/518 (95.37 %), six gaps); using the LSU sequence it is *N. modesta* (strain CCFFEE 5672, GenBank

Colour illustrations. Dry-drilling and sampling site in West Spitsbergen. *Neodevriesia kalakoutsii* VKM F-4872: colony after 30 d at 20 °C on PDA; conidiophores; conidia and fragmenting conidial segments from aerial hyphae. Scale bars = 10 µm.


KF310026; Identities = 756/758 (99.74 %), two gaps); using the SSU sequence these are *Neodevriesia* sp. (strain MUT 4891, GenBank KT587311; Identities = 1599/1601 (99.88 %), no gaps) and *Neodevriesia* sp. (strain CPC 11876, GenBank GU214564; Identities = 1595/1601 (99.63 %), no gaps); using *tef1* it is *N. strelitziae* (strain CBS 122379, GenBank GU349049; Identities = 841/900 (93.44 %), two gaps); using *rpb2* it is *N. strelitziae* (strain CBS 122379, GenBank GU371738; Identities = 849/1071 (79.27 %), 14 gaps); using *rpb1* it is '*Capnodiales*' sp. (strain S2, GenBank GU250946; Identities = 669/770 (86.88 %), 15 gaps).

Neodevriesia kalakoutsii micromorphologically differs from the phylogenetically closely related species *N. modesta* (Egidi et al. 2014), also growing at lower temperatures, in having distinctly differentiated conidiophores bearing long branched chains and having ramoconidia. Conidia are predominantly aseptate and not so uniform in size and shape.



Galina A. Kochkina, Nataliya E. Ivanushkina, Vladimir E. Trubitsyn & Svetlana M. Ozerskaya, All-Russian Collection of Microorganisms, G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Pushchino Center for Biological Research of the Russian Academy of Sciences, 142290, Pushchino, pr. Nauki, 5, Russia; e-mail: gak@dol.ru, nei@dol.ru, lichoradkin43@gmail.com & smovkm@gmail.com

Aleksey V. Kachalkin, Lomonosov Moscow State University, 119234, Moscow, Leninskie Gory Str. 1/12, Russia, and All-Russian Collection of Microorganisms, G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Pushchino Center for Biological Research of the Russian Academy of Sciences, 142290, Pushchino, pr. Nauki, 5, Russia; e-mail: kachalkin_a@mail.ru



Fungal Planet 1369 – 24 December 2021

Papiliotrema horticola Kachalkin, A.M. Glushakova & M.A. Tomashevskaya, *sp. nov.*

Etymology. Named after 'hortus', as many known strains were isolated from the horticultural plants.

Classification — *Rhynchogastremataceae*, *Tremellales*, *Tremellomycetes*.

On glucose peptone yeast extract agar (GPYA) and 5 % malt extract agar (MEA), after 7 d at 21 °C, *streak* is cream to ivory coloured, smooth and semi-glossy, with an entire margin. *Cells* are subglobose and ellipsoidal (1.8–4.5 × 3.2–8.0 µm), occurring singly or in pairs, dividing by polar budding. *Sexual structures*, *pseudohyphae*, *true hyphae* and *ballistoconidia* have not been observed during 6 wk at 21 °C in culture (pure cultures and in mating test) grown on GPYA, MEA, potato dextrose agar (PDA), cornmeal agar (CMA), McClary acetate agar. Glucose is not fermented. Glucose, galactose, sucrose, maltose, trehalose, lactose, melibiose, cellobiose, raffinose, melezitose, soluble starch (delayed and weak), D-xylose, L-arabinose, D-arabinose, D-ribose, L-rhamnose, D-glucosamine, ethanol, glycerol, ribitol, galactitol, D-mannitol, D-glucitol, myo-inositol, methyl alpha-D-glucoside, salicin, DL-lactic acid (weak), succinic acid, citric acid (weak), 2-keto-D-gluconate, 5-keto-D-gluconate, arbutin, L-malic acid and saccharic acid are assimilated; no growth occurs on L-sorbose, inulin, erythritol, methanol, m-hydroxybenzoic acid, p-hydroxybenzoic acid, gallic acid, ferulic acid, vanillic acid, protocatechuic acid and veratric acid. Assimilation of nitrogen compounds: positive for ammonium sulphate, L-lysine and cadaverine, and negative for potassium nitrate. Growth on vitamin-free medium, on 50 % w/w glucose / yeast extract (0.5 %) agar, on MEA with 10 % NaCl is positive. Growth with 0.01 % and 0.1 % cycloheximide is negative. Starch-like compounds are produced. Diazonium blue B colour and urease reactions are positive. Maximum growth temperature is 25–26 °C.

Typus. RUSSIA, Vladimir Oblast, isolated as endophyte from the hypanthium of apples (*Malus communis*), Oct. 2019, A.M. Glushakova, ok6-19 (holotype KBP Y-6685, preserved in a metabolically inactive state, culture ex-type VKM Y-3061 = DSM 111206 = CBS 16715, ITS-D1/D2 domains of LSU nrDNA, SSU, *tef1*, *rpb1* and *rpb2* sequences GenBank MW579431, MT636882, LR814014, LR814015 and HG993247, MycoBank MB 840888).

Additional materials examined. RUSSIA, Moscow, isolated as endophyte from leaves of *Lactuca sativa*, Apr. 2020, A.M. Glushakova, KBP YE-0975, ITS and D1 domain of LSU nrDNA sequence GenBank MZ749747; isolated as endophyte from leaves of *Brassica rapa* subsp. *pekinensis*, Apr. 2020, A.M. Glushakova, KBP YE-0998, ITS and D1 domain of LSU nrDNA sequence GenBank MZ749748.

Colour illustrations. Russia, Vladimir Oblast, apples on local market. *Papiliotrema horticola* KBP Y-6685: growth of yeast colonies and yeast cells on MEA (after 7 d at 21 °C). Scale bar = 5 µm.


Notes — Based on a blastn search of NCBI's GenBank nucleotide database, the closest hits using the **ITS** sequence are '*Cryptococcus*' sp. from *Vitis vinifera* collected in USA (P40A001, GenBank JX188145; Identities = 527/527 (100 %)), strain under patent deposit *Papiliotrema cf. aureus* (NRRL Y-30215, GenBank EF056299; Identities = 384/385 (99 %), no gaps) and endophytic strains *Papiliotrema aurea* from wheat collected in Italy (AD12 and AD17, GenBank MN913559 and MN913560; Identities = 484/485 (99 %), no gaps), but this result differs from the ex-type strain of *P. aurea* (CBS 318^T, GenBank AB035045; Identities = 424/440 (97 %), one gap (0 %)). The closest hits using the **LSU** sequence are '*Cryptococcus*' sp. (P40A001, GenBank JX188145; Identities = 844/844 (100 %)) and *P. aurea* (CBS 12077, GenBank KY108729; Identities = 842/844 (99 %), no gaps), but this result also differs from the ex-type strain of *P. aurea* (CBS 318^T, GenBank AB035041; Identities = 562/570 (99 %), no gaps); using **SSU** it is *P. aurea* (CBS 318^T, GenBank NG_062615; Identities = 1658/1664 (99 %), no gaps), using **tef1** it is *P. aurea* (CBS 318^T, GenBank KF037035; Identities = 480/531 (90 %), 12 gaps (2 %)), using **rpb1** it is *P. aurea* (CBS 318^T, GenBank LK024017; Identities = 621/680 (91 %), two gaps (0 %)), using **rpb2** it is *P. aurea* (CBS 318^T, GenBank KF036763; Identities = 744/825 (90 %), seven gaps (0 %)). Based on the thresholds for nrDNA as DNA barcodes (Vu et al. 2016), strains *P. aurea* CBS 318^T and KBP Y-6685 belong to different species. Additionally, the study of the neighbouring clade including *P. flavescens* and *P. terrestris* strains (Yurkov et al. 2015) was able to reveal that the pairwise sequence alignments using blast have intraspecific differences in *Papiliotrema* much less than 10 % for the *tef1* and *rpb1* sequences. The phylogenetic position of the new species is demonstrated using the combined ITS and LSU rDNA phylogeny and multi-locus phylogeny based on the ribosomal DNA cistron, *rpb1* and *tef1* sequences. *Papiliotrema horticola* sp. nov. can be phenotypically distinguished from the closest species *P. aurea* by assimilation ethanol, DL-lactic acid, saccharic acid (data for ex-type strain VKM Y-328), growth on vitamin-free medium, on media with 50 % glucose and 10 % NaCl, maximum temperature of growth, and size of cells.

Supplementary material**FP1369** Phylogenetic tree.

Aleksey V. Kachalkin, Lomonosov Moscow State University, 119234, Moscow, Leninskie Gory Str. 1/12, Russia, and All-Russian Collection of Microorganisms, G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms RAS, 142290, Pushchino, pr. Nauki 5, Russia; e-mail: kachalkin_a@mail.ru

Anna M. Glushakova, Lomonosov Moscow State University, 119234, Moscow, Leninskie Gory Str. 1/12, Russia, and Mechnikov Research Institute for Vaccines and Sera, 105064, Moscow, Maly Kazenny by-street, 5A, Russia; e-mail: glushakova.anya@yandex.ru
Maria A. Tomashevskaya, All-Russian Collection of Microorganisms, G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms RAS, 142290, Pushchino, pr. Nauki 5, Russia; e-mail: tomkotik@rambler.ru

Paraphoma variabilis



Fungal Planet 1370 – 24 December 2021

Paraphoma variabilis Magaña-Dueñas, Cano-Lira & Stchigel, *sp. nov.*

Etymology. From Latin *variabilis*, due to the variable shape of the conidia.

Classification — *Didymellaceae*, *Pleosporales*, *Dothideo-mycetes*.


Hyphae hyaline, septate, smooth- and thin-walled, 1.5–3 µm wide. **Conidiomata** pycnidial, immersed to semi-immersed, solitary, scattered, pale brown at the base, with an ostiolate neck, subglobose to pyriform, 150–180 × 175–200 µm, covered with very short setae. **Conidiomatal wall** 3–5-layered, 10–20 µm wide, translucent, of *textura angularis*, composed by an outer layer of very pale brown to dark brown, flattened polygonal cells of 4–6 µm diam; setae subhyaline, rounded at the tip, finger-like, 7–25 × 2.5–3.0 µm, ostiolar neck brown to dark brown, conical-truncate, 55–60 × 25–30 µm. **Conidiophores** absent. **Conidiogenous cells** phialidic, ampulliform to cylindrical, determinate, hyaline, smooth-walled, 5–8 × 1.5–3 µm. **Conidia** aseptate, hyaline, smooth- and thin-walled, variable in shape, cylindrical, ellipsoidal, ovate, reniform, sigmoid, 4–8 × 2–3 µm, truncate at the base.

Culture characteristics — (after 7 d at 25 °C). Colonies on PDA reaching 37–41 mm diam, radiated, velvety, margin undulate, with abundant aerial mycelium, grey to yellowish grey (29F1/29C1; Methuen Handbook of Colour); reverse bluish grey to yellowish grey (23F3/4B2), diffusible pigment not produced. Colonies on OA reaching 40 mm diam, convex, velvety, margin regular, with abundant aerial mycelium, surface and reverse grey (20A1), diffusible pigment not produced. Colonies on MEA

2 % reaching 25 mm diam, flattened, velvety, margin lobate, with abundant aerial mycelium, grey to orange grey (30E1/5B1); reverse grey to dark grey (30D1/30F1), diffusible pigment not produced. Cardinal growing temperatures: optimum 25 °C, maximum 30 °C, minimum 5 °C

Typus. SPAIN, Tarragona Province, Els Ports de Tortosa-Beseit, from dung, Oct. 2017, coll. I.A. Iturrieta-González, isol. V. Magaña-Dueñas (holotype CBS H-24765, cultures ex-type CBS 147695 = FMR 17160; ITS, LSU, *rpb2* and *tub2* sequences GenBank LR993310, LR993311, LR993313 and LR993314, MycoBank MB 839143).

Notes — Based on a megablast search of NCBI's GenBank nucleotide database, the closest hit using the **LSU** sequence was *Phaeosphaeriopsis nolinae* (strain CBS 102205, GenBank KY090667; Identities = 552/552 (100 %), no gaps). The closest hit using the **ITS** sequence was *Paraphoma fimeti* (strain UTHSC DI16-296, GenBank LT796872; Identities = 528/585 (90 %), eight gaps (1 %)). The closest hit using the **rpb2** sequence was *Paraphoma fimeti* (strain UTHSC DI16-296, GenBank LT797032; Identities = 842/905 (93 %), no gaps). The closest hit using the **tub2** sequence was *Paraphoma fimeti* (strain UTHSC DI16-296, GenBank LT796952; Identities = 294/320 (92 %), five gaps (1 %)). Based on the combined analysis of ITS, LSU, *rpb2* and *tub2* sequences, the closest relative of *Paraphoma variabilis* is *Paraphoma fimeti*. The latter differs from our novel species in that it produces smaller (3–5 × 2–3 µm vs 4–8 × 2–3 µm) ellipsoidal conidia (Boerema et al. 2004), which are cylindrical, ellipsoidal, ovate, reniform and sigmoid, and truncated at the base, in *P. variabilis*. Also, *P. fimeti* produce a yellowish diffusible pigment, which is absent in *P. variabilis*.



Colour illustrations. Ports de Totosa-Beseit natural park, Tarragona, Spain. Colony on PDA and OA after 14 d at 25 + 1 °C; pyriform conidioma; conidiogenous cells; conidia. Scale bars = 25 µm (conidioma) and 10 µm (all others).

Maximum likelihood tree obtained from the combined DNA sequences dataset from four loci (ITS, LSU, *rpb2* and *tub2*) of our isolate and sequences retrieved from the GenBank database. Ex-type strains of the different species are indicated with superscript [†]. The new species proposed in this study is indicated in **bold**. The RAxML v. 8.2.10 (Stamatakis 2014) bootstrap support values (> 70 %) are provided at the nodes. *Neophaeosphaeria agaves* CPC 21264 and *Neophaeosphaeria filamentosa* CBS 102202 were used as outgroup.

Phaeococcomyces kinklidotomophilus



Fungal Planet 1371 – 24 December 2021

Phaeococcomyces kinklomatophilus Sastoque, Stchigel & Cano-Lira, *sp. nov.*

Etymology. From Greek κίγκλιδωμα-, railing, and - φίλος, friend, because the environmental origin of the fungus.


Classification — *Phaeococcomycetaceae*, *Lichenostigmatales*, *Arthoniomycetes*.

Colonies lacking mycelium but consisting of a mass of yeast-like cells embedded in a mucous matrix, which are aseptate, hyaline, smooth- and thin-walled, ellipsoid to globose, 3–6 × 2–5 µm when young, producing blastoconidia via multilateral budding, becoming brown, verrucose and globose with the age, 5–8 µm diam, remaining attached to one another to form up to 5-celled chains.

Culture characteristics — (after 14 d at 25 °C in the dark). Colonies reaching 3.5 mm diam on malt extract agar (MEA), 2.0 mm on oatmeal agar (OA) and potato carrot agar (PCA), and 2.5 mm on potato dextrose agar (PDA), circular, margins entire, flat to slightly convex, smooth and glistening on OA and PCA, but crater-like, dry and slightly roughened and tough with lobate margins when older on MEA and PDA. Surface and reverse black on all culture media tested. Minimum, optimum and maximum temperature of growth: 5 °C, 15–25 °C, and 25 °C, respectively.

Typus. SPAIN, Tarragona Province, Els Pallaresos, isolated from a blackened metal railing of an industrial warehouse, 20 Nov. 2020, coll. J. F. Cano-Lira & A. M. Stchigel, isol. A.P. Sastoque (holotype CBS H-24766, culture ex-type FMR 18615 = CBS 147696; ITS and LSU sequences GenBank HG995460 and HG995431, MycoBank MB 839137).


Notes — Based on a megablast search of NCBI's GenBank nucleotide database, the closest hit using the **ITS** sequence was *Phaeococcomyces eucalypti* (strain CPC 17606, GenBank NR_120226; Identities = 494/535 (92.34 %), eight gaps (1 %)), and using the **LSU** sequence it was *Phaeococcomyces eucalypti* (strain CPC 17606, GenBank NG_042748; Identities = 463/473 (97.89 %), two gaps (0 %)). Our ITS phylogenetic tree corroborated the placement of our isolate as a new species of the genus, being phylogenetically close to *P. eucalypti*. *Phaeococcomyces kinklomatophilus* was isolated by swabbing from the surface of a metal railing in Els Pallaresos village, Tarragona province, Catalonia, Spain. *Phaeococcomyces* is a genus of black yeasts with slimy, folded, slow-growing colonies lacking of aerial mycelium. The genus currently contains four species: *P. eucalypti*, *P. mexicanus*, *P. nigricans* (the type species) and *P. rothmanniae*. *Phaeococcomyces kinklomatophilus* is morphologically very similar to other species of the genus, but differs from *P. eucalypti*, its phylogenetically closest species, in size (3.5 mm vs 25 mm diam on MEA), shape (crater-shaped vs folded on MEA and PDA) and colour (nearly black vs iron-grey on MEA and PDA) of the colonies. Moreover, *P. kinklomatophilus* is molecularly easily distinguishable from the other species of the genus.



Colour illustrations. Railing with chromatic alteration in Els Pallaresos village, Tarragona Province, Spain. Colonies growing on PDA culture media (at 25 °C); blastoconidia (DIC and PH respectively). Scale bars = 10 µm.

Maximum likelihood tree obtained from the ITS alignment of our isolate and sequences retrieved from GenBank. The tree was built by using RAXML CIPRES (http://www.phylo.org/sub_sections/portal/) (Miller et al. 2010). Bootstrap support values ≥ 70 % are presented at the nodes. *Anteaglonium gordoniae* SGSF172 and *Flammeascoa lignicola* MFLUCC 10-0128a were used as outgroups. The new species proposed in this study is indicated in **bold**. ^TRepresents the ex-type strain of the species.

Penicillium xyleborini



Fungal Planet 1372 – 24 December 2021

***Penicillium xyleborini* Visagie & W.J. Nel, sp. nov.**

Etymology. Latin, *xyleborini*, refers to the genus of the beetle host (*Xyleborinus saxesenii*) from which this species was isolated.

Classification — *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*.


Conidiophores biverticillate to terverticillate; **stipes** smooth-walled, 100–200(–500) × 2.5–4(–4.5) µm; **branches** 10–30 µm; **metulae** 4–6 per stipe/branch, (8–)10–16 × 3–4 µm; **phialides** 6–10 per metula, 8–11 × 3–4 µm (av. = 9.5 ± 0.23 × 3.3 ± 0.2), av. length metula/phialide 1.3; **conidia** smooth-walled, globose to subglobose, 2.5–3.5 × 2.5–3.5 µm (av. = 3.1 ± 0.2 × 2.94 ± 0.2), av. width/length = 0.95, n = 57.

Culture characteristics — (25 °C, 7 d): On Czapek yeast autolysate agar (CYA): colonies moderately deep, concentrically and radially sulcate, crateriform; margins low, narrow (1 mm), entire; mycelia white; texture velutinous; sporulation moderately dense, conidia *en masse* greyish to dull green (26D4–F4–5; Kornerup & Wanscher 1967); soluble pigments absent; exudates clear to inconspicuously yellow; reverse greyish yellow, olive brown (4B3, 4E4–5). On malt extract agar (MEA): colonies moderately deep, concentrically sulcate, crateriform; margins low, narrow (1 mm), somewhat irregular; mycelia white; texture velutinous; sporulation moderately dense, conidia *en masse* dull to greyish green (25D4–5); soluble pigments absent; exudates absent; reverse brown (6E6), dark brown (6F6). On yeast extract sucrose agar (YES): colonies deep, radially and concentrically sulcate, crateriform; margins low, narrow (1 mm), somewhat irregular; mycelia white; texture velutinous; sporulation moderately dense, conidia *en masse* greyish green (25C4); soluble pigments absent; exudates absent; reverse yellowish white, greyish yellow, olive brown (4A2, 4B3, 4E4–5). On dichloran 18 % glycerol agar (DG18): colonies moderately deep, radially and concentrically sulcate; margins low, narrow (1 mm), entire; mycelia white; texture velutinous; sporulation moderately dense, conidia *en masse* greyish to dull green (26D4–F4–5);

soluble pigments inconspicuously yellow; exudates absent; reverse greyish yellow, greenish white (4B4–C4, 29A2–30A2). On oatmeal agar (OA): colonies low, plane; margins low, narrow (1 mm), entire; mycelia white, inconspicuously yellow; texture velutinous; sporulation moderately dense, conidia *en masse* dull green (27D4–F5); soluble pigments absent; exudates clear; reverse yellowish white, (2A2), olive yellow (2C6). On creatine sucrose agar (CREA): colonies weak growth, no acid production. Colony diam (in mm, after 7 d, at 25 °C): CYA 15–16; CYA 15 °C 18–20; CYA 20 °C 20–24; CYA 30 °C no growth; CYA 37 °C no growth; CYAS 18–21; MEA 12–14; DG18 15–18; YES 18–20; OA 15–18; CREA 13–18.

Typus. SOUTH AFRICA, Limpopo Province, Tzaneen, (–23.708192, 30.099344), from an ambrosia beetle (*Xyleborinus saxesenii*), 30 Sept. 2019, *W. Nel* (holotype PREM 63078 (dried specimen), culture ex-type CMW 56800 = CN 001D4 = WN 2019.10.32; LSU, ITS, *BenA*, *CaM* and *rpb2* sequences GenBank MW504290, MW504356, MW480817, MW480823 and MW480824, MycoBank MB 840990).

Notes — A multigene phylogeny resolves *Penicillium xyleborini* as a close relative of *P. lusitanum* in sect *Ramosum* series *Soppiorum* (Houbraken et al. 2020). The new species produces greyish to dull green conidia, compared to the dark green conidia of *P. lusitanum*. Also, *P. xyleborini* conidiophores are more commonly terverticillate, with longer phialides and larger conidia when compared with *P. lusitanum* (Gonçalves et al. 2019). Other *Penicillium* species isolated from ambrosia beetles or their galleries during this project included *P. brevicompactum*, *P. cairnsense*, *P. citreonigrum*, *P. coprophilum*, *P. glabrum*, *P. ibericum*, *P. pancosmium*, *P. pasqualense*, *P. soliforme*, *P. ubiquetum*, *P. viticola* and a new section *Sclerotiorum* species that will be described elsewhere. Sequences of these species were submitted to GenBank and accessioned with numbers MW480792–MW480820.



Colour illustrations. The ambrosia beetle (*Xyleborinus saxesenii*) from which the new species was isolated. *Penicillium xyleborini* colonies on CYA15C, CYA20C, CYA, CYAS, MEA, YES, DG18, OA, CREA (from top to bottom); conidiophores; conidia. Scale bars = 10 µm.

Combined phylogeny of *Penicillium* section *Ramosum* based on ITS, *BenA*, *CaM* and *rpb2*. Aligned data sets (MAFFT v. 7.454; Katoh et al. 2013) were analysed using maximum likelihood (IQ-TREE v. 1.6.12; Minh et al. 2020) and Bayesian Inference (MrBayes v. 3.2.7a; Ronquist et al. 2012). Maximum likelihood bootstrap support values (≥ 80 %) and Bayesian posterior probability values (≥ 0.95) are given above branches. The new species is indicated by **bold text**, T = ex-type strain and GenBank accession numbers are shown in a smaller font next to the culture accession number (ITS = green, *BenA* = blue, *CaM* = red, *rpb2* = purple).



Fungal Planet 1373 – 24 December 2021

***Phylloporia tamilnadensis* Sridhar, Palani & Decock, sp. nov.**

Etymology. Name refers to the southwestern State of Tamil Nadu, where this fungus was collected.

Classification — *Hymenochaetaceae*, *Hymenochaetales*, *Basidiomycota*.

Basidiomata annual, mostly solitary, pileate, sessile; *pileus* spongy to leathery when fresh, turning horny to brittle on drying, initially arises as small dot gradually widening in all direction to form distinct basidiomata, infrequently laterally fused with adjacent basidiomata, easily separable from host, 0.8–20 × 0.6–10 mm, 0.8–20 mm thick near point of attachment, applanate to circular, hoof-shaped, broadly to narrowly attached, more or less bent at margin, pale orange (5A3, Kornerup & Wanscher 1978); *pilear* surface tomentose to velvety, irregularly zonate, pitted, melon (5A6) to pale orange (5A3); *margin* 0.6–1 mm wide, sterile, entire, more or less lobed at maturity, reddish blond (5C4) to pale orange; *hymenophore* poroid; pores round to angular mostly, 3–4 per mm; dissepiments thin, entire to lacerate; *context* up to 3 mm thick, distinctly duplex, the lower part homogenous, dense, brownish orange (5C4), very thin into margin up to 0.4–1.0 mm at base, with a distinct to indistinct thin black line subtending an upper tomentum, porous spongy; *tubes* 0.6–10 mm long, orange when fresh (6A8), the tubes starting 0.3 mm from the margin. **Hyphal system** monomitic in all parts; generative hyphae simple septate, thin- to slightly thick-walled, hyaline, yellowish to light golden brown, darker, brownish in KOH, scarcely ramified, the branches constricted at their emergence point, soon growing parallel to mother hyphae; *in the tomentum*, hyphae parallel, adpressed first, soon erected, loosely packed, straight to sinuous, mostly unbranched, slightly thick-walled, sub-hyaline to pale golden brown, from 3.5 µm diam near the base, gradually enlarging up to 4.5–8.0 µm; *in the context* hyphae adpressed to oblique but mostly erected, with a near parallel orientation, moderately thick-walled with the lumen widely open, septate, but with long aseptate segments, 3.5–4.0 µm diam; *in the hymenophoral trama* hyphae with a subparallel disposition, straight, occasionally geniculated, the lumen widely open, septate, with aseptate segments or with occasional secondary septa, 2.5–3.5 µm diam. **Hymenium:** *basidia* ~11 × 5.5 µm, clavate, septate at base, with four sterigmata up to 5 µm long; *basidioles* ~10.5 × 5 µm; *basidiospores* 3.5–5.0 × 2.3–3.6 µm (av. 4.3 × 2.9 µm), Q = 1.3, ellipsoid, moderately thick-walled, smooth, hyaline to pale yellow, inamyloid, acyanophilic.

Colour illustrations. India, Tamil Nadu State, Ranipet District, Sholinghur Taluk, *Catunaregam spinosa* growing in Mettu kunnathur village. Habitat: basidiomata pore view; transverse section of basidiomata; pilear surface details; basidium and basidiospores. Scale bars = 1 cm (basidiomata, pore view and T.S. of basidiomata); 50 µm (pilear surface details); 10 µm (basidium and basidiospores).

Typus. INDIA, Tamil Nadu State, Ranipet District, Sholinghur Taluk, Mettu kunnathur Village, N13°02'17" E79°31'10", growing on thin branches of living plant of *Catunaregam spinosa* (Rubiaceae), 10 Dec. 2017, P. Sridhar, MKRF 05 (holotype CAL #1840, LSU sequence GenBank MN525278, MycoBank MB 838470).

Additional material examined. INDIA, Tamil Nadu State, Ranipet District, Sholinghur Taluk, Mettu kunnathur Village, N13°02'17" E79°31'10", growing on thin branches of living plant of *C. spinosa*, 10 Nov. 2018, P. Sridhar, MKRF 21 (LSU sequence GenBank MN525279).

Notes — The olive yellow cap and ash grey dots on the stipe surface, as well as larger spores, distinguish *S. praetermissus* from the closest species of the *Suillus acidus* group. According to phylogenetic analysis, the nearest species is *S. subalutaceus*, p-distance is 1.8 %, 12 nucleotide substitutions per 666 alignment positions of ITS1-5.8S-ITS2. According to a microscopic examination of the type specimen of *S. subalutaceus* (MICH 50221) carried out by E. Zvyagina, its spores are much shorter with the same width. The cap surface of *S. subalutaceus* is 'pinkish-buff' (Smith & Thiers 1964). *Suillus subalutaceus* forms mycorrhiza with two-needle pines (Nguyen et al. 2016), while the new taxon *S. praetermissus* is associated with five-needle pines. In our study all examined specimens were collected under *Pinus sibirica*. However, for specimens and mycorrhizal samples from Japan, of which sequences were used in the phylogenetic tree, an association with *P. corayensis* and *P. pumila* was also noted. *Suillus subalutaceus* is an American species, while *S. praetermissus* has an Asian distribution.

Supplementary material

FP1373 Phylogenetic tree.



Fungal Planet 1374 – 24 December 2021

***Phytophthora docyniae* Q.N. Dang, T.H. Nguyen & T.I. Burgess, sp. nov.**

Etymology. Refers to the genus name of the only known host, *Docynia indica*.

Classification — *Peronosporaceae*, *Peronosporidae*, *Oomycota*.


Sporangia produced abundantly in non-sterile soil extract after 2 d, terminally borne on unbranched sporangiophores, non-papillate, ellipsoid to ovoid, $34.5 \pm 2.7 \times 23.8 \pm 2.7 \mu\text{m}$ (overall range $30\text{--}39.5 \times 19.8\text{--}27.9 \mu\text{m}$), length/breadth ratio 1.45 ± 0.9 , with internal proliferation sporangia in both a nested and extended way. Direct germination of sporangia with unusual shapes was often observed. *Hyphal swellings* are globose, subglobose, irregular, mostly catenulate and sometimes clustered. *Chlamydospores* produced abundantly on all media (potato dextrose agar (PDA), carrot agar (CA), V8 agar) and in soil extract, mostly intercalary, globose, subglobose and irregular, $43.4\text{--}96.3 \mu\text{m}$ diam, very thick-walled on PDA, 4.2 ± 1.1 ($2.0\text{--}6.9 \mu\text{m}$). Chlamydospores often germinated and produced sporangia in soil extract. *Gametangia* not produced in single culture or when paired with A1 and A2 tester strains of *P. cinnamomi* (MP75, DCE60). Radial growth rates on V8 agar at optimum temperature ($20\text{--}32.5^\circ\text{C}$) and near the maximum temperature (35°C), $4.2 \pm 0.8 \text{ mm/d}$ and $2.2 \pm 0.13 \text{ mm/d}$, respectively. Minimum temperature for growth is 15°C .

Culture characteristics — Colonies are chrysanthemum on PDA and V8 agar, while growth on CA showed no distinct pattern. Growth on CA and V8A was faster than on PDA.

Typus. VIETNAM, Yen Bai Province, Mu Cang Chai, isolated from the soil and root of *Docynia indica* (*Rosaceae*), Oct. 2019, V.L. Binh & X.T. Hung (holotype CBS H-24780, culture ex-type CBS 148031 = FP77, ITS, *cox1*, β -tubulin and LSU sequences GenBank MZ420209, MZ435257, MZ435255 and MZ435254, MycoBank MB 840241).

Additional material examined. VIETNAM, Yen Bai Province, Mu Cang Chai, isolated from the soil and roots of *D. indica*, Oct. 2019, V.L. Binh & X.T. Hung, culture FP78.

Notes — Phylogenetically, *Phytophthora docyniae* resides in clade 9c of the *Phytophthora* phylogeny (Abad et al. 2019) most closely related to *P. macrochlamydospora*, *P. quininea* and *P. richardiae*. These species have a few similar features, such as hypha swelling and unbranched sporangiophores; however, there are many differences. *Phytophthora docyniae*, *P. quininea*, and *P. richardiae* produce non-papillate sporangia, while *P. macrochlamydospora* has semi-papillate sporangia (Buisman 1927, Crandall 1947, Irwin 1991). The size of sporangia varies; the smallest belong to *P. docyniae*, followed by *P. macrochlamydospora*, *P. quininea* and then *P. richardiae*. Chlamydospores are absent in *P. richardiae* and very large for *P. macrochlamydospora*. Furthermore, a distinguishing feature of *P. docyniae* is that the chlamydospores frequently germinate and immediately produce sporangia and direct germination of sporangia with unusual shapes was often observed.



Maximum likelihood phylogeny produced using RAxML v. 8.2.11 (Stamatakis 2014) from concatenated ITS, BT and *cox1* using the GTR + I model showing the phylogenetic position of *Phytophthora docyniae* in clade 9c. Sequences are all available from IDPhy (Abad et al. 2019) and Bose et al. (2021). Bayesian analysis was conducted on the same dataset with MrBayes v. 3.2.6 (Ronquist & Huelsenbeck 2003) and resulted in the same topology. Numbers above the branches reflect support obtained from the analysis of the same dataset (bootstrap values estimated by RAxML/Bayesian posterior probabilities only values above 60 % or > 0.6 are shown). Representative species from other clades were used as outgroup taxa (not shown). The scale bar corresponds to expected number of substitutions per nucleotide site.

Colour illustrations. A flowering shoot of *Docynia indica*, the host of *Phytophthora docyniae*. Top to bottom: chlamydospores germinating to produce sporangia; chlamydospores and hyphal swelling with projections; chrysanthemum colony morphology on PDA; ovoid sporangia with direct germination; internal nested and extended proliferation; intercalary chlamydospore; thick-walled chlamydospore on PDA. Scale bars = $40 \mu\text{m}$.

Quynh. N. Dang, Thu. H. Nguyen, Hung. X. Tran & Thu. Q. Pham, Forest Protection Research Centre, Vietnamese Academy of Forest Sciences, 46 Duc Thang Ward, Bac Tu Liem District, Hanoi City, Vietnam;
e-mail: nhuquynh.dang@murdoch.edu.au, hoaithu.fsv@gmail.com, trahungfuv@gmail.com & phamquangthu@vafs.gov.vn
Treena I. Burgess, Phytophthora Science and Management, Harry Butler Institute, Murdoch University, 90 South Street, Murdoch, WA 6150, Australia; e-mail: tburgess@murdoch.edu.au

Pluteus brunneovenosus



Fungal Planet 1375 – 24 December 2021

***Pluteus brunneovenosus* Ferisin, Fellin, Justo & Dovana, sp. nov.**

Etymology. The epithet '*brunneovenosus*' refers to the colour of the cap with brown veins at the centre.

Classification — *Pluteaceae*, *Agaricales*, *Agaricomycetes*.

Basidiomata medium-sized, agaricoid. *Pileus* up to 35 mm diam, initially campanulate, then expanded to applanate, with low umbo at centre; surface glabrous, brown (Mu 10YR6/2) to dark brown (Mu 10YR3/2, 2/2) in centre pallescent towards the margin, hygrophanous, weakly to strongly venose at centre, strongly striate-sulcate at the margin up to the half of the radius. *Lamellae* fairly distant, free, slightly ventricose, up to 4 mm broad, first whitish later pink with whitish edge. *Stipe* 30–40 × 3–4 mm, cylindrical usually broadened towards base, striate lengthwise and flocculose, whitish in the upper or totally greyish, dark grey at the base. *Context* white, greyish in the stipe base. *Smell* and *taste* not distinctive. *Basidia* 26–30 × 8–10 µm, clavate, 4-spored. *Basidiospores* (6.5–)7.0–7.4–7.9(–8.5) × (5.0–)5.5–5.9–6.3(–7.0) µm, Q = (1.06–)1.21–1.26–1.35(–1.64); mainly broadly ellipsoid, more rarely ellipsoid, thick-walled, non-amyloid, cyanophilous. *Pleurocystidia* scattered and rare, similar to cheilocystidia. *Cheilocystidia* 30–140 × 15–31 µm, so numerous as to make the lamellar edge sterile; shape variable from clavate to utriform, fusiform, fusiform with rostrate apex, (rostrum length from 10 up to 35 µm), hyaline or with oily contents, slightly thick-walled. *Pileipellis* a trichohymeniderm made up of appressed clavate to fusiform elements 40–120 × 20–32 µm, sometimes with rostrate apex, (rostrum length up to 25 µm), pigment intracellular pale brown to brown. *Stipitipellis* a cutis of pale brown-grey hyphae with oily contents, 4–10 µm wide. *Caulocystidia* 40–60 × 18–22 µm, clavate to fusiform generally with rostrate apex (rostrum length up to 25 µm), generally in small clusters in the apical part of the stipe, rare towards the base, but present in all stipe length, hyaline or with oily contents. *Clamp connections* absent in all tissues.

Habitat & Distribution — Generally solitary or in small groups, on twigs of broadleaved trees (willow, oak, and poplar). Found, so far as we know, only in North Italy in Trentino Alto Adige and Friuli Venezia Giulia Region.

Typus. ITALY, Farra d'Isonzo, Bosco di Sotto, on twigs of broadleaved trees on the ground, 23 June 2020, G. Ferisin (holotype GDOR5090, ITS sequence GenBank OK090867, MycoBank MB 841180).

Additional materials examined. ITALY, Friuli-Venezia Giulia, Farra d'Isonzo, Bosco di Sotto, on twigs of broadleaved trees on the ground, 14 Oct. 2019, G. Ferisin, GDOR5091, ITS sequence GenBank OK090865; Trentino-Alto Adige, Ton, Masi di Vigo, on twigs on the ground, 11 Aug. 2019, A. Fellin, GDOR5089, ITS sequence GenBank OK090866.


Colour illustrations. Italy, Farra d'Isonzo, deciduous natural forest where the holotype collection was collected. Fresh basidiomes in field (holotype GDOR5090); pileipellis; stipitipellis with caulocystidia; cheilocystidia and basidiospores. Scale bars = 20 µm.

Notes — Terminology for descriptive terms is according to Vellinga (1988) and colour codes are taken from Munsell (1994). Spore dimensions are expressed as (a)–b–c–d(–e), where (a) = minimum value, b = (average - standard deviation), c = average, d = (average + standard deviation) and (e) = maximum value. Maximum-likelihood analysis of the ITS region was performed with RAXML v. 8.2.11 (Stamatakis 2014) using the GTR+G model as implemented in Geneious v. 11.1.5. *Pluteus brunneovenosus* is macroscopically characterised by its relatively small-sized basidiomes with pale brown to brown pileus, strongly veined in the centre; stipe cylindrical, pubescent, with grey to dark grey base; broadly ellipsoid spores; pileipellis a trichohymeniderm made up of clavate to fusiform elements sometime with rostrate apex, cheilocystidia utriform to broadly fusiform also with rostrate apex, pleurocystidia absent; caulocystidia present in all the stipe length. In the ITS phylogenetic tree, the three collections of *P. brunneovenosus* clustered in a fully supported clade (maximum likelihood bootstrap support value (ML-BS) = 100 %) which is sister (ML-BS = 95 %) to another clade that includes two sequences (GenBank FJ816665 of *P. dominicanus* var. *hyalinus* and GenBank KM983712, coll. AJ491, which is probably an undescribed species from Virgin Islands in the USA). *Pluteus dominicanus* var. *hyalinus* differs mainly for the very rare cheilocystidia, the absence of cystidia with rostrate apex and, for having an hymeniderm pileipellis made of sphaeropedunculate elements (Menolli et al. 2010). Phylogenetically, *P. brunneovenosus* is also close to sequences named *P. thomsonii* (GenBank HM562053 and HM562197) and *P. divarticulatus* (GenBank FJ375247). *Pluteus thomsonii* is very similar to *P. brunneovenosus* and can be distinguished on account of the more developed veins on the pileus and for the absence of rostrate terminal elements of the pileipellis (Vellinga 1990). *Pluteus divarticulatus*, originally described from France in 2003 and subsequently considered *Pluteus thomsonii* var. *divarticulatus* by Ludwig (2007) differs mainly by the presence of divarticulate caulocystidia, cheilocystidia and terminal elements of the pileipellis (Corriol 2003). Morphologically, *P. brunneovenosus* is also similar to *P. insidiosus*, that differs mainly for having a hymenidermal pileipellis made of sphaeropedunculate or narrowly clavate elements and the absence of caulocystidia (Vellinga 1990).

Supplementary material

FP1375 Phylogenetic tree.

Pythium serotinoosporum



Fungal Planet 1376 – 24 December 2021

***Pythium serotinoosporum* Abrinbana, Abdollahz. & Badali, sp. nov.**

Etymology. Name refers to the late production of oospores in culture.

Classification — *Pythiaceae*, *Peronosporales*, *Oomycota*, *Straminipila*.

Main *hyphae* up to 8 µm diam. *Sporangia* rarely produced zoospores through short discharge tubes, globose to subglobose, internally proliferating, terminal, occasionally intercalary, 15–33 µm diam (av. 25.21 µm diam). *Hyphal swellings* mostly germinating with 1–4 germ tubes, variously shaped, typically globose, subglobose, ellipsoidal, limoniform, ovoid, pyriform or irregular in shape, intercalary, occasionally terminal, occasionally in chains of 2–3, globose hyphal swellings 14–42 µm diam (av. 29.23 µm diam), elongated hyphal swellings 20–50 × 10–34 µm (av. 32.82 × 22.19 µm). Sexual organs produced infrequently after at least 3 wk in water culture. *Oogonia* smooth-walled, globose, subglobose, rarely elongated, terminal or intercalary, globose oogonia 17–29 µm diam (av. 22.65 µm diam), elongated oogonia up to 50 µm long. *Antheridia* 1–2(–7) per oogonium, diclinous. Antheridial cells globose, clavate, swollen, crooked or elongated. *Oospores* mainly globose, rarely subglobose or ovoid, plerotic or nearly plerotic, occasionally aplerotic, mostly one, but sometimes two per oogonium, 17–28 µm diam (av. 22.52 µm diam), the oospore wall up to 1 µm thick.


Culture characteristics — Colonies on potato carrot agar (PCA) submerged with a narrow chrysanthemum pattern, on cornmeal agar (CMA) submerged with a vague chrysanthemum pattern, on potato dextrose agar (PDA) profuse and dense aerial mycelium with no special pattern. Daily growth on PCA was 3 mm at 5 °C, 7 mm at 10 °C, 10 mm at 15 °C, 11 mm at 20 °C, 17 mm at 25 °C, 15 mm at 30 °C, 14 mm at 35 °C and 0 mm at 40 °C. The isolates were not viable at 40 °C.

Typus. IRAN, West Azarbaijan Province, Mahabad, from soil under *Prunus dulcis* (*Rosaceae*), 15 Aug. 2012, *M. Abrinbana* (holotype IRAN 16694 F, ex-type IRAN 2393 C, ITS, *COXI*, *COXII* and LSU sequences GenBank MG182707, MG182700, OK482595, OK428852, MycoBank MB 823188).

Additional material examined. IRAN, West Azarbaijan Province, Mahabad, from soil under *Prunus dulcis*, 15 Aug. 2012, *M. Abrinbana* (IRAN 2394 C = CBS 143583, ITS, *COXI*, *COXII* and LSU sequences GenBank MG182708, MG182701, OK482596, OK428853).

Colour illustrations. Almond orchard in West Azarbaijan Province. Various shapes of hyphal swellings; empty sporangium and zoospores; empty sporangium with internal proliferation; diclinous antheridium with crook-necked antheridial cell and oogonium; two diclinous antheridia applied to intercalary oogonium; six diclinous antheridia attached to oogonium; plerotic oospore; aplerotic oospores; double oospore; ovoid oospore. Scale bars = 10 µm.

Notes — Phylogenetic analyses of DNA sequence data supported the phylogenetic position of *P. serotinoosporum* in clade E2 sensu Lévesque & de Cock (2004). This species is indistinguishable from *P. middletonii* by ITS but, in the combined *COXI* and *COXII* phylogeny, *P. serotinoosporum* isolates form a fully-supported distinct clade sister to *P. middletonii* with 8 and 22 bp differences in *COXI* and *COXII* sequences, respectively. Morphologically, it differs from *P. middletonii* by having hyphal swellings, mostly plerotic oospores and a higher number of antheridia per oogonium (1–7 vs 1–2 per oogonium) and the absence of hypogynous as well as sessile antheridia.



The 50 % majority rule phylograms inferred from the Bayesian analysis of the ITS (a) and concatenated *COXI* and *COXII* (b) sequences alignment of *Pythium* clade E2, indicating the position of *P. serotinoosporum*. Bayesian posterior probability/maximum parsimony bootstrap support values are given at the nodes. Bayesian and maximum parsimony analyses were performed using MrBayes v. 3.2.7 (Ronquist et al. 2012) and PAUP v. 4.0b10 (Swofford 2003), respectively. The new species is indicated in **bold face** and a coloured block. *Pythium acanthophoron* (CBS 33729) from clade J was used as outgroup. † indicates ex-type strains. The scale bars represent 0.1 (a) and 0.01 (b) expected changes per site.

Masoud Abrinbana & Farzaneh Badali, Department of Plant Protection, Faculty of Agriculture, Urmia University, P.O. Box 165, Urmia, Iran;
e-mail: m.abrinbana@urmia.ac.ir & farzaneh.badali@gmail.com

Jafar Abdollahzadeh, Department of Plant Protection, Agriculture Faculty, University of Kurdistan, P.O. Box 416, Sanandaj, Iran;
e-mail: j.abdollahzadeh@uok.ac.ir

Thecaphora stajsicii



Fungal Planet 1377 – 24 December 2021

***Thecaphora stajscii* J. Kruse, R.G. Shivas & McTaggart, sp. nov.**


Etymology. In acknowledgement of Val Stajsic, an Australian botanist based at the National Herbarium of Victoria (MEL), who first recognised and collected smut fungi on *Oxalis* in Australia.

Classification — *Glomosporiaceae*, *Ustilaginales*, *Ustilaginomycetes*.

Sori in all or most capsules on an infected plant, replace the seeds with a powdery, cinnamon-brown spore mass causing the capsules to split longitudinally. **Spores** solitary when mature, globose to subglobose, 12–16 µm diam, pale to medium yellowish brown; wall even, 1–2 µm thick including moderately dense and coarse verruculose, warts c. 1 µm high. **Asexual morph** not seen.

Typus. AUSTRALIA, Australian Capital Territory, Canberra, Pialligo Redwood Forest, S35°19'12" E149°12'00", 595 m a.s.l., in capsules of *Oxalis radicata* (*Oxalidaceae*), 13 Nov. 2018, J. Kruse 945 (holotype BRIP 68991, isotype MEL 2496300A; ITS sequence GenBank MW892416, MycoBank MB 839329).

Notes — *Thecaphora stajscii* is the second species described on *Oxalis* in Australia. The first, *T. australiensis*, was described from *Oxalis exilis* in Victoria (Crous et al. 2018b). Capsules of *O. radicata* infected by *T. stajscii* are about half the length of healthy capsules. *Oxalis exilis* and *O. radicata* are both native species in Australia (Stajsic 2020a), where 30 species of *Oxalis* have been recorded, of which 23 are introduced (Stajsic 2020b). Worldwide, two other species, *T. oxalidis* and *T. capensis*, are known to infect *Oxalis* (Vánky et al. 2008). *Thecaphora stajscii* has smaller spores than *T. australiensis* (14–26 × 14–19 µm), and has spores similar in size to those of *T. oxalidis* (12–17 × 13.5–21(–24) µm) (Vánky 2011). The ITS sequence of *T. australiensis* (GenBank MW892417) was obtained from the ex-type specimen for the phylogenetic analysis. The four *Thecaphora* species on *Oxalis* form a monophyletic clade. *Thecaphora australiensis* and *T. stajscii* are closely related, which may indicate further biodiversity of host specific species on *Oxalis*.



Phylogram based on an alignment of the ITS region of rDNA. Tree searched in IQTree v. 2.1.3 (Minh et al. 2020) with a model test (command -m TEST), and support from 10 000 replicates of an approximate likelihood ratio test (≥ 0.9) and ultra-fast bootstraps ($\geq 95\%$) above nodes. Phylogenetic network made in SplitsTree v. 4.14.8 (Huson & Klopper 2005) showing genetic distance and putative recombination events among species of *Thecaphora*.

Colour illustrations. *Oxalis radicata* infected with *Thecaphora stajscii*, Canberra. Fungarium specimen; spores emerging from capsules; spores (equatorial and upper surface view). Scale bars = 1 cm (plant material), 10 µm (spores).

Julia Kruse, Pfalzmuseum für Naturkunde – POLLICHIA-Museum, Hermann-Schäfer-Str. 17, 67098 Bad Dürkheim, Germany;
e-mail: j.kruse@pfaizmuseum.bv-pfalz.de

Roger G. Shivas, Centre for Crop Health, University of Southern Queensland, Toowoomba 4350, Queensland, Australia;
e-mail: roger.shivas@usq.edu.au

Alistair R. McTaggart, Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Brisbane 4001, Australia;
e-mail: a.mctaggart@uq.edu.au

Trechispora patawaensis



Fungal Planet 1378 – 24 December 2021

***Trechispora patawaensis* V. Papp, Dima & L. Nagy, sp. nov.**

Etymology. Named after the place (Patawa camp) where the type specimen was collected.

Classification — *Hydnodontaceae*, *Trechisporales*, *Agaricomycetes*.

Basidiocarp annual, resupinate, soft and fragile, easily separable from the substrate; **margin** arachnoid, consisting of radiating hyphal mats with cord-like structures; **context** soft; **hymenophore** poroid, pore surface when fresh pure white, becoming pale yellowish when dried; the pores round to slightly irregular with thin dissepiments, small sized (6–7 per mm); **hyphal system** monomitic, thin-walled subicular hyphae hyaline, branched, with clamp connections and ampullate septate, up to 5 µm wide; Subhymenial hyphae thin-walled, branched, short-celled, isodiametric, up to 4 µm wide, with clamps in all septa. Single, rhomboid crystals abundantly present in subiculum; cystidia or other sterile hymenial elements not observed. **Basidia** short, cylindrical with four sterigmata and a basal clamp connection, 8–10 × 4–5 µm; **basidiospores** ovoid to subglobose (Qav = 1.36), hyaline, echinulate, (3.01–)3.07–3.35(–3.50) × (2.11–)2.21–2.44(–2.60) µm (excluding the ornamentation), L = 3.18 µm, W = 2.3 µm (n = 30), not cyanophilous.

(Line-drawn micromorphological structure of *T. patawaensis* basidiomata in FP1378 suppl. mat.)

Habitat & Distribution — The type was found in the Kaw Mts of French Guiana, on decayed dead branch of unknown angiosperm wood lying on the ground. A similar unpublished ITS sequence of a specimen collected in Ecuador (deposited at the UNITE database, no. UDB014091; Identities = 604/617 (98 %), seven gaps (1 %)), suggests that *T. patawaensis* presumably occurs throughout the Amazon basin.

Typus. FRENCH GUIANA, Kaw Mts, Camp Patawa, N4.54437° W52.15277°, on decayed branch of unknown angiosperm tree, 19 Dec. 2019, V. Papp & B. Dima (holotype WU-Mykol. 44791, WU); ITS and LSU sequences GenBank OL314550 and OL314546, MycoBank MB841876).

Colour illustrations. Camp Patawa, Kaw Mts, French Guiana (type locality). Basidiomata *in situ* (holotype). Scale bar = 1 mm.

Notes — In our phylogenetic analysis based on ITS and LSU sequences, *T. patawaensis* is located in a unique fully-supported clade (phylogenetic tree shown on FP1378 & 1379 suppl. mat.). In our phylogram, *T. patawaensis* clusters together with *T. farinacea* (MA-Fungi 79474, KHL 8454), *T. araneosa* (KHL 8570), *T. thelephora* (1984a AMV, MVL109) and *T. hymenocystis* (KHL 8795) in a well-supported clade. *Trechispora hymenocystis* has similar rhomboid crystals in the subiculum and cords but differs from *T. patawaensis* by its fairly large sphaerocysts on the subicular hyphae, smaller pores and larger basidiospores (Larsson 1994). The poroid *Trechispora* species formerly described from tropical regions, are easily distinguished from *T. patawaensis*, because *T. brasiliensis* has a dimitic hyphal system, *T. polygonospora* is characterised by angular and mainly hexagonal basidiospores, while *T. regularis* is well delimited by its cystidia-like incrustated hyphal ends (Larsson 1994). According to Hjortstam & Ryvarden (2007), three further poroid species originally described from Europe were reported from tropical areas: *T. candidissima*, *T. clancularis* and *T. mollusca*. The new species *T. patawaensis* morphologically differs from these species by the rhomboid crystal structures, smaller pores and smaller basidiospores. *Trechispora suberosa* was described from Guangxi (Southern China) and also has a poroid hymenophore, but differs from *T. patawaensis* by corky basidiocarps, thin- to slightly thick-walled generative hyphae without ampullate septa, and by lacking crystals (Yuan & Dai 2012).

Supplementary material

FP1378 Line-drawn micromorphological structure of *Trechispora patawaensis* basidiomata (WU-Mykol. 44791, type). a. Hyphae from subiculum; b. hyphae from cords; c. hyphae from dissepiment edge; d. subhymenium and hymenium; e. basidiospores. Scale bars = 10 µm (a–d), 5 µm (e).

FP1378 & 1379 Phylogenetic tree.

Viktor Papp, Department of Botany, Hungarian University of Agriculture and Life Sciences, Ménézi út 44. H-1118 Budapest, Hungary;

e-mail: agaricum@gmail.com

Bálint Dima, Department of Plant Anatomy, Institute of Biology, Eötvös Loránd University, Pázmány Péter sétány 1/C, H-1117, Budapest, Hungary;

e-mail: cortinarius1@gmail.com

László G. Nagy, Institute of Biochemistry, Biological Research Centre of the Eötvös Loránd Research Network, Temesvári blvd. 62, Szeged, H-6726, Hungary; e-mail: lnagy@funkenomelab.com



Fungal Planet 1379 – 24 December 2021

***Trechispora subregularis* V. Papp, Dima & L. Nagy, sp. nov.**

Etymology. The specific epithet '*subregularis*' refers to the morphological similarity to *Trechispora regularis*.

Classification — *Hydnodontaceae*, *Trechisporales*, *Agaricomycetes*.

Basidiocarp annual, effused, resupinate, soft and fragile, easily separable from the substrate, *margin* abrupt, slightly byssoid; *hymenophore* poroid, pores up to 0.5 mm long; pore surface whitish when young, turns pale yellowish in spots when dried; the pores angular, small sized, 6–7 per mm; *hyphal system* monomitic, thin-walled subicular hyphae hyaline, branched, with clamp connections and ampullate septate, up to 6 µm wide; crystals formed as jagged ridges along the hyphae (in subiculum and cords) and thick-walled cystidia-like hyphal segments. **Basidia** short, cylindrical with four sterigmata and a basal clamp connection, 9–11 × 4–5 µm; **basidiospores** ovoid to subglobose (Qav = 1.34), hyaline, echinulate, (3.10–)3.22–3.41(–3.45) × (2.32–)2.41–2.55(–2.62) µm (excluding the ornamentation), L = 3.26 µm, W = 2.5 µm (n = 30), not cyanophilous.

(Line-drawn micromorphological structure of *T. subregularis* basidiomata in FP1379 suppl. mat.)

Habitat & Distribution — Only known from the type locality in the Kaw Mts of French Guiana, tropical South America, on dead log of unknown angiosperm wood.

Typus. FRENCH GUIANA, Kaw Mts, near Camp Patawa, N4.55515° W52.14227°, on decayed log of unknown angiosperm tree, 21 Dec. 2019, V. Papp & B. Dima (holotype WU-Mykol. 44792, WU); ITS and LSU sequences GenBank OL331097 and OL314548, MycoBank MB 841878).

Colour illustrations. Fragment of tropical forest in Kaw Mts, French Guiana (type locality). Basidiomata *in situ* (holotype). Scale bar = 1 mm.

Notes — BLASTn searches based on ITS rDNA did not closely match with any known *Trechispora* species in the UNITE and GenBank databases. Our phylogenetic analysis based on ITS and LSU sequences (phylogenetic tree shown on FP1378 & 1379 suppl. mat.), the holotype of *T. subregularis* forms a unique lineage, while the collection of *T. regularis* (GenBank AF347087) from Jamaica is closely related. Amongst the eight poroid species currently accepted in *Trechispora*, only *T. regularis* has cystidia-like incrustated hyphal ends like *T. subregularis* (Larsson 1994, Yuan & Dai 2012). *Trechispora subregularis* is rather similar to *T. regularis*, but the latter has larger pores as 3–4 per mm (Larsson 1994), and is phylogenetically well delimited. Based on thorough morphological examinations, Larsson (1994) concluded, that the specimens of *T. regularis* are not homogenous, and the studied collections may even contain several species. Our recent findings supported this observation, and suggest that further studies are needed to clarify the taxonomic position of those species previously synonymised with *T. regularis* (Lowe 1966, Liberta 1973, Rajchenberg 1987, Meiras-Ottoni et al. 2021).

Supplementary material

FP1379 Line-drawn micromorphological structure of *Trechispora subregularis* basidiomata (WU-Mykol. 44792, type). a. Hyphae from subiculum; b. hyphae from cords; c. hyphae from dissepiment edge; d. cystidia-like hyphal segment; e. subhymenium and hymenium; f. basidiospores. Scale bars = 10 µm (a–e), 5 µm (f).

FP1378 & 1379 Phylogenetic tree.

Viktor Papp, Department of Botany, Hungarian University of Agriculture and Life Sciences, Ménézi út 44. H-1118 Budapest, Hungary;

e-mail: agaricum@gmail.com

Bálint Dima, Department of Plant Anatomy, Institute of Biology, Eötvös Loránd University, Pázmány Péter sétány 1/C, H-1117, Budapest, Hungary;

e-mail: cortinarius1@gmail.com

László G. Nagy, Institute of Biochemistry, Biological Research Centre of the Eötvös Loránd Research Network, Temesvári blvd. 62, Szeged,

H-6726, Hungary; e-mail: lnagy@funkenomelab.com



Fungal Planet 1380 – 24 December 2021

Tuber suaveolens Ant. Rodr. & Morte, sp. nov.

Etymology. Referring to its mild odour.

Classification — *Tuberaceae*, *Pezizales*, *Pezizomycetes*.


Ascomata hypogeous, 1–3 cm in size, subglobose, sometimes with a rudimentary basal cavity, covered with brown-black pyramidal warts, 4–6-sided, 2–4(–5) mm across, 1–3 mm high, often depressed at the apex. *Peridium* 150–300 µm thick, pseudoparenchymatous, composed of subglobose, angular cells, 10–20 µm diam, dark red-brown and thickened walls in the outermost layers, pale yellow and thin-walled in the innermost layers, merging with interwoven hyphae of glebal tissue. *Gleba* firm, solid, white when immature, becoming dark brown at maturity, marbled with numerous, thin, white, meandering veins. Mild phenolic *odour*. Bitter *taste*. *Asci* inamyloid, 50–90 × 50–70 µm, walls thickened, 1–2 µm, ellipsoid to subglobose, with a short stalk, 10–30 × 5–7 µm, 1–5(–6)-spored. *Ascospores* 25–55 × 25–37 µm, Q = 1.1–1.4, excluding ornamentation, yellowish, ellipsoid to subglobose, ornamented with a coarse irregular reticulum, 3–5 µm high, sometimes bending at the top. Meshes variable, usually 3–5 across width of spore and often with incomplete secondary crests inside.

Ecology & Distribution — *Tuber suaveolens* grows in calcareous soil of limestone mountains of the southeast of the Iberian Peninsula, associated to *Quercus* spp. in winter.

Typus. SPAIN, Albacete, Nerpio, in calcareous soil, under *Quercus faginea* (*Fagaceae*), 23 Jan. 2008, A. Rodríguez (holotype MUB Fung-931; ITS and LSU sequences GenBank MZ616376 and MZ618623, MycoBank MB 840650).

Additional material examined. SPAIN, Albacete, Nerpio, under *Quercus ilex* subsp. *ballota*, 11 Feb. 2009, A. Rodríguez, MUB Fung-1013; ITS sequence GenBank MZ616377.


Notes — *Tuber suaveolens* is a black truffle of the *aestivum* clade characterised by its brown-black warty peridium, brown gleba marbled with thin white veins, mild odour and reticulate-alveolate spores. It resembles *T. mesentericum*, but in addition to genetic differences, it differs from *T. mesentericum* (Vittadini 1831) by having a mild phenolic odour, rudimentary or absent basal cavity and different vein arrangement. It also resembles *T. alcaracense*, but *T. alcaracense* has a pleasant odour and lacking a basal cavity (Crous et al. 2020b).



Colour illustrations. Spain, Nerpio (Albacete), wild truffière of *Quercus faginea*. Ascocarp; mature ascospores. Scale bar = 20 µm.

Antonio Rodríguez, Alfonso Navarro-Ródenas, Francisco Arenas, Angel Luigi Guarnizo & Asunción Morte, Departamento de Biología Vegetal (Botánica), Facultad de Biología, Universidad de Murcia, 30100 Murcia, Spain; e-mail: antonio@trufamania.com, anr@um.es, f.arenasjimenez@um.es, angelluigi.guarnizo@um.es & amorte@um.es

Variabilispora viridis



Fungal Planet 1381 – 24 December 2021

Variabilispora viridis V.A. Iliushin, I.Y. Kirtsideli & E.G. Lukina, *sp. nov.**Etymology.* Named after its green (Lat.: *viridis*) colonies.*Classification* — *Tympanidaceae*, *Helotiales*, *Leotiomyces*.


Vegetative hyphae hyaline, smooth-walled, septate, branched, 1.5–3.8 µm wide. Sporulation abundant, conidia formed on hyphal cells and by microcyclic conidiation. *Conidiophores on hyphae* reduced to conidiogenous cells. *Conidiogenous cells* enteroblastic, hyaline, smooth-walled, terminally and intercalary, reduced to short necks or collarettes formed directly on hyphal cells; discrete phialides often observed, cylindrical to ampulliform, 5–10 × 1.5–3.5 µm; collarettes cylindrical, thin-walled, inconspicuous, 0.5–1 µm long, opening 0.5–1.2 µm wide; necks cylindrical, 0.5–2.5 µm long, 0.5–1.5 µm wide. *Conidia* aggregated in masses around the hyphae, hyaline, smooth-walled, aseptate, subglobose, ellipsoidal, obovate, 7–10.5 × 2.5–5 µm. *Conidiomata* and *endoconidia* not observed. *Microcyclic conidiation* occurs from collarettes at one or sometimes both ends of conidia that have developed into mother cells, often thick-walled, sometimes septate, > 8 µm long, 3.5–5.5 µm wide.

Culture characteristics — Colonies on Czapek agar (CZ) after 4 wk at 23 °C reaching 20–29 mm diam, flat to low convex with dentate to fimbriate margin, moist, lacking aerial mycelium; zonate, successively deep olive green (#232f00; ISCC-NBS Centroid Color Charts) and moderate olive green (#4a5d23) from the centre towards the edge; reverse deep olive green (#232f00). Colonies on malt extract agar (MEA) after 4 wk at 23 °C reaching 10–20 mm diam, moist, lacking aerial mycelium;

non-zonate, deep olive green (#232f00); reverse same colour. Colonies on oatmeal agar (OA) after 4 wk at 23 °C reaching 20–32 mm diam, flat, moist, lacking aerial mycelium; successively moderate olive green (#4a5d23) and light greenish yellow (#eae679) from the centre towards the edge; reverse same colour. Minimum temperature for growth 8 °C, optimum 23 °C, maximum 30 °C.

Typus. NORWAY, Svalbard Archipelago, coastal area of the Grunfjord of the Greenland Sea, from driftwood of *Picea abies* (*Pinaceae*), 2020, E.G. Lukina (holotype LE F-341002, culture ex-type CBS 148030, ITS and LSU sequences GenBank MW201500.1 and MZ798432.1, MycoBank MB 840872).

Notes — Bien et al. (2020) introduced a new genus *Variabilispora* for collophorina-like fungi. Phylogenetic analyses show *V. viridis* as the closest species to *V. flava*, but several morphological differences were found. Morphologically, *V. viridis* is different from *V. flava* in size of the phialides (2–9 × 1.5–2.5 µm vs 5–10 × 1.5–3.5 µm in *V. viridis*), conidia (2.5–6.5 × 1.5–2 µm vs 7–10.5 × 2.5–5 µm in *V. viridis*), and mother cells of microcyclic conidiation (> 5 × 2.5–3.5 µm vs > 8 × 3.5–5.5 µm in *V. viridis*). Furthermore, *V. viridis* also differs from *V. flava* by its moderate olive green to light greenish yellow colour on OA (*V. flava* is sulphur yellow to pure yellow). Based on a megablast search of NCBI's GenBank nucleotide database, the closest hit using the ITS sequence is with the ex-type strain of *V. flava* (culture CBS 144845, GenBank NR_165906.1; Identities = 460/477 (96 %), two gaps (0 %)); and using the LSU sequence it is with the same strain of *V. flava* (GenBank NG_068608.1; Identities = 526/542 (97 %), no gaps).



Colour illustrations. Svalbard Archipelago, coastal area of the Grunfjord of the Greenland Sea with driftwoods. Colony on CZ and OA after 4 wk at 23 °C; conidiogenous cells formed on hyphal cells; conidia; mother cell. Scale bars = 20 µm.



Fungal Planet 1382 – 24 December 2021

Microdochium maculosum A.P. de Souza, B.W. Ferreira, R.W. Barreto & B.S. Vieira, *sp. nov.**Etymology.* Referring to leaf spots (maculas) caused by the fungus.Classification — *Microdochiaceae*, *Xylariales*, *Sordariomycetes*.

Lesion on living leaves, starting as small necrotic dots, becoming irregular, 5–50 mm diam, pale brown, with dark brown margins, surrounded by a yellow halo, coalescing and leading to extensive leaf blight. *Mycelium* superficial and immersed; hyaline to pale brown, branched, septate, hyphae 1–4 µm diam. *Conidiophores* cylindrical to filiform, 7–55 × 1–3 µm, sometimes reduced to conidiogenous cells borne directly from the hyphae. *Conidiogenous cells* terminal or intercalary, holoblastic, mono- or polyblastic, cylindrical to filiform, denticulate, sympodial, 7–39 × 1–3 µm, hyaline, smooth. *Conidia* fusoid, straight or curved, 6–15 × 2–4 µm, 1–3-septate, hyaline, occasionally curved at the tip, base tapering to a distinct hilum, 1–2 × 1–2 µm. *Sexual morph* not observed. *Chlamydospores* not observed.


Culture characteristics — Reaching up to 55 mm diam on oatmeal agar (OA) and 50 mm diam on malt extract agar (MEA), after 7 d at 25 °C. Colony flat, edge entire, aerial mycelium dense and felt-like; on OA, centre olivaceous to grey olivaceous and white toward the periphery; on MEA, centre flesh to rosy buff, margins, reverse rosy buff. Sporulation abundant on OA.

Typus. BRAZIL, Minas Gerais, Viçosa, Floricultura, campus of the Universidade Federal de Viçosa, on living leaves of *Digitaria insularis* (*Poaceae*), 21 May 2020, R.W. Barreto (holotype VIC 47532, culture ex-type COAD 3358, ITS, LSU and *rpb2* sequences GenBank OK966953, OK966954 and OL310501, MycoBank MB 841887).

Notes — Based on a megablast search of NCBI's GenBank nucleotide database, the LSU and *rpb2* sequences of *Microdochium maculosum* showed a similarity of 99.77 % and 91.31 % respectively, with the sequences of *Microdochium indocalami* (SAUCC1016, GenBank MT199878 and MT510550); whereas the ITS sequence was 98.61 % similar to that of *Microdochium* sp. (JP103, GenBank AB255278). *Microdochium maculosum* had a similarity of 97.62 % with ITS sequence of *Microdochium fisheri* strain CBS 242.91 (GenBank NR_155374), 98.70 % similarity with LSU and 89.29 % with *rpb2* of *M. fisheri* strain CBS 242.90, (GenBank KP858951 and KP859124).

Colour illustrations. *Microdochium maculosum* causing leaf spots in *Digitaria insularis*. Conidiophores, with denticulate conidiogenous cells; fusiform to allantoid conidia; colonies in OA and MEA media. Scale bars = 10 µm.

The phylogenetic reconstruction using ITS, LSU and *rpb2* of accepted *Microdochium* species, showed that the new species was located in an highly-supported clade together with *M. indocalami*, *M. fisheri*, *M. lycopodium*, *M. phragmitis*, and *M. rhopalostylidis*, being closely related with the former two species. However, *M. maculosum* has clear morphological differences distinguishing it from *M. fisheri* and *M. indocalami*. *Microdochium fisheri* has fusoid, obovoid, subpyriform to clavate (7–12 × 3–4 µm, 0–1-septate) conidia (Hernández-Restrepo et al. 2016a), and *M. indocalami* has conidia which are cylindrical, clavate to obovoid (13–15.5 × 3.5–5.5 µm, 1–3-septate) and have a flattened base and no distinct hila (Huang et al. 2020). On the other hand, the new species *M. maculosum* has fusoid, straight or curved conidia (6–15 × 2–4 µm, 1–3-septate). *Microdochium fisheri* is known from *Oryza sativa* in India and the UK, and *M. indocalami* is known from *Indocalamus longiauritus* in China (Farr & Rossman 2021). No species of *Microdochium* have ever been reported on members of *Digitaria* prior to the present work.



Bayesian tree inferred from the combined datasets of ITS, LSU and *rpb2* sequences from species belonging to the genus *Microdochium*, including the specimen of *Microdochium maculosum* obtained in this study (indicated in **bold**). The analysis was performed with 10 M generations in MrBayes v. 3.2.6 (Ronquist et al. 2012). Bayesian probabilities > 0.50 are given at each node. The tree is rooted to *Idriella lunata* CBS 204.56. GenBank accession numbers of the sequences used in this study are from Huang et al. (2020).

REFERENCES

- Abad ZG, Burgess TI, Bienapfl JC, et al. 2019. IDphy: Molecular and morphological identification of Phytophthora based on the types. USDAAPHIS PPQ S&T Beltsville Lab, USDA APHIS PPQ S&T ITP, Centre for Phytophthora Science and Management, and World Phytophthora Collection.
- Abdollahzadeh J, Groenewald JZ, Coetzee MPA, et al. 2020. Evolution of lifestyles in Capnodiales. *Studies in Mycology* 95: 381–414.
- Alvarado P, Cabero J, Moreno G, et al. 2014. Species diversity of *Genea* (Ascomycota, Pezizales) in Europe. *Ascomycete.org* 6: 41–51.
- Alvarado P, Cabero J, Moreno G, et al. 2016. Phylogenetic overview of the genus *Genea* (Pezizales, Ascomycota) with an emphasis on European taxa. *Mycologia* 108: 441–456.
- Alvarado P, Pérez J-B, Van Vooren N, et al. 2020. *Genea coronata* (Pyrenomataceae, Pezizales), a cryptic new species in a highly polymorphic genus. *Sydowia* 72: 231–242.
- Anonymous. 1969. Flora of British Fungi – Colour Identification Chart. Her Majesty's Stationery Office, Edinburgh.
- Antonin V, Noordeloos M. 2010. A monograph of marasmioid and collybioid fungi in Europe. IHW Verlag.
- Arzanlou A, Groenewald JZ, Gams W, et al. 2007. Phylogenetic and morphotaxonomic revision of *Ramichloridium* and allied genera. *Studies in Mycology* 58: 57–93.
- Aveskamp MM, Verkley GJM, De Gruyter J, et al. 2009. DNA phylogeny reveals polyphyly of *Phoma* section *Peyronellaea* and multiple taxonomic novelties. *Mycologia* 101: 363–382.
- Bandini D, Oertel B, Ploch S, et al. 2019. Revision of some central European species of *Inocybe* (Fr.: Fr.) Fr. subgenus *Inocybe*, with description of five new species. *Mycological Progress* 18: 247–294.
- Bandini D, Sesli E, Oertel B, et al. 2020. *Inocybe antoniniana*, a new species of *Inocybe* section *Marginatae* with nodulose spores. *Sydowia* 72: 95–106.
- Bao DF, Wanasinghe DN, Luo ZL, et al. 2019. *Murispora aquatica* sp. nov. and *Murispora fagicola*, a new record from freshwater habitat in China. *Phytotaxa* 416: 1–13.
- Baral H-O. 1987. Lugol's solution / IKI versus Melzer's reagent: hemiamyloidity, a universal feature of the ascus wall. *Mycotaxon* 29: 399–450.
- Bates ST, Roberson RW, Desjardin DE. 2009. *Arizona gasteroid fungi I: Lycoperdaceae* (Agaricales, Basidiomycota). *Fungal Diversity* 37: 153–207.
- Bensch K, Groenewald JZ, Braun U, et al. 2015. Common but different: The expanding realm of *Cladosporium*. *Studies in Mycology* 82: 23–74.
- Bensch K, Groenewald JZ, Meijer M, et al. 2018. *Cladosporium* species in indoor environments. *Studies in Mycology* 89: 177–301.
- Berkeley MJ. 1845. Decades of Fungi: Australian Fungi. *The London Journal of Botany* 4: 42–73.
- Bertault R. 1965. *Amanites du Maroc*. Bulletin trimestriel de la Société mycologique de France 81: 345–371.
- Bezerra JDP, Sandoval-Denis M, Paiva LM, et al. 2017. New endophytic *Toxicocladosporium* species from cacti in Brazil, and description of *Neocladosporium* gen. nov. *IMA Fungus* 8: 77–97.
- Bien S, Kraus C, Damm U. 2020. Novel collophorina-like genera and species from *Prunus* trees and vineyards in Germany. *Persoonia* 45: 46–67.
- Boerema GH, De Gruyter J, Noordeloos ME, et al. 2004. *Phoma* identification manual. Differentiation of specific and intra-specific and intra-specific taxa in culture. CABI Publishing, Wallingford, UK.
- Bose T, Wingfield MJ, Roux J, et al. 2021. *Phytophthora* species associated with roots of native and non-native trees in natural and managed forests. *Microbial Ecology* 81: 122–133.
- Braun U. 1998. A monograph of *Cercosporiella*, *Ramularia* and allied genera (phytopathogenic hyphomycetes), Vol. 2, IHW-Verlag, Eching, Munich, Germany.
- Buisman CJ. 1927. Root rots caused by Phycomycetes. Mededelingen uit het Phytopathologisch laboratorium Willie Commelin Scholten 11: 1–65.
- Bukharova NV, Zmitrovich IV. 2014. Aphylloroid fungi of the Bastak Reserve. *Mikologiya i fitopatologiya* 48: 343–354.
- Castañeda-Ruiz RF. 1984. Nuevos taxones de Deuteromycotina: *Arnoldiella robusta* gen. et sp. nov.; *Roigiella lignicola* gen. et sp. nov.; *Sporidesmium pseudolmediae* sp. nov. y *Thozetella havanensis* sp. nov. *Revista del Jardín Botánico Nacional Universidad de la Habana* 5: 57–87.
- Castañeda-Ruiz RF, Gams W, Saikawa M. 1997. Three new conidial fungi (hyphomycetes) from Cuba. *Nova Hedwigia* 64: 473–483.
- Chen C, Verkley GJM, Sun G, et al. 2016. Redefining common endophytes and plant pathogens in *Neofabraea*, *Pezizula*, and related genera. *Fungal Biology* 120: 1291–1322.
- Chen Q, Hou LW, Duan WJ, et al. 2017. *Didymellaceae* revisited. *Studies in Mycology* 87: 105–159.
- Chen Q, Jiang JR, Zhang GZ, et al. 2015. Resolving the *Phoma* enigma. *Studies in Mycology* 82: 137–217.
- Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Systematic Biology* 65: 997–1008.
- Contu M. 1985a '1984'. *Novitates* (2). *Documents Mycologiques* 14: 26.
- Contu M. 1985b. *Amanita mairei* Foley sensu Foley. A. *mairei* sensu Bertault: Nome nuovo per la specie di Bertault: *Amanita bertaultii* sp. nov. *Funghi Ambiente* 1: 11–13.
- Contu M. 2000. Chiave per la determinazione delle specie europee del genere *Amanita* sez. *Vaginatae*. *Bollettino del gruppo micologico G. Bresadola* 43: 233–240.
- Contu M, Pacioni G. 1998. *Amanita cistetorum* and *Psathyrella liciosae*, two new Mediterranean species. *Mycotaxon* 69: 437–446.
- Corriol G. 2003. *Pluteus diverticulatus* sp. nov., une espèce nouvelle de la sous-section *Mixtini*. *Bulletin Trimestriel de la Société Mycologique de France* 119: 231–244.
- Corriol G, Moreau PA, Bellanger JM. 2017. *Calocybe hymenoderma* sp. nov., *Calocybella juncicola* comb. nov. et les contours taxinomiques de la section *Rugosomyces*. *Errotari* 14: 35–46.
- Crandall BS. 1947. A new *Phytophthora* causing root and collar rot of cinchona in Peru. *Mycologia* 39: 218–223.
- Crous PW, Braun U, McDonald BA, et al. 2020a. Redefining genera of cereal pathogens: *Oculimacula*, *Rhynchosporium* and *Spermospora*. *Fungal Systematics and Evolution* 7: 67–98.
- Crous PW, Braun U, Wingfield MJ, et al. 2009. Phylogeny and taxonomy of obscure genera of microfungi. *Persoonia* 22: 139–161.
- Crous PW, Carnegie AJ, Wingfield MJ, et al. 2019a. Fungal Planet description sheets: 868–950. *Persoonia* 42: 291–473.
- Crous PW, Cowan DA, Maggs-Kölling G, et al. 2021a. Fungal Planet description sheets: 1182–1283. *Persoonia* 46: 313–528.
- Crous PW, Groenewald JZ, Mansilla JP, et al. 2004. Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus*. *Studies in Mycology* 50: 195–214.
- Crous PW, Lombard L, Sandoval-Denis M, et al. 2021b. *Fusarium*: more than a node or a foot-shaped basal cell. *Studies in Mycology* 98: 100116.
- Crous PW, Luangsa-ard JJ, Wingfield MJ, et al. 2018a. Fungal Planet description sheets: 785–867. *Persoonia* 41: 238–417.
- Crous PW, Schumacher RK, Akulov A, et al. 2019b. New and Interesting Fungi. 2. *Fungal Systematics and Evolution* 3: 57–134.
- Crous PW, Schumacher RK, Wingfield MJ, et al. 2018b. New and interesting fungi. 1. *Fungal Systematics and Evolution* 1: 169–215.
- Crous PW, Shivas RG, Quaedvlieg W, et al. 2014a. Fungal Planet description sheets: 214–280. *Persoonia* 32: 184–306.
- Crous PW, Summerell BA, Shivas RG, et al. 2011. Fungal Planet description sheets: 92–106. *Persoonia* 27: 130–162.
- Crous PW, Verkley GJM, Christensen M, et al. 2012. How important are conidial appendages? *Persoonia* 28: 126–137.
- Crous PW, Wingfield MJ, Burgess TI, et al. 2017. Fungal Planet description sheets: 625–715. *Persoonia* 39: 270–467.
- Crous PW, Wingfield MJ, Cheewangkoon R, et al. 2019c. Foliar pathogens of eucalypts. *Studies in Mycology* 94: 125–298.
- Crous PW, Wingfield MJ, Chooi Y-H, et al. 2020b. Fungal Planet description sheets: 1042–1111. *Persoonia* 44: 301–459.
- Crous PW, Wingfield MJ, le Roux JJ, et al. 2015. Fungal Planet description sheets: 371–399. *Persoonia* 35: 264–327.
- Crous PW, Wingfield MJ, Lombard L, et al. 2019d. Fungal Planet description sheets: 951–1041. *Persoonia* 43: 223–425.
- Crous PW, Wingfield MJ, Richardson DM, et al. 2016. Fungal Planet description sheets: 400–468. *Persoonia* 36: 316–458.
- Crous PW, Wingfield MJ, Schumacher RK, et al. 2014b. Fungal Planet description sheets 281–319. *Persoonia* 33: 212–289.
- Crous PW, Wingfield MJ, Schumacher RK, et al. 2020c. New and interesting fungi 3. *Fungal Systematics and Evolution* 6: 157–231.
- Darriba D, Taboada GL, Doallo R, et al. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 772.
- De Gruyter J, Noordeloos ME. 1992. Contributions towards a monograph of *Phoma* (Coelomycetes) – I. 1. Section *Phoma*: Taxa with very small conidia in vitro. *Persoonia* 15: 71–92.
- De Gruyter J, Noordeloos ME, Boerema GH. 1993. Contributions towards a monograph of *Phoma* (Coelomycetes) – I. 2. Section *Phoma*: Additional taxa with very small conidia and taxa with conidia up to 7 µm long. *Persoonia* 15: 369–400.
- De Hoog GS. 1977. *Rhinocladiella* and allied genera. *Studies in Mycology* 15: 1–140.
- De Hoog GS, Mayser P, Haase G, et al. 2000. A new species, *Phialophora europaea*, causing superficial infections in humans. *Mycoses* 43: 409–416.

- Denk T, Grimm GW. 2010. The oaks of western Eurasia: Traditional classifications and evidence from two nuclear markers. *Taxon* 59: 351–366.
- Diederich P, Braun U. 2009. First lichenicolous record of *Acremonium hyphomatis* (anamorphic Ascomycota). *Bulletin de la Société des naturalistes luxembourgeois* 110: 97–99.
- Domsch KH, Gams W, Anderson TH. 1993. *Compendium of soil fungi*. Reprinted by IHW-Verlag with supplement.
- Domsch KH, Gams W, Anderson T-H. 2007. *Compendium of soil fungi*. 2nd ed. IHW-Verlag, Eching, Germany.
- Dong W, Hyde KD, Jeewon R, et al. 2021. Towards a natural classification of annulatascaceae-like taxa II: introducing five new genera and eighteen new species from freshwater. *Mycosphere* 12: 1–88.
- Drummond AJ, Suchard MA, Xie D, et al. 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* 29: 1969–1973.
- Duan JX, Wu WP, Liu XZ. 2007. *Dinemasporium* (coelomycetes). *Fungal Diversity* 26: 205–218.
- Edler D, Klein J, Antonelli A, et al. 2020. raxmlGUI 2.0: a graphical interface and toolkit for phylogenetic analyses using RAXML. *Methods in Ecology and Evolution* 12: 373–377.
- Egidi E, De Hoog GS, Isola D, et al. 2014. Phylogeny and taxonomy of meristematic rock-inhabiting black fungi in the Dothideomycetes based on multi-locus phylogenies. *Fungal Diversity* 65: 127–165.
- Ekanayaka AH, Hyde KD, Gentekaki E, et al. 2019. Preliminary classification of Leotiomycetes. *Mycosphere* 10: 310–489.
- Esteve-Raventós F, Moreno G, Bizio E, et al. 2015. *Inocybe flavobrunnescens*, a new species in section *Marginatae*. *Mycological Progress* 14: 14.
- Farr DF, Rossman AY. *Fungal Databases*. U.S. National Fungus Collections, ARS, USDA. Retrieved October 14, 2021, from <https://nt.ars-grin.gov/fungaldatabases/>.
- Fassatiová O. 1986. Moulds and filamentous fungi in technical microbiology. Elsevier, Amsterdam.
- Fechner N, Bonito G, Bougher NL, et al. 2017. New species of *Austroboletus* (Boletaceae) in Australia. *Mycological Progress* 16: 769–775.
- Feng P, Lu Q, Najafzadeh MJ, et al. 2014. Cyphellophora and its relatives in Phialophora biodiversity and possible role in human infection. *Fungal Diversity* 65: 17–45.
- Fries E. 1821. *Systema mycologicum* 1. Ex officina Berlingiana, Lund, Sweden.
- Gams W. 1971. *Cephalosporium-artige Schimmelpilze* (Hyphomycetes), Fischer, Stuttgart, Germany.
- Gams W. 1975. *Cephalosporium-like Hyphomycetes: Some tropical species*. *Transactions of the British Mycological Society* 64: 389–404.
- Gams W. 1976. Some new or noteworthy species of *Mortierella*. *Persoonia* 9: 111–140.
- Giraldo A, Gené J, Sutton DA, et al. 2015. Phylogeny of *Sarocladium* (Hypocreales). *Persoonia* 34: 10–24.
- Gonçalves MFM, Santos L, Silva BMV, et al. 2019. Biodiversity of *Penicillium* species from marine environments in Portugal and description of *Penicillium lusitanum* sp. nov., a novel species isolated from sea water. *International Journal of Systematic and Evolutionary Microbiology* 69: 3014–3021.
- Gouy M, Guindon S, Gascuel O. 2010. SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution* 27: 221–224.
- Gräfenhan T, Schroers H-J, Nirenberg HI, et al. 2011. An overview of the taxonomy, phylogeny, and typification of nectriaceous fungi in *Cosmospora*, *Acremonium*, *Fusarium*, *Stilbella*, and *Volutella*. *Studies in Mycology* 68: 79–113.
- Gryzenhout M, Tarigan M, Clegg PA, et al. 2010. *Cryptometron aestuescens* gen. sp. nov. (Cryphonectriaceae) pathogenic to *Eucalyptus* in Indonesia. *Australasian Plant Pathology* 39: 161–169.
- Guindon S, Lethiec F, Duroux P et al. 2010. PHYML Online – a web server for fast maximum likelihood-based phylogenetic inference. *Nucleic Acids Research* 33: W557–W559.
- Guzmán-Dávalos L, Pradeep CK, Vrinda KB, et al. 2017. A new stipitate species of *Crepidotus* from India and Thailand, with notes on other tropical species. *Mycologia* 109: 804–814.
- Haelewaters D, De Kesel A. 2020. Checklist of thallus-forming Laboulbeniomyces from Belgium and the Netherlands, including *Hesperomyces halyziae* and *Laboulbenia quarantanae* spp. nov. *MycKeys* 71: 23–86.
- Haelewaters D, De Kesel A, Pfister DH. 2018. Integrative taxonomy reveals hidden species within a common fungal parasite of ladybirds. *Scientific Reports* 8: 15966.
- Hall TA. 1999. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Halling RE. 1996. Notes on *Collybia* V. *Gymnopus* section *Levipedes* in tropical South America, with comments on *Collybia*. *Brittonia* 48: 487–494.
- Hanss J-M, Moreau P-A. 2020. Une Révision des Amanites «vaginées» (*Amanita* sect. *Vaginatae*) en Europe, 1re partie: quelques Amanites argentées. *Bulletin de la Société mycologique de France* 133: 67–141.
- Hashimoto A, Sato G, Matsuda T, et al. 2015. Taxonomic revision of *Pseudodolachnea* and *Pseudodolachnella* and establishment of *Neopseudodolachnella* and *Pseudodinemasporium* gen. nov. *Mycologia* 107: 383–408.
- Hawksworth DL. 1972. A large-spored species of *Acremonium* sect. *Nectrioidea*. *Transactions of the British Mycological Society* 58: 510–512.
- Hawksworth DL. 1979. The lichenicolous hyphomycetes. *Bulletin of the British Museum for Natural History* 6: 183–300.
- He M-Q, Hyde KD, Ratchadawan C, et al. 2020. Two new species of *Microspalliota* (Agaricaceae/Agaricales) from Thailand. *Phytotaxa* 453: 140–141.
- Heinemann P, Leelavathy KM. 1991. The genus *Microspalliota* (Agaricaceae) in Kerala State, India. *Mycological Research* 95: 343–344.
- Heinemann P, Little Flower S. 1983. *Microspalliota* de Kerala (Inde). *Bulletin du Jardin Botanique National de Belgique* 53: 75–84.
- Hernández-Restrepo M, Groenewald JZ, Crous PW, et al. 2016a. Taxonomic and phylogenetic re-evaluation of *Microdochium*, *Monographella* and *Idriella*. *Persoonia* 36: 57–82.
- Hernández-Restrepo M, Groenewald JZ, Elliott ML, et al. 2016b. Take-all or nothing. *Studies in Mycology* 83: 19–48.
- Hipp AL, Manos PS, Hahn M, et al. 2020. Genomic landscape of the global oak phylogeny. *New Phytologist* 226: 1198–1212.
- Hjortstam K, Ryvarden L. 2007. Checklist of corticioid fungi (Basidiomycotina) from the tropics, subtropics and the southern hemisphere. *Synopsis Fungorum* 22: 27–146.
- Hoang DT, Chernomor O, von Haeseler A, et al. 2018. UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* 35: 518–522.
- Holubová-Jechová V. 1974. A revision of the genus *Olpitrichum* Atk. *Folia Geobotanica et Phytotaxonomica* 9: 425–432.
- Hongo T. 1971. *Notulae mycologicae*. *Mem Shiga University* 21: 62–68.
- Horak E. 1976. On cuboid spored species of *Entoloma*. *Sydowia* 28: 171–236.
- Horak E. 1977. Additions to 'On cuboid spored species of *Entoloma*'. *Sydowia* 29: 289–299.
- Horak E. 1978 '1977'. *Crepidotus episphaeria* and related species from the Southern Hemisphere. *Berichte der Schweizerischen Botanischen Gesellschaft* 87: 227–235.
- Horak E. 2018. Agaricales (Basidiomycota) of New Zealand 2. Brown spored genera p.p. *Crepidotus*, *Flammulaster*, *Inocybe*, *Phaeocollybia*, *Phaeomarasmius*, *Pleuroflammula*, *Pyrrhoglossum*, *Simocybe*, *Tubaria* and *Tympanella*. *Fungi of New Zealand Nga Hekaheka o Aotearoa Volume 6*. *Westerdijk Biodiversity Series* 16: 1–205. *Westerdijk Fungal Biodiversity Institute*, Utrecht, the Netherlands.
- Hou C-L, Lin Y-R, Piepenbring M. 2005. Species of *Rhytismataceae* on needles of *Juniperus* spp. from China. *Canadian Journal of Botany* 83: 37–46.
- Hou LW, Groenewald JZ, Pfenning LH, et al. 2020a. The phoma-like dilemma. *Studies in Mycology* 96: 309–396.
- Hou LW, Hernandez-Restrepo M, Groenewald JZ, et al. 2020b. Citizen science project reveals high diversity in Didymellaceae (Pleosporales, Dothideomycetes). *MycKeys* 65: 49–99.
- Houbraken J, Kocube S, Visagie CM, et al. 2020. Classification of *Aspergillus*, *Penicillium*, *Talaromyces* and related genera (Eurotiales): an overview of families, genera, subgenera, sections, series and species. *Studies in Mycology* 95: 5–169.
- Huang S, Xia J, Zhang X, et al. 2020. Two new species of *Microdochium* from *Indocalamus longiauritus* in south-western China. *MycKeys* 72: 93–108.
- Hubka V, Dudová Z, Kubátová A, et al. 2017. Taxonomic novelties in *Aspergillus* section *Fumigati*: *A. tasmanicus* sp. nov., induction of sexual stage in *A. turcosus* and overview of related species. *Plant Systematics and Evolution* 303: 787–806.
- Hubka V, Peterson SW, Frisvad JC, et al. 2013. *Aspergillus waksmanii* sp. nov. and *Aspergillus marvanovae* sp. nov., two closely related species in section *Fumigati*. *International Journal of Systematic and Evolutionary Microbiology* 63: 783–789.
- Huelsenbeck JP, Ronquist F. 2001. MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–755.
- Huson DH, Kloepper TH. 2005. Computing recombination networks from binary sequences. *Bioinformatics*, 21: ii159–ii165.
- Hyde KD, Dong Y, Phookamsak R, et al. 2020. Fungal diversity notes 1151–1276: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 100: 5–277.
- Hyde KD, Norphanphoun C, Abreu VP, et al. 2017. Fungal diversity notes 603–708: taxonomic and phylogenetic notes on genera and species. *Fungal Diversity* 87: 1–235.
- Hyde KD, Tennakoon DS, Jeewon R. et al. 2019. Fungal diversity notes 1036–1150: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 96: 1–242.

- Hymphries Z, Seifert KA, Hirooka Y, et al. 2017. A new family and genus in Dothideales for Aureobasidium-like species isolated from house dust. *IMA Fungus* 8: 299–315.
- Irwin JAG. 1991. *Phytophthora macrochlamydospora*, a new species from Australia. *Mycologia* 83: 517–519.
- Jiang H-B, Hyde KD, Yang E-F, et al. 2021. Morphological and phylogenetic appraisal of *Ophioceras* (Ophiocercaceae, Magnaporthales). *PLoS ONE* 16: e0253853.
- Jiang N, Fan X, Tian C, et al. 2020. Reevaluating Cryphonectriaceae and allied families in Diaporthales. *Mycologia* 112: 267–292.
- Johnston PR. 1986. Rhytismataceae in New Zealand 1. Some foliicolous species of *Coccomyces* de Notaris and *Propolis* (Fries) Corda. *New Zealand Journal of Botany* 24: 89–124.
- Johnston PR. 1994. Ascospore sheaths of some *Coccomyces*, *Hypoderma*, and *Lophodermium* species (Rhytismataceae). *Mycotaxon* 52: 221–239.
- Johnston PR, Gamundi IJ. 2000. *Torrendiella* (Ascomycota, Helotiales) on *Nothofagus*. *New Zealand Journal of Botany* 38: 493–513.
- Ju Y-M, Rogers JD. 2001. New and interesting *Biscogniauxia* taxa, with a key to the world species. *Mycological Research* 9: 1123–1133.
- Kalyanamoorthy S, Minh BQ, Wong TFK, et al. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* 14: 587–589.
- Karstedt F, Baroni TJ, Largent DL, et al. 2019. Phylogenetic and morphological analyses of species of the Entolomataceae (Agaricales, Basidiomycota) with cuboid basidiospores. *Phytotaxa* 391: 001–027.
- Katoh K, Rozewicki J, Yamada KD. 2019. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20: 1160–1166.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30: 772–780.
- Khalabuda TV. 1973. Fungi of the genus *Mortierella* Coemans. *Institut mikrobiologii (Akademii nauk SSSR)*.
- Kirk PM. 1982. New or interesting microfungi VI. *Sporidesmiella* gen. nov. (hyphomycetes). *Transactions of the British Mycological Society* 79: 479–489.
- Klaubauf S, Tharreau D, Fournier E, et al. 2014. Resolving the polyphyletic nature of *Pyricularia* (Pyriculariaceae). *Studies in Mycology* 79: 85–120.
- Knapp DG, Kovács GM, Zajta E, et al. 2015. Dark septate endophytic pleosporalean genera from semiarid areas. *Persoonia* 35: 87–100.
- Kobayashi T, Courtecuisse R. 2000. Two new species of *Inocybe*, section *Marginatae* (Agaricales, Cortinariaceae) from Japan. *Mycoscience* 41: 161–166.
- Kornerup A, Wanscher JH. 1967. *Methuen Handbook of Colour*. Methuen & Co Ltd, London, England.
- Kornerup A, Wanscher JH. 1978. *Methuen Handbook of Colour*, 3rd edn. Eyre Methuen, Ltd., London.
- Kornerup A, Wanscher JH. 1983. *Methuen Handbook of Colour*, 3rd edn. Eyre Methuen, Ltd., London.
- Kozlov AM, Darriba D, Flouri T, et al. 2019. RAXML-NG: a fast, scalable, and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35: 4453–4455.
- Kruys A, Huhndorf SM, Miller AN. 2015. Coprophilous contributions to the phylogeny of Lasiosphaeriaceae and allied taxa within Sordariales (Ascomycota, Fungi). *Fungal Diversity* 70: 101–113.
- Kühner R. 1955. Compléments à la "Flore Analytique" VI) *Inocybe* goniosporés et *Inocybe* acystidiés. Espèces nouvelles ou critiques. *Bulletin de la Société Mycologique de France* 71: 279–311.
- Kumar S, Stecher G, Li M, et al. 2016. MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* 35: 1547–1549.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7 for Bigger Datasets. *Molecular Biology and Evolution* 33: 1870–1874.
- Kušan I, Matočec N, Antičić O, et al. 2014. Biogeographical variability and re-description of an imperfectly known species *Hamatocanthoscypha rotundispora* (Helotiales, Hyaloscyphaceae). *Phytotaxa* 170: 1–12.
- Kuyper TW. 1986. A revision of the genus *Inocybe* in Europe. I. Subgenus *Inosperma* and the smooth-spored species of subgenus *Inocybe*. *Persoonia Supplement* 3: 1–247.
- Lanfear R, Frandsen PB, Wright AM, et al. 2017. PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* 34: 772–773.
- Laraba I, McCormick SP, Vaughan MM, et al. 2021. Phylogenetic diversity, trichothecene potential, and pathogenicity within *Fusarium sambucinum* species complex. *Plos One* 16: e0245037.
- Larsson KH. 1994. Poroid species in *Trechispora* and use of calcium oxalate crystals for species identification. *Mycological Research* 98: 1153–1172.
- Latha KD, Raj KA, Manimohan P. 2019. *Laccaria violaceotincta*: a new species from tropical India based on morphology and molecular phylogeny. *Phytotaxa* 392: 140–146.
- Latha KPD, Raj KNA, Thushara C, et al. 2016. Three new species of *Calocybella* from India based on morphology and molecular phylogeny. *Phytotaxa* 255: 133–143.
- Le Gal M, Mangelot MF. 1961. Contribution à l'étude des Mollisioïdées. IV. (3e série). *Revue de Mycologie* 26: 263–331.
- Lechat C, Fournier J. 2016. *Varicosporellopsis*, a new aquatic genus from South of France. *Ascomycete.org* 8: 96–100.
- Leonardi M, Furtado ANM, Comandini O, et al. 2020. *Halimium* as an ectomycorrhizal symbiont: new records and an appreciation of known fungal diversity. *Mycological Progress* 19: 1495–1509.
- Leslie JF, Summerell BA. 2006. *The Fusarium laboratory manual*. Blackwell Publishing, Ames.
- Lévesque CA, De Cock AWAM. 2004. Molecular phylogeny and taxonomy of the genus *Pythium*. *Mycological Research* 108: 1363–1383.
- Li HY, Sun GY, Zhai XR, et al. 2012. *Dissoconiaceae* associated with sooty blotch and flyspeck on fruits in China and the United States. *Persoonia* 28: 113–125.
- Li J-X, He M-Q, Zhao R-L. 2021. Three new species of *Micropsalliota* (Agaricaceae, Agaricales) from China. *Phytotaxa* 491: 174.
- Liberta AE. 1973. The genus *Trechispora* (Basidiomycetes, Corticiaceae). *Canadian Journal of Botany* 51: 1871–1892.
- Lima CS, Pfenning LH, Costa SS, et al. 2019. A new *Fusarium* lineage within the *Gibberella fujikuroi* species complex is the main causal agent of mango malformation disease in Brazil. *Plant Pathology* 58: 33–42.
- Liu LL, Yang J, Liu NG, et al. 2019. *Sporidesmium guizhouense* sp. nov. (Sordariomycetes incertae sedis), a new species from a freshwater habitat in Guizhou Province, China. *Phytotaxa* 422: 144–156.
- Liu ZY, Plautz HL, Lumyong S, et al. 2016. Fungal diversity notes 603–708: taxonomic and phylogenetic notes on genera and species. *Fungal Diversity* 87: 1–235.
- Loizides M, Bellanger J-M, Yiangou Y, et al. 2018. Preliminary phylogenetic investigations into the genus *Amanita* (Agaricales) in Cyprus, with a review of previous records and poisoning incidents. *Documents Mycologiques XXXVII*: 201–218.
- Lombard L, Houbraken J, Decock C, et al. 2016. Generic hyper-diversity in *Stachybotriaceae*. *Persoonia* 36: 156–246.
- Lombard L, Van der Merwe NA, Groenewald JZ, et al. 2015. Generic concepts in *Nectriaceae*. *Studies in Mycology* 80: 189–245.
- Lowe JL. 1966. *Polyporaceae of North America*. The genus *Poria*. New York State University College of Forestry Technical Publication 90.
- Lu YZ, Liu JK, Hyde KD, et al. 2018. A taxonomic reassessment of *Tubeufiales* based on multi-locus phylogeny and morphology. *Fungal Diversity* 92: 131–344.
- Ludwig E. 2007. *Pilzkompendium* (Eching), 2. Berlin. Ed. Fungicon.
- Lundquist N. 1972. *Nordic Sordariaceae s. lat. Symbolae Botanicae Upsalienses* 20: 1–374.
- Luo ZL, Hyde KD, Bhat DJ, et al. 2018. Morphological and molecular taxonomy of novel species *Pleurotheciaceae* from freshwater habitats in Yunnan, China. *Mycological Progress* 17: 511–530.
- Luo ZL, Hyde KD, Liu JK, et al. 2019. Freshwater *Sordariomycetes*. *Fungal Diversity* 99: 451–660.
- Madrid H, da Cunha KC, Gené J, et al. 2014. Novel *Curvularia* species from clinical specimens. *Persoonia* 33: 48–60.
- Manoj KA, Vrinda KB, Pradeep CK. 2018. New and noteworthy crepidotoid agarics from India. *Cryptogamie, Mycologie* 39: 287–298.
- Maryani N, Sandoval-Denis M, Lombard L, et al. 2019. New endemic *Fusarium* species hitch-hiking with pathogenic *Fusarium* strains causing Panama disease in small-holder banana plots in Indonesia. *Persoonia* 43: 48–69.
- Marin-Felix Y, Guarro J, Cano-Lira JF, et al. 2018. *Melanospora* (Sordariomycetes, Ascomycota) and its relatives. *MycKeys* 44: 81–122.
- Mata JL, Hughes KW, Petersen RH. 2006. An investigation of *omphalotaceae* (Fungi: Euagarics) with emphasis on the genus *Gymnopus*. *Sydowia* 58: 191–289.
- Matheny BP, Swenie RA. 2018. The *Inocybe* *geophylla* group in North America: a revision of the lilac species surrounding *I. lilacina*. *Mycologia* 110: 618–634.
- McKenzie EHC, Pinnoi A, Wong MKM, et al. 2002. Two new hyaline *Chalara* species, and a key to species described since 1975. *Fungal Diversity* 11: 129–139.
- Meiras-Otttoni A, Larsson K-H, Gibertoni TB. 2021. Additions to *Trechispora* and the status of *Scytinopogon* (Trechisporales, Basidiomycota). *Mycological Progress* 20: 203–222.

- Menolli Jr N, Asai T, Capelari M. 2010. Records and new species of *Pluteus* from Brazil based on morphological and molecular data. *Mycologia* 1: 130–153.
- Miller AN, Huhndorf SM. 2004a. A natural classification of *Lasiosphaeria* based on nuclear LSU rDNA sequences. *Mycological Research* 108: 26–34.
- Miller AN, Huhndorf SM. 2004b. Using phylogenetic species recognition to delimit species boundaries within *Lasiosphaeria*. *Mycologia* 96: 1106–1127.
- Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Proceedings of the Gateway Computing Environments Workshop (GCE), 14 Nov. 2010, New Orleans, LA: 1–8.
- Minh BQ, H.A. Schmidt, Chernomor O, et al. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution* 37: 1530–1534.
- Minter DW. 1996. *Colpoma juniperi*. IMI descriptions of Fungi and bacteria No 1294. *Mycopathologia* 136: 159–161.
- Misra PC, Srivastava P. 1976. A new species of *Tetracoccosporium*. *Mycotaxon* 4: 276–278.
- Morgan AP. 1890. North American Fungi. The Gasteromycetes. III. The Journal of the Cincinnati Society of Natural History 12: 163–172.
- Morozova OV, Alexandrova AV, Popov ES, et al. 2016. New data on the agaricoid basidiomycetes of the Tver Region. *Novosti Sistematiki Nizshikh Rastenii*. 50: 174–186. [In Russian.]
- Morozova OV, Popov ES, Kovalenko AE. 2012. Studies on mycobiota of Vietnam. I. Genus *Entoloma*: new records and new species. *Mikologiya i fitopatologiya* 46: 184–200.
- Munsell Color 1994. Soil Color Charts (revised edition). Macbeth Division of Kollmorgen Instruments Corporation. New Windsor. New York. USA.
- Muroi T, Udagawa S, Otani Y. 1987. Some coprophilous ascomycetes from Peru. In: H Inoue (ed), *Studies on Cryptogams in Southern Peru*. Tokai University Press, Tokyo: 151–168.
- Nag-Raj TR, Kendrick WB. 1975. Monograph of *Chalara* and allied genera. Wilfrid Laurier University Press, Waterloo, Ontario, Canada.
- Nagy LG, Petkovits T, Kovács GM, et al. 2011. Where is the unseen fungal diversity hidden? A study of *Mortierella* reveals a large contribution of reference collections to the identification of fungal environmental sequences. *New Phytologist* 191: 789e794.
- Nguyen L-T, Schmidt HA, Von Haeseler A, et al. 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32: 268–274.
- Noordeloos ME. 1992. *Entoloma* s.l. Fungi Europaei, vol. 5. Giovanna Biella, Saronno, Italy.
- Noordeloos ME, Morozova OV. 2010. New and noteworthy *Entoloma* species from the Primorsky Territory, Russian Far East. *Mycotaxon* 112: 231–255.
- Nováková A, Hubka V, Dudová Z, et al. 2014. New species in *Aspergillus* section *Fumigati* from reclamation sites in Wyoming (U.S.A.) and revision of *A. viridinutans* complex. *Fungal Diversity* 64: 253–274.
- O'Donnell K, Cigelnik E, Nirenberg H. 1998. Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia* 90: 465–493.
- O'Donnell K, Lutzoni FM, Ward TJ, et al. 2001. Evolutionary relationships among mucoralean fungi (Zygomycota): Evidence for family polyphyly on a large scale. *Mycologia* 93: 286–296.
- O'Donnell K, McCormick SP, Busman M, et al. 2018. Marasas et al. 1984 'Toxigenic *Fusarium* species: identity and mycotoxicology' revisited. *Mycologia* 110: 1058–1080.
- Ou JH, Lin GC, Chen CY. 2020. *Sarocladium* species associated with rice in Taiwan. *Mycological Progress* 19: 67–80.
- Peterson SW, Manitchotpisit P, Leathers TD. 2013. *Aureobasidium thailandense* sp. nov. isolated from leaves and wooden surfaces. *International Journal of Systematic and Evolutionary Microbiology* 63: 790–795.
- Pitt JI, Lantz H, Pettersson OV, et al. 2013. *Xerochrysium* gen. nov. and *Bettsia*, genera encompassing xerophilic species of *Chrysosporium*. *IMA Fungus* 4: 229–241.
- Rajchenberg M. 1987. Type studies of Polyporaceae (Aphylliphorales) described by J. Rick. *Nordic Journal of Botany* 7: 553–568.
- Rajeshkumar KC, Braun U, Groenewald JZ, et al. 2021. Phylogenetic placement and reassessment of *Asperisporium pongamiae* as *Pedrocrousiella pongamiae* gen. et comb. nov. (Mycosphaerellaceae). *Fungal Systematics and Evolution* 7: 165–176.
- Rayner RW. 1970. A Mycological Colour Chart. Commonwealth Mycological Institute, Kew and British Mycological Society.
- Réblová M, Untereiner WA, Réblová K. 2013. Novel evolutionary lineages revealed in the Chaetothyriales (Fungi) based on multigene phylogenetic analyses and comparison of ITS secondary structure. *PLoS ONE* 8: e63547.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Ronquist F, Teslenko M, Van der Mark P, et al. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542.
- Rossman AY. 2000. Towards monophyletic genera in the holomorphic Hypocreales. *Studies in Mycology* 45: 27–34.
- Rossman AY, Samuels GJ, Rogerson CT, et al. 1999. Genera of Bionectriaceae, Hypocreaceae and Nectriaceae (Hypocreales, Ascomycetes). *Studies in Mycology* 42: 1–248.
- Ryberg M, Nilsson RH, Kristiansson E, et al. 2008. Mining metadata from unidentified ITS sequences in GenBank: a case study in *Inocybe* (Basidiomycota). *BMC Evolutionary Biology* 8: 50.
- Saccardo PA. 1889. *Sylloge fungorum omnium hucusque cognitorum* VIII. Saccardo, Padova.
- Schipper MAA. 1978. On the genera *Rhizomucor* and *Parasitella*. *Studies in Mycology* 17: 52–71.
- Schultes NP, Murtishi B, Li DW. 2017. Phylogenetic relationships of *Chlamydomyces*, *Harzia*, *Olpitrichum*, and their sexual allies, *Melanospora* and *Sphaerodes*. *Fungal Biology* 121 (10): 890–904.
- Seifert K, Morgan-Jones G, Gams W, et al. 2011. The Genera of Hyphomycetes. CBS Biodiversity Series no. 9: 1–997. CBS-KNAW Fungal Biodiversity Centre, Utrecht, Netherlands.
- Shang QJ, Hyde KD, Camporesi E, et al. 2020. Additions to the genus *Cytospora* with sexual morph in Cytosporaceae. *Mycosphere* 11: 189–224.
- Shearer CA, Crane JL, Chen W. 1999. Freshwater ascomycetes: *Ophioceras* species. *Mycologia* 91: 145–156.
- Sherwood MA. 1980. Taxonomic studies in the Phacidiales: The genus *Coccomyces* (Rhytismataceae). *Occasional Papers of the Farlow Herbarium of Cryptogamic Botany* 15: 1–120.
- Silvestro D, Michalak I. 2012. raxmlGUI: a graphical frontend for RAXML. *Organisms Diversity and Evolution* 12: 335–337.
- Singer R. 1947. Contributions towards a monograph of the genus *Crepidotus*. *Lilloa* 13: 59–95.
- Spotts RA, Seifert KA, Wallis KM, et al. 2009. Description of *Cryptosporopsis kienholzii* and species profiles of *Neofabraea* in major pome fruit growing districts in the Pacific Northwest USA. *Mycological Research* 113: 1301–1311.
- Stadler T. 2009. On incomplete sampling under birth-death models and connections to the sampling-based coalescent. *Journal of Theoretical Biology* 261: 58–66.
- Stajsic V. 2020a. In VICFLORA: Flora of Victoria, *Oxalis* <https://vicflora.rbg.vic.gov.au/flora/taxon/0ed554c2-95c9-4cb2-b85e-da75e2655077> [accessed 6 April 2021].
- Stajsic V. 2020b. Oxalidaceae. In: Kodala PG. (ed), *Flora of Australia*, Australian Biological Resources Study, Canberra. <https://profiles.ala.org.au/opus/foa/profile/Oxalidaceae> [accessed: 6 April 2021].
- Stamatidakis A. 2014. RAXML version 8: A toll for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313.
- Summerbell RC, Gueidan C, Guarro J, et al. 2018. The Protean *Acremonium*. *A. sclerotigenum/egyptiacum*: Revision, Food Contaminant, and Human Disease. *Microorganisms* 6: 88.
- Summerbell RC, Gueidan C, Schroers HJ, et al. 2011. *Acremonium* phylogenetic overview and revision of *Gliomastix*, *Trichothecium* and *Sarocladium*. *Studies in Mycology* 68: 139–162.
- Swofford DL. 2003. PAUP* 4.0b10. Phylogenetic Analysis Using Parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland, MA, USA.
- Tamura K, Stecher G, Peterson D et al. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30: 2725–2729.
- Tanney JB, Seifert KA. 2020. Mollisiaceae: An overlooked lineage of diverse endophytes. *Studies in Mycology* 95: 293–380.
- Trifinopoulos J, Nguyen L-T, von Haeseler A, et al. 2016. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research*: 44: W232–W235.
- Tulloss RE. 1994. Type studies in *Amanita* section *Vaginatae* I: some taxa described in this century (studies 1–23) with notes on description of spores and refractive hyphae in *Amanita*. *Mycotaxon* 52: 305–396.
- Valenzuela-Lopez N, Cano-Lira JF, Guarro J, et al. 2018. Coelomycetous Dothideomycetes with emphasis on the families Cucurbitariaceae and Didymellaceae. *Studies in Mycology* 90: 1–69.
- Vandepol N, Liber J, Desiro A, et al. 2020. Resolving the Mortierellaceae phylogeny through synthesis of multi-gene phylogenetics and phylogenomics. *Fungal Diversity* 104: 267–289.
- Vánky K. 2011 '2012'. *Smut fungi of the world*. St. Paul, Minnesota, APS Press.
- Vánky K, Lutz M, Bauer R. 2008. About the genus *Thecaphora* (Glomosporiaceae) and its new synonyms. *Mycological Progress* 7: 31–39.

- Vauras J. 1994. *Inocybe diabolica*, a new agaric from subarctic regions. *Mycologia Helvetica* 2: 121–128.
- Vauras J, Kokkonen K. 2009. Finnish records of the genus *Inocybe*. The new species *Inocybe saliceticola*. *Karstenia* 48: 57–67.
- Vauras J, Larsson E. 2016. *Inocybe caprimulgi* and *I. lacunarum*, two new nodulose-spored species from Fennoscandia. *Karstenia* 55: 1–18.
- Vellinga EC. 1988. Glossary. In: Bas C, Kuyper TW, Noordeloos ME, Vellinga EC (eds). *Flora Agaricina Neerlandica* vol. 1: 54–64. Blakema, Rotterdam, The Netherlands.
- Vellinga EC. 1990. *Pluteus*. In: Bas C, Kuyper TW, Noordeloos ME, Vellinga EC (eds), *Flora Agaricina Neerlandica*, vol 2: 31–55. Blakema, Rotterdam, The Netherlands.
- Vellinga EC. 2001. *Lepiota*. In: Noordeloos ME, Kuyper ThW, Vellinga EC. (eds), *Flora Agaricina Neerlandica* 5: 109–151. Balkema publishers.
- Vellinga EC. 2010. *Lepiota* in California: species with a hymeniform pileus covering. *Mycologia* 102: 664–674.
- Viana FMP, Sousa JA, Araújo JDM, et al. 2012. Fungos Associados a Algumas Espécies Florestais Madeireiras no Distrito de Irrigação de Acaraú-Marco, CE. Documentos / Embrapa Agroindústria Tropical.
- Videira SIR, Groenewald JZ, Braun U, et al. 2016. All that glitters is not *Ramularia*. *Studies in Mycology* 83: 49–163.
- Videira SIR, Groenewald JZ, Nakashima C, et al. 2017. *Mycosphaerellaceae* – Chaos or clarity? *Studies in Mycology* 87: 257–421.
- Vilgalys R, Miller Jr OK. 1983. Biological species in the *Collybia dryophila* group in North America. *Mycologia* 75: 707–722.
- Visagie CM, Houbraken J. 2020. Updating the taxonomy of *Aspergillus* in South Africa. *Studies in Mycology* 95: 253–292.
- Vitelli M, Vessella F, Cardoni S, et al. 2017. Phylogeographic structuring of plastome diversity in Mediterranean oaks (*Quercus* Group *Ilex*, Fagaceae). *Tree Genetics & Genomes* 13.3.
- Vittadini C. 1831. *Monographia Tuberacearum*. Typographia Rusconi, Milano, Italy.
- Vizzini A, Angelini C, Ercole E. 2017. Is the species diversity in the lyophylloid genera *Calocybella* and *Gerhardtia* (Agaricales, Basidiomycota) underestimated? Two new species from the Dominican Republic. *Phytotaxa* 291: 241–252.
- Vizzini A, Consiglio G, Setti L, et al. 2015. *Calocybella*, a new genus for *Rugosomyces pudicus* (Agaricales, Lyophyllaceae) and emendation of the genus *Gerhardtia*. *IMA Fungus* 6: 1–11.
- Vizzini A, Liang JF, Jančovičová S, et al. 2014. *Lepiota coloratipes*, a new species for *Lepiota rufipes* ss. *Auct. europ. non ss. orig.* *Mycological Progress* 13: 171–179.
- Vizzini A, Tatti A, Huijser HA, et al. 2019. Looking for *Lepiota psalion* Huijser & Vellinga (Agaricales, Agaricaceae). *MycKeys* 52: 45–69.
- Vizzini A, Zotti M, Traverso M, et al. 2016. Variability, host range, delimitation and neotypification of *Amanita simulans* (*Amanita* section *Vaginatae*): collections associated with *Helianthemum* grasslands, and epitypification of *A. lividopallescens*. *Phytotaxa* 280: 1–22.
- Voitk A, Saar I, Lodge DJ, et al. 2020a. From Esteri's to Bond's mushroom — some new and old *Cuphophylli* in NL. *Omphalina* 11: 71–80.
- Voitk A, Saar I, Lodge DJ, et al. 2020b. New species and reports of *Cuphophyllus* from northern North America compared with related Eurasian species. *Mycologia* 112: 438–452.
- Vu D, Groenewald M, Szöke S et al. 2016. DNA barcoding analysis of more than 9 000 yeast isolates contributes to quantitative thresholds for yeast species and genera delimitation. *Studies in Mycology* 85: 91–105.
- Wanasinghe DN, Jones EBG, Camporesi E, et al. 2015. The genus *Murispora*. *Cryptogamie Mycologie* 36: 419–448.
- Wang H, Gu ZR, Zhao CL. 2021. *Hyphodermella zixishanensis* (Polyporales, Basidiomycota), a new species with reddish hymenial surface. *Nordic Journal of Botany* 39: e03329.
- Wu WP, Zhuang WY. 2005. *Sporidesmium*, *Endophragmiella* and related genera from China. *Fungal Diversity Research Series* 15: 1–351.
- Yilmaz N, Sandoval-Denis M, Lombard L, et al. 2021. Redefining species limits in the *Fusarium fujikuroi* species complex. *Persoonia* 46: 129–162.
- Yuan H-S, Dai Y-C. 2012. Wood-inhabiting fungi in southern China 5. Polypores from Guangxi Autonomous Region. *Annales Botanici Fennici* 49: 341–351.
- Yurkov A, Guerreiro MA, Sharma L et al. 2015. Multigene assessment of the species boundaries and sexual status of the basidiomycetous yeasts *Cryptococcus flavesens* and *C. terrestris* (Tremellales). *PLoS One* 10: e0120400.
- Zalar P, Gostinčar C, De Hoog GS, et al. 2008. Redefinition of *Aureobasidium pullulans* and its varieties. *Studies in Mycology* 61: 21–38.
- Zeller SM, Smith AH. 1964. The genus *Calvatia* in North America. *Lloydia* 27: 148–186.
- Zhao R-L, Desjardin DE, Soyong K, et al. 2010. A monograph of *Microp-salliota* in northern Thailand based on morphological and molecular data. *Fungal Diversity* 45: 33–79.